

ИЗВЕСТИЯ АКАДЕМИИ НАУК СССР СЕРИЯ ГЕОЛОГИЧЕСКАЯ

IZVESTIYA AKAD. NAUK SSSR

SERIYA GEOLOGICHESKAYA

961

CONTENTS

No. 9, September

<p>ETROCHEMICAL CRITERIA ESTABLISHING THE RELATION BETWEEN MINERALIZATION AND GRANITOIDS (AS EXEMPLIFIED IN THE MYAO-CHANSKIY REGION), by M.G. Rub, V.V. Onikhimovskiy, and B.V. Makeyev</p>	1
<p>MAGNETIC-FRACTIONAL-MINERALOGICAL STUDIES OF ROCKS, by F.N. Yefimov</p>	21
<p>THE TECTONICS OF THE NORTHERN PARTS OF PATOMSK UPLAND, by S.V. Ruzhentsev, and Chang Bu-Chung</p>	32
<p>AGE AND STRATIFICATION SEQUENCE OF DEPOSITS IN THE UPPER PART OF THE KARATAU SERIES IN THE SOUTHERN URALS, by Yu.R. Bekker</p>	43
<p>OUTLINE OF THE STRATIGRAPHY AND TECTONICS OF THE TAS-KHAYAKHTAKH RANGE, by N.A. Bogdanov</p>	54
<p>THE FACIES, DYNAMIC PHASES, AND FORMATIONS OF ALLUVIUM, by I.P. Kartashov</p>	67
<p>THE PROBLEM OF THE RELATIONSHIPS BETWEEN MARINE AND CONTINENTAL DEPOSITS IN THE LOWER AND MIDDLE VOLGA REGIONS, by Yu.M. Vasil'yev and P.V. Fedorov</p>	79
METHODS	
<p>SEPARATION OF CLAY PARTICLES BY ELECTROPHORESIS, by N.V. Logvinenko, and A.A. Lazarenko</p>	87
<p>THE 1961 LAUREATES OF THE LENIN PRIZES, N.M. Strakhov — Originator of the Theory of Lithogenesis</p>	91
REVIEWS AND DISCUSSIONS	
<p>THE PROBLEM OF THE NOMENCLATURE OF EFFUSIVE ROCKS, by T.Ya. Goncharova</p>	93
<p>THE NOMENCLATURE OF EFFUSIVE ROCKS, by I.M. Speranskaya</p>	94
<p>MAIN RESULTS OF THE DISCUSSION ON THE NOMENCLATURE OF EFFUSIVE ROCKS, by Ye.K. Ustiyev</p>	96

ON A BOOK BY G. M. ZARIDZE AND N. F. TATRISHVILI
 "MAGMATISM IN GEORGIA AND ASSOCIATED
 ORE FORMATIONS", by A. P. Lebedev 102

BIBLIOGRAPHY 105

EDITORIAL COLLEGIUM

G. D. Afanas'yev (Chief Editor), Corresponding Member of the ASUSSR
 G. P. Barsanov, Doctor of Geological and Mineralogical Sciences
 K. A. Vlasov, Corresponding Member of the ASUSSR
 D. S. Korzhinskiy, Academician
 M. F. Mirchink, Corresponding Member of the ASUSSR
 Ye. V. Pavlovskiy, Doctor of Geological and Mineralogical Sciences
 A. V. Peyve, Corresponding Member of the ASUSSR
 V. I. Smirnov, Corresponding Member of the ASUSSR
 F. V. Chukhrov, Corresponding Member of the ASUSSR
 D. I. Shcherbakov, Academician
 V. S. Yablokov, Candidate for Geological and Mineralogical Sciences
 (Secretary of the Editorial Collegium)

TWENTY-SIXTH YEAR OF PUBLICATION

Technical Editor - V. Ya. Karasev

T-10220. Approved for printing August 24, 1961.

Circulation - 3, 250 copies. Order 3902

Paper size 70 x 108-1/16

Paper 4

Printing sheets 10.96 x 1

Publ. sheets 12.7

Second printing office of the USSR Academy of Sciences Publishing House.
 Moscow, Shubinskiy per. 10.

PETROCHEMICAL CRITERIA ESTABLISHING THE RELATION BETWEEN MINERALIZATION AND GRANITOIDS (AS EXEMPLIFIED IN THE MYAO CHANSKIY REGION)¹

by

M. G. Rub, V. V. Onikhimovskiy, and B. V. Makeyev

Studies of the geochemical characteristics of the Upper-Cretaceous intrusions and the accompanying post-magmatic formations in the Myao-Chanskiy region (Khabarovsk Krai) produced some interesting new material which confirms the opinion we expressed in 1956 regarding the considerable importance of the petrochemical criteria associated with the relationship between mineralization and intrusions. In addition, these investigations confirmed the existence of specialized intrusions and showed that intrusive bodies emplaced at different times (the tin-bearing variety, in particular) and located in different structural zones may — along with other common features — also possess certain specific individual peculiarities. Our new material is partially described in this article.

Brief Geological Outline of the Myao-Chanskiy Region

According to I. Ya. Zinter, V. V. Onikhimovskiy, P. A. Epov, and other investigators, the mineralized zone in the Myao-Chanskiy region is confined to the zone of Cretaceous folding, composed of Jurassic rocks and unconformably overlain by Lower-Cretaceous formations, this zone is covered by Upper Cretaceous, Cenozoic, and Early Quaternary effusives.

The principal structural feature in this region is a major anticline (the Badzhalskiy anticline) which strikes in a north easterly direction. The center of the geanticline consists of Jurassic rocks represented by sandstones and siltstones with intercalated cherty clay shales and argillaceous slate, with occasional gravels and fine pebble conglomerates. The total thickness of the Jurassic formations is 3500 m. They are overlain by Lower Cretaceous sandstones, siltstones, clay shales and conglomerates containing beds of quartz porphyries, their tuffs, gritstones, tufogenic sandstones and tuffites containing fossilized Lower Cretaceous flora. According to the

data of P. A. Epov, V. V. Onikhimovskiy and other investigators, the Lower Cretaceous formations attain a thickness of 4000 meters.

The Upper Cretaceous deposits, which are well developed in the investigated region (Fig. 1), are represented by a variety of porphyrites and their tuffs comprising the Amutskaya series. The observed thickness of the Upper Cretaceous formations, whose ages were determined from fossil flora is 550 meters.

In addition to these Jurassic and Cretaceous formations, there are extensive Tertiary and Quaternary rocks in the Myao-Chanskiy region of these, only the basalts and dolerites are extensive.


The data of V. V. Onikhimovskiy and others indicate that the structure of the Myao-Chanskiy region is the result of Variscian, Cretaceous, and Cenozoic tectonic stresses. Therefore, the history of its development is long and complex.

Brief Geologic, Petrographic, and Geochemical Characterization of the Intrusive Rocks

Intrusives are widespread in the Myao-Chanskiy region and consist of gabbro-diorites, quartz-diorites, diorite-monzonites, monzonites, granodiorites, various granites, and other varieties. In addition, this area abounds in dikes of acid (fine-grained granites, aplites, granite-porphyrines, quartz-porphyrines, felsites), average (granodiorite-porphyrines, monzonite-porphyrines, diorite-porphyrines) and basic (diabasic porphyries, gabbro-diabases) varieties.

All of these intrusives cut the Jurassic and Cretaceous sedimentary deposits as well as the Lower and Upper Cretaceous effusive formations. Thus, the lower age limit of these intrusives is clearly Upper Cretaceous. Their upper age limit has not been precisely determined. The only known fact is that these rocks are overlain by Neogene dolerites and basalts. An absolute age determination of these intrusives made at the All-Union Geological Scientific-

¹ Petrogeokhimicheskiye kriterii svyazi orudeniya s granitoidami (na primere Myao-Chanskogo rayona). pp. 3 - 23.



A - B Section




FIGURE 1. Schematic geologic map of the Myao-Chanskiy region compiled by V.V. Onikhimovskiy, et al.

Quaternary System: 1 - alluvial deposits; 2 - basalts, dolerites, and their tuffs. Tertiary System: 3 - dolerites, basalts and their tuffs; pebble-beds, clays, sandy loams, lignites. Cretaceous System: 4 - porphyrites and their tuffs; 5 - conglomerates with beds of quartz-porphyrines. Jurassic System: 6 - clay shales, siltstones with interbedded sandstones; 7 - fine- and medium-grained sandstones with intercalations of clay and cherty shales; Myao-Chanskiy intrusive complex granitoids; 8 - non-articulated granitoids of Myao-Chanskiy

Explanation for FIGURE 1 continued on next page.

rusive complex; 9 - porphyritic biotite and alaskite granites, the third phase of the
 o-Chanskiy intrusive complex. Upper Cretaceous granitoids of the Myao-Chanskiy intrusive
 plex; 10 - granitoids, quartz-diorites, diorites, facies of endomorphic granites of the
 ond phase; 11 - biotite and biotite-hornblende granites, second phase of the Myao-Chanskiy
 rusive complex; 12 - gabbro-diorites, diorites, diorite-monzonites, monzonites, quartz-
 zonites, granodiorites and other varieties, the first phase of the Myao-Chanskiy intrusive
 plex. Dikes of the Upper Cretaceous intrusive complex; 13 - Dikes of acid, average, and
 ic composition; 14 - Fracture zones containing quartz-tourmaline mineralization.





FIGURE 2. Stratigraphic Column

1 - diabase porphyrites; 2 - diorite porphyrites, diorite-monzonite-porphyries, granodiorite-porphyries; 3 - porphyritic, biotite, alaskite tourmaline-bearing granites, third phase of the Myao-Chanskiy intrusive complex, 4 - granodiorites, quartz-diorites, granodiorite-porphyries, marginal phase of the second-phase granites; 5 - biotite and biotite-hornblende granites, second phase of the Myao-Chanskiy intrusive complex; 6 - porphyritic diorite-monzonites, diorite-monzonite-porphyries, diorite porphyrites, marginal phase of the first-phase hybrid rocks; 7 - diorite-monzonites, quartz-diorites, and other varieties, the first phase of the Myao-Chanskiy intrusive complex.

Research Institute (VSEGEI) by N.I. Poleyeva using the argon method produced a figure of 80 to 100 million years which corresponds to the Upper Cretaceous. In speaking of the age and of the correlation of the described rocks, one should mention the fact that the views of earlier investigators diverge considerably in this respect. I. Ya. Zinter, E. P. Izokh, O. N. Kabakov, and P. A. Epov, for example, believe that the formation of the intrusives in the Myao-Chanskiy region occurred in several different age-phases. The first to intrude were the rocks of basic and average composition, and the acid rocks were intruded later. Moreover, some authors (P. A. Epov) differentiate two phases which, in their opinion, should be related to the Upper Cretaceous intrusive complex. E. P. Izokh distinguishes six phases in the formative process of the granitoids in the Myao-Chanskiy intrusive complex. He considers the gabbro and gabbro-diorites as belonging to the first phase, the monzonites and monzonite-granodiorites - to the second phase, the monzonite-granites to the third. The fourth, fifth, and sixth phases are represented by various granites.

Another group of investigators attributes the diversity of the intrusives in this region to facies varieties of a large, deep-seated granitoid mass. They assign these granitoids to the Upper Cretaceous.

Our special study of the intrusives and related postmagmatic formations permits us to make the following assumption. The formation of the various intrusives in the Myao-Chanskiy region occurred in three successive phases related to a single intrusive complex known as the Myao-Chanskiy.

The first phase is represented by gabbro-diorites, diorites, quartz-diorites, diorite-monzonites, quartz-monzonites and other varieties. The second phase includes biotite and biotite-hornblende granites, granodiorites, and quartz-diorites. The third phase consists of alaskitic and porphyritic-biotite granites (Figure 2). Each phase is accompanied by its dike series.

The diorite porphyry dikes are the youngest manifestations of magmatic activity completing the formation of the Myao-Chanskiy intrusive complex.²

The first phase intrusives are widely distributed in the basin of Silinka and Kholdomi

Rivers where they form a number of massifs covering an area of from 1-2 to 30-40 square kilometers. They are accompanied by their associated dike series represented by diorite-porphyrries, diorite-monzonite-porphyrries, monzonite-porphyrries, granodiorite-porphyrries and other varieties.

An intersection of first-phase hybrid rocks and second-phase granites was observed on the right bank of Silinka River by M. G. Rub and B. V. Makeyev. Potassic metasomatism is observed in the hybrid rocks at the contact with the granites. Moreover, in a number of sections near the granites the hybrid rocks are intruded by dikes of fine-grained granites, aplites, and granite-porphyrries which M. G. Rub and B. V. Makeyev classify as derivatives of the second-phase granites.

Detailed petrographic analysis revealed the following as characteristic of the intrusive rocks belonging to the first phase of the Myao-Chanskiy complex: uneven distribution of dark-colored minerals, a rapid change of structures, an unbalanced quantitative and mineralogical composition, associations of minerals unusual for normal magmatic rocks (along with 40% of zonal plagioclase represented by labradorite, the presence of orthoclase - 30 percent and 10-15 percent quartz), and other indications of their hybrid origin. Examination of thin sections and artificial concentrates revealed that the predominant accessory mineral in the intrusives belonging to the first phase of the Myao-Chanskiy complex are: zircon, apatite, pyrite, cassiterite, tourmaline, magnetite, garnet, orthite, and ilmenite.

The chemical analyses of the rocks in question (Table 1, and Figure 3) do not correspond to the average types of rocks, as defined by Daly. In particular, samples c443 and c188 occupy an intermediate position between diorites and monzonites, and we have called these rocks quartzitic diorite-monzonites. The rocks differ from monzonites in their high content of SiO_2 and ferrous iron, and in the small amount of alkaline and ferric iron. They differ from diorites in their high percentages of SiO_2 , K_2O and smaller quantity of Na_2O , and, in addition, in the presence of minor amounts of Li_2O , Rb_2O and B - the characteristic accessory minerals for the granitoids of all three phases of the complex. Numerous spectral analyses bear evident to the fact that the usual secondary elements of the described rocks are: barium, strontium, lithium, rubidium, zirconium, boron, lead, copper, chromium, nickel, and cobalt. Thus, the hybrid rocks referred to above contain, on the one hand, a number of elements characteristic for granites belonging to the later phases of this complex (barium, strontium, zirconium, boron, and silicon), and on the other hand, those typical for basic rocks (phosphorus, chromium, nickel and cobalt).

²V. V. Onikhimovskiy believes that it is possible to differentiate three interrelated groups among other intrusive rocks. Of these, the first and the second (which correspond to our first and second phases) are usually spatially separated, whereas the third represents the final and residual derivatives of granites.




FIGURE 3. Chemical composition of the granitoids of the Myao-Chanskiy intrusive complex.

443, 188 - diorite-monzonite, first phase; 511, 11, 13, 14 - biotite granites, second phase; 489 - alaskite granites, third phase; 481 - tourmaline granites; 12, 16 - granodiorites, facies variety of the second phase; 116 - average composition of alaskites, after Daly; 10 - average composition of alkaline granites, after Daly; 30 - average composition of monzonites, after Daly; 47 - average composition of diorites, after Daly; 45 - average composition of granodiorites, after Daly; 3,4 - average composition of granites, after Daly.

Our studies of the composition of the intrusive rocks constituting the first phase of the Krasnoyarsk-Chanskii complex (up to 500 m deep) showed that oxidation does not increase with depth. This fact is yet another proof contradicting the view-point of certain geologists who tend to explain the great diversity of the granites in the investigated area as facies varieties of the large granitoid mass buried at depth.³

All the aforesaid justifies the supposition that the first-phase intrusives of the Myo-ranskiy complex were formed as a result of cationic hybridization (in early structural stage) of the limestones, porphyrites, and other rocks rich in magnesium and iron by granitic magma.

Analysis of the relationships between the st-phase intrusives and the sandy-shale entry rocks (basin of the Silinka River, the

Silinskiy and Kholdominskiy areas) showed that a certain modification in the composition of the intrusive rocks is observed at their contact with the country rocks. This change is less noticeable away from the contact and is no longer traceable 30 to 50 meters away from the contact. These changes attest to the fact that assimilation and hybridization must have taken place in the upper structural stage when the first-phase intrusives of the complex were being intruded. These processes must have been far more intensive during the formation of the second-phase granites. The first-phase intrusives were subjected to silification, sericitization, and tourmalinization. Moreover, in a number of areas these processes were so intensively developed that the rocks were transformed into quartz-sericite and quartz-tourmaline formations. The second-phase intrusives are widely distributed throughout the Chalba River basin, where they form the great Chalba massif (covering an area of about 180 square kilometers) and a number of smaller massifs. They may also be found in the basin of the Silinka River and in other localities.

According to this point of view, gradual oxidation
ould occur and normal granites should be found
epth.

Table 1

Results of Chemical Analyses of Granitoid Samples of the Myao-Chanskiy Intrusive Complex from M. G. Rub's Materials.
The analyses were made in the Chemical Laboratory of the Institute of Ore Geology of the USSR Academy of Science (IGEM).
The data (in %) are taken from P. A. Epov's report.

Elements	First Phase										Second Phase																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																															
	Quartzite diorite-monzonites					Diorite-monzonite porphyries					Granite-monzonite porphyries					Quartz diorites					Coarse-grained biotite granites					Hybrid rocks of the endomorphic facies																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																
	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample	Sample

¹Samples 188, 443, 511, 481 were analyzed in the Chemical Laboratory of the IGEM of the USSR Academy of Sciences by I. B. Nikitina; Sample c492 - by Ye. Kuznetsova; Samples 148, 148, 2, 301, 447, 560, 87, 1001, 515, 519, 490, 9, 462, 489, 504, 4, 75, and 482 - by G. Ye. Kalenchuk and I. F. Kolosovaya; L. Krutetskaya; Samples 11, 12, 13, 15, are taken from P. A. Epov's report (1957). Analyses for boron were made in the IGEM laboratory by V. V. Danilova.

NOTE: Comma represents decimal point.

Table 1 continued

Elements	Third phase									
	Fine-grained biotite granites					Tourmaline-granites				
	Sample c489	Sample c489	Sample c504	Sample c475	Sample c481	Sample c482	Average composition of monzonites after Daly	Average composition of diorites, after Daly	Average composition of post-cambrian granites after Daly	Average composition of alkaline granites, after Daly
SiO ₂	76.26	—	—	—	75.28	—	56.12	58.90	70.18	73.80
TiO ₂	0.01	—	—	—	0.01	—	1.10	0.76	0.39	0.11
Al ₂ O ₃	12.45	—	—	—	12.02	—	16.96	16.47	14.47	12.33
Fe ₂ O ₃	0.07	—	—	—	0.63	—	2.93	2.89	1.57	2.58
FeO	4.29	—	—	—	1.14	—	4.01	4.04	1.78	1.28
MnO	0.02	—	—	—	0.02	—	0.16	0.12	0.12	0.02
MgO	0.06	—	—	—	0.14	—	3.27	9.57	0.88	0.26
CaO	0.96	—	—	—	0.72	—	6.50	6.14	1.99	0.46
Na ₂ O	3.43	3.52	2.54	3.12	2.22	2.46	3.67	3.46	3.48	4.55
K ₂ O	4.98	5.28	5.94	5.46	5.65	5.60	3.76	2.41	4.11	4.20
H ₂ O ⁻	0.20	—	—	—	0.18	—	—	—	—	—
H ₂ O ⁺	0.15	—	—	—	1.50	—	1.05	1.27	0.84	0.86
CO ₂	0.08	—	—	—	0.35	—	0.47	0.27	0.19	0.05
P ₂ O ₅	0.03	—	—	—	—	—	—	—	—	—
Li ₂ O	0.02	0.04	0.028	0.04	0.03	0.045	—	—	—	—
Rb ₂ O	0.06	0.09	0.037	0.07	0.05	0.07	—	—	—	—
Cs ₂ O	0.019	0.01	0.005	0.01	0.006	0.02	—	—	—	—
B ₂ O ₃	0.01	Not analyzed	Not analyzed	Not analyzed	0.60	Not analyzed	—	—	—	—
	No out-crop	No out-crop	No out-crop	No out-crop	No out-crop	No out-crop	—	—	—	—
Total	100.08	—	—	—	100.54	—	—	—	—	—

NOTE: Comma represents decimal point.

Table 2

Results of Incomplete Chemical Analyses of Orthoclase Feldspars and of Biotite Granites in the Myao-Chanskiy intrusive complex (in weight %).

Oxides	Sample c1001 ^k	Sample c511 ^k	Sample c489 ^k	Sample 511	Sample c511 ^b	Sample c489 ^b	Sample 489
Na ₂ O	3,32	3,09	3,78	0,20	0,20	0,46	0,25
K ₂ O	11,78	12,32	11,73	8,49	8,80	8,16	8,39
Li ₂ O	0,27	0,025	0,035	0,17	0,192	0,244	0,32
Rb ₂ O	0,085	0,073	0,091	0,19	0,182	0,318	0,28
CS ₂ O	0,01	0,008	0,01	0,06	0,051	0,10	0,06
F	Not analyzed	Not analyzed	Not analyzed	0,95	Not analyzed	Not analyzed	0,75

NOTE; Comma represents decimal point.

The second-phase intrusives are represented by coarsegrained biotite and biotite-hornblende granites which, in the peripheral parts of the massifs, become granodiorites, quartz-diorites, and diorite-monzonites.

The granitoids of this phase cut and metamorphose the Jurassic deposits of sandy shales and the Upper Cretaceous porphyrites (Pereval'-nyy section). At the contact with granites the sandstones and shales are metamorphosed into hornfels and are sometimes slightly greisenized. Tourmaline is found in them in small quantities. Porphyrites along the granite contact are also silicified and sometimes slightly tourmalinized. The thickness of contact aureoles around the massifs produced by second-phase granites ranges from tens to hundreds of meters depending on the character of the contacts. Microscopic examinations have disclosed that the second-phase granites are composed of orthoclase feldspar, quartz, plagioclase (represented by oligoclase), biotite, and hornblende. The auxiliary minerals include zircon, apatite, orthite, monazite, rutile, sphene, garnet, magnetite, cassiterite, and tourmaline. The rock texture is hypidiomorphic-granular, and in some sections, micropegmatitic. Sometimes porphyritic texture may also be observed.

Orthoclase feldspar always predominates over plagioclase in granites. Its contents varies from 30 to 50-55 percent and usually amounts to 40-45 percent. The results of chemical analyses of the orthoclase for K₂O, Na₂O, Li₂O, Rb₂O, and CS₂O in the discussed granites (samples 1001, c511^k) are shown in Table 2.

It may be seen from Table 2 that lithium, rubidium, and cesium are always present in orthoclase feldspars. The Na₂O content in them (from 3.09 to 3.32 percent) is attributable

to the large number of perthite inclusions in the albite as shown by petrographic examinations.

The results of spectral analyses of the orthoclase feldspars in the described granites revealed that, apart from rubidium, the characteristic secondary elements of these minerals are: barium (0.0n, rarely 0.n%), rubidium (0.0n, rarely 0.n%), gallium (0.00n to 0.0n%), and copper (from 0.000n to 0.00n%). Moreover, both in the orthoclase feldspars and, particularly, in the plagioclases, boron is invariably present, while strontium (0.0n%) and tin (0.00n to 0.0n%) also frequently occur.

Of the dark-colored minerals, biotite predominates in the discussed granites. The incomplete analysis results of the biotite (sample c511^b) are given in Table 2. Always present in it are Li₂O, Rb₂O, and CS₂O, and to a greater extent, in the biotite of third-phase granites.

The results of semi-quantitative spectral analyses of the biotite reveal that rubidium, lithium, barium, zirconium, zinc, gallium are usually present in quantities of 0.0n% with tin and copper amounting to 0.00n%. In certain cases yttrium was found in 0.00n-0.n% niobium in 0.00n% and lead in 0.00n-0.0n% values. Scandium was present in isolated samples.

Of the ferrides, apart from titanium, manganese, and vanadium, small quantities of chromium and nickel are usually present in negligible amounts in the biotite.

The characteristic accessory elements for biotites in the second-phase granites of the complex are, therefore: rubidium, chromium

ium, lithium, zirconium, tin,⁴ gallium, and copper. It is interesting that beryllium is usually absent, and in those cases when it occurs, its contents never exceeded 0.00n%. The leading accessory minerals of the second-phase granites are zircon, apatite, orthite, cassiterite, pyrite, tourmaline, magnetite, molybdenite, whereas the principal accessory elements are barium, strontium, lithium, niobium, chromium, zirconium, boron, gallium, lead, zinc, copper, and molybdenum.

An investigation of the exomorphic and endomorphic zones of the massifs composed of second-phase granites disclosed that the granites themselves are also altered at their contacts with the country rocks. The degree of metamorphism diminishes proportionally to the distance from the contact. For example, in the Chalba massif, in the region of Zabolonnyy spring and the headwaters of the Chalba River, one may observe granodiorites and diorite-granites at the very contact with the country rocks. As one moves away from the contact they give way to biotite-hornblende granites and then to biotite granites. The diorite-granodiorites and biotite-hornblende granites are conspicuous for the uneven distribution of dark-colored minerals, unbalanced quantitative and mineralogical composition, and other manifestations betraying their hybrid origin. These rocks were probably formed as a result of the assimilation of the country rocks (represented by hybrid rocks of the first phase, gneisses and schists) and by granitic magma at the upper structural stage at the time of granite formation.

The assay results for the second-phase granites, as well as for the hybrid facies of these granites, are given in Table 1 and are shown in the diagram of chemical composition (Figure 3).

From this Table it is obvious that the second-phase granites — as evidenced by comparison of analyses results with average types of rocks, according to Daly — approximate, in content the Post Cambrian granite and differ from it in higher SiO_2 and K_2O content and lower MgO , CaO , Na_2O , and Fe_2O amounts. Rb_2O , Rb_2O , Cs_2O and a small amount of iron are invariably present in the second-phase granites, just as they are in the first-phase hybrid rocks.

⁴Tin, as previously established by us, is normally present in the biotites of tin-bearing granites and is not found in non-stanniferous granites. Consequently, the presence in biotites of this or that type of granite, along with other indications, bespeaks their stanniferous nature, just as the presence of beryllium in the biotites of this or that type of granites, along with other indications, signifies that they contain beryllium.

The hybrid facies of the described rocks (samples c491 and 12) correspond to the average types of rocks and occupy an intermediate position between granites and granodiorites (sample 12). The second-phase granites are intruded by dikes of fine-grained biotite granites, aplites, pegmatites and granite-porphyrries with north westerly (320° - 330°) and north-easterly (30 - 40°) strike and a steep dip. The dikes range from several centimeters to several tens of meters in thickness. The contact of the dikes with the enclosing granites is sharp. However, the latter display slight silification and greisenization in the immediate vicinity of the contact.

In some sections the second-phase granites are greisenized. Greisenization is manifested in the formation of lithium-bearing muscovite and tourmaline which replace both the feldspars and dark-colored minerals. Sometimes greisenization is so advanced that the granites are transformed into quartzose-micaceous-tourmaline greisens. Zones of intensive silification and tourmalinization associated with lead mineralization occur rather frequently in these granites.

The third phase of the Myao-Chanskiy intrusive complex is represented by fine-grained biotite, alaskite, and tourmaline-bearing granites. In comparison to the second-phase granites, these granites are less widespread. They consist of small massifs not exceeding two square kilometers. The third-phase granites intrude and metamorphose the Jurassic sandy-shale deposits and the granites of the second phase.

A detailed petrochemical study of the third-phase granites and their comparison with the second-phase facies shows that, along with common features, they possess a number of distinguishing characteristics.

Thus, the biotite granites of the third phase are characterized by:

1. An even greater amount of orthoclase feldspar (50%) and respectively a correspondingly smaller proportion of plagioclase.
2. A more acid plagioclase represented by acid oligoclase.
3. Lack of hornblende and less biotite, whose content does not exceed 2%.

The results of incomplete assays of the biotites of the third-phase granites indicate that they contain more Li_2O , Rb_2O , and Cs_2O than the biotites of second-phase granites.

4. A greater quantity of cassiterite and tourmaline among its accessory minerals, and the presence of fergusonite. Thus, the leading

Table 3

Results of X-Ray-chemical Analyses of Zircons and Apatites from the Granitoids of the Myao-Chanskiy Intrusive Complex (in %).

Elements	First phase				Second phase				Third phase	
	Zircon from diorite-monzonites		Apatite for diorite-monzonites		Zircon from biotite granites		Apatite from biotite granites		Zircon from biotite granites	Apatite from biotite granites
	c511	c230	c548 ^a	c511 ^a	c487	c462	c487	c462 ^a	c489	c481 ^a
Hf	0,1	0,1	—	—	0,5	0,3	—	—	0,5	—
Y	0,1	—	0,1	0,2	0,5	0,3	2	0,5	1	2
Ge	—	—	0,05	0,05	0,1	0,05	0,1	0,1	0,5	0,5
La	—	—	—	—	0,5	—	—	—	0,3	—
Pr	—	—	—	—	—	—	—	—	0,05	—
N	—	—	0,05	0,05	—	—	0,1	0,1	0,3	0,3
Gd	—	—	—	—	—	—	—	—	0,05	—
Dy	—	—	—	—	—	—	—	—	0,05	—
Er	—	—	—	—	—	—	—	—	0,05	—
Th	—	—	—	—	—	—	—	0,2	2	0,3
U	—	—	—	—	—	—	—	—	0,3	—
Pb	0,3	—	—	—	—	—	—	—	—	—
Sn	—	0,3	—	—	—	—	—	—	—	—
As	—	—	—	—	—	0,05	—	—	—	—

NOTE: Comma represents decimal point.

accessory minerals of third-phase granites are zircon, apatite, cassiterite, pyrite, tourmaline,⁵ fergusonite, molybdenum and magnetite.

5. A large proportion of yttrium and elements of the rare-earth group in zircons and apatites of the third phase as compared to second-phase granites (Table 3). Also present in the zircons of the granites are uranium and thorium lacking in the zircons of the second-phase granites.

6. A greater amount of silica and potassium oxide, and smaller proportions of ferric oxide, calcium oxide and magnesium (Table 1, Figure 3).

7. A greater quantity of lithium, rubidium, cesium, and boron (Table 1).

The tourmaline-bearing granites differ from the biotite granites of the third phase by the following characteristics:

⁵Particularly interesting is the presence of acicular tourmaline crystals in one of the principal rock-forming minerals — quartz. This tourmaline, which was formed simultaneously with the quartz during the final phases of the magmatic stage, indicates how rich the magma was in boron which began to separate even prior to the end of the actual magmatic stage.

1. Two generations of tourmaline are to be observed in them, the tourmaline amounting to 8-10%.⁶ The results of incomplete chemical analyses of the tourmalines and tourmaline-granites are recorded below in Table 4.

As this analysis shows, lithium is always present in the tourmalines of tourmaline-bearing greisens and tourmaline rocks, while rubidium and cesium are absent. A somewhat higher percentage of potassium oxide is observable in the tourmalines of the tourmaline-bearing granites as compared to those of the tourmaline rocks.

The results of the spectral analyses of tourmalines extracted from tourmaline-bearing granites showed that the ever-present lithium, zirconium

⁶The first-generation tourmaline occurs in the form of well defined prismatic crystals and has a deep reddish-brown or blue color. The first-generation tourmaline is eroded and replaced by quartz and apatite. Sometimes it reveals idiomorphic crystals of zircon surrounded by pleochroic aureoles. Second-generation tourmaline also fills in the small fissures in the rock. In contrast with first-generation tourmaline, it has a light-blue or brownish tint. First generation tourmaline crystallized, probably, by the end of the magmatic process (somewhat earlier than apatite and quartz and simultaneously with them), whereas the crystallization of second-generation tourmaline occurred later in connection with the post-magmatic processes.

Table 4

Results of Chemical Analyses for Individual Tourmaline Components¹
from various rocks of the Myao-Chanskiy Intrusive Complex (in %).

Elements	Tourmaline-bearing granites, Sample c482	Quartzitic- micaceous tourmalinic greisens, Sample 488	Tourmalinized diorite-mon- zonites, Sample c556	Tourmaline rocks, Sample c120
N ₂ O	2,19	2,12	2,19	2,28
K ₂ O	0,27	0,24	0,18	0,19
Li ₂ O	0,024	0,022	0,011	0,028
Rb ₂ O	0,00	0,00	0,00	0,00
Cs ₂ O	0,00	0,00	0,00	0,00

¹The analysis was made at the chemical laboratory of the IGEM
of the USSR Academy of Sciences by G. Ye. Kalenchuk.

NOTE: Comma represents decimal point.

ic, copper, gallium, and vanadium, are characteristic of them, while yttrium, zinc, strontium, and lead also often occur.

2. Biotite is either altogether absent in the tourmaline-bearing granites, or occurs in negligible amounts.

3. Among the accessory minerals there is zircon and fergusonite and correspondingly zircon apatite.

4. Albitization processes are well developed.

5. Tourmaline-bearing granites, like the biotite granites of the third phase, in chemical composition (Table 1, Figure 3) approximate the average composition of alaskites, as defined by Daly, but differ from it by a somewhat lower Na₂O and Na₂O content and a larger amount of K₂O. In comparing tourmaline-bearing granites with third-phase biotite granites, we find that they closely approximate the latter and are distinguishable from them by a slightly higher percentage of CaO and Na₂O as well as by a considerably larger amount of boron.

In a number of sections the biotite and tourmaline-bearing third-phase granites are greisenized. Tin ore manifestations are observable in the exomorphic zones of the tourmaline-bearing granites.

As mentioned earlier, dike formations (of average and basic composition) are widespread in the Myao-Chanskiy region even though they are still inadequately studied. A part of these dikes is directly related to the hybrid rocks of the first phase. Another part, in our opinion, is, probably associated with the mantle of young Tertiary and Quaternary effusives (Figure 2). Finally, apparently, there are the dikes and small intrusive bodies which completed the formation of the Myao-Chanskiy intrusive complex.

The specific characteristics of the three groups of dikes are not yet entirely defined and should become the object of further investigations. At this stage we shall note only that according to available preliminary data, dikes associated with the hybrid rocks of the first phase are characterized by higher potassium content, whereas those that are related to the younger effusive formations are distinguishable by a more basic composition and lack of any mineralization.

In the conclusion of this brief geologic, petrographic, and petrochemical characterization of the Myao-Chanskiy complex intrusive rocks we shall make the following statement. A detailed geologic, petrographic and petrochemical study of the intrusive rocks of all three phases of the Myao-Chanskiy complex makes it possible to postulate that these phases are the derivatives of a single slowly evolving magmatic chamber. The assimilation and hybridization phenomena, which occurred in the initial developmental stages of the magmatic chamber, must have upset the normal course of magmatic differentiation, and caused the appearance of hybrid rocks, on the one hand, and of third-phase granites, on the other, the latter being enriched by alkalis, volatile elements, and ore components.

Comparison of the intrusive rocks belonging to various phases shows that, in spite of a number of substantial differences in their petrographic contents, a set of common features may still be discerned. For example, for the representatives of all phases, even for the first-phase hybrid rocks, a high percentage of potassium and the constant presence of lithium, boron, rubidium, cesium, and tin are characteristic. A gradual decrease in content of iron, magnesium, and calcium, and a gradual increase of silica and alkalis, particularly potassium, are observed from the

first phase to the third. The amount of lithium, cesium, boron, tin, as well as certain other elements, steadily increases from the first phase to the tourmaline-bearing granites of the third phase.

As to the accessory minerals, the proportion of zircon, tourmaline, and cassiterite increases gradually from the first-phase hybrid rocks to the tourmaline-bearing third-phase granites. Apart from zircon, apatite, cassiterite, tourmaline, and molybdenite, fergusonite also may be listed among the leading accessory minerals of the third-phase granites. The amount of yttrium, rare-earth elements, as well as uranium and thorium (Table 3), increases in the zircons and apatites belonging to the more recent phases of the Myao-Chanskiy intrusive complex. This fact allowed us to conclude that the accumulation of yttrium, rare earths, uranium and thorium occurred in the final stages of the magmatic chamber development during the crystallization of the third-phase granites.

All of these factors indicate that the development process in the described magmatic chamber must gradually lead to greater acidification of the magma and its enrichment with volatile constituents and ore components. At the same time the role of calcium, magnesium, and iron gradually diminishes as the significance of silicic acid increased. The observable deviations from this scheme in the Myao-Chanskiy region are attributable to the influences of the enclosing medium.

It is important to note that the intrusives of all three phases of the Myao-Chanskiy complex contains, sometimes more, sometimes less clearly expressed traces of hybridization. However, abyssal assimilation is characteristic for the first phase in the lower structural stage,⁷ whereas for the second and third phases, hybridism is local and traceable directly to the country rocks. Thus, in a number of places in the endomorphic zones of the second-phase granites (the eastern contact of Chalba massif) quartzose diorite-monzonites and granodiorites are observable at the contact with the country rocks. Away from the contact these dioritic facies are gradually replaced by normal granites.

The factual material in our possession attests to the fact that the processes of assimilation and hybridization which developed quite intensively in the initial stage of the magma chamber development, must have somewhat affected the

second and third-phase granites during the formation of the first-phase hybrid rocks. Only such an influence explains the presence in third-phase granites (sample 48) — which by their composition resemble alaskites — of a high percentage of calcium oxide and parameter c (Table 1). A study of the young (Upper Cretaceous) granitoids in the Myao-Chanskiy intrusive complex has confirmed the conclusions made earlier by M. G. Rub [9], in the case of the granitoids in the Upper Paleozoic complex (Prikhankayskiy region), to the effect that hybridization processes accompanying the formation of complex multi-phase intrusions occur both in the upper and in the lower structural stages. Moreover, the latter upset the normal development of magmatic differentiation and exerted a certain influence on the specialization of intrusions.

SEQUENCE OF POSTMAGMATIC FORMATIONS ACCOMPANYING THE GRANITOIDS OF THE MYAO-CHANSKIY INTRUSIVE COMPLEX

In the Myao-Chanskiy region various types of extensive post magmatic formations occur in close areal association with the granitoids of the Myao-Chanskiy intrusive complex (Figure 1). Postmagmatic formations usually coincide with dislocation zones having submeridional trends, or at intersections of dislocation zones having submeridional and latitudinal trends. Studies of the composition and the formative peculiarities of the mineralized zones revealed that their formation progressed in a rather complicated manner and consisted of several stages successively replacing each other in terms of time. These stages are separated by clearly defined periods of inter-mineralization adjustment.

The following postmagmatic formations were revealed in various mineralized zones of the investigated area (enumeration is made from the earliest to the latest): quartzo-feldspathic metasomatites containing tourmaline or axinite and sphene; quartz-micaceous-tourmaline greisens; tourmaline and quartz tourmaline rocks and the accompanying essentially quartz and quartz-sericite rocks formed as a result of near-vein metasomatic manifestations;⁸ combed vein-quartz with a cassiterite matrix, quartz-sulfide, and quartz-carbonate-sulfide ores.

Such a sequence of the described rocks becomes apparent from the structural interrelations of

⁸There is no unanimity of opinion on this matter. In particular, V. V. Onikhimovskiy believes that the formation of quartz-sericitic rocks, the essentially quartz rocks ("secondary quartzites"), quartz-tourmalin and tourmaline rocks occurred in four different phases of the hydrothermal stage.

⁷As previously stated, the intrusives of the first phase were probably formed as a result of deep-seated magma assimilation of sandstones, porphyrites, and other rocks rich in iron and magnesium.



FIGURE 4. Quartzite fragments cemented by younger quartz.
(3/4 natural size).

the facies. In speaking of the essentially quartz and quartz-sericite rocks, one should note that they are confined to the outer portions of the tourmaline and quartz-tourmaline zones. Moreover, the quartz-sericite rocks occur between the essentially quartz and sericitized country rocks. The essentially quartz, quartz-sericite, quartz-tourmaline and tourmaline rocks are closely associated with each other. They were formed in a single process involving the action of postmagmatic acid solutions containing a high percentage of boron, on sandy shales, effusives and intrusives. Here the principal role is the metasomatic replacement of the country rocks and, to a lesser degree, by those of deposition in open cavities.

Investigation of the essentially quartz and quartz-sericite rocks makes it possible to assume that their formation is related to the manifestation of infiltrative metasomatic zoning. Indeed, according to D.S. Korzhinitsy's data [7, 8], several metasomatic zones must occur along the path of the percolation of acid solutions into the replaceable rocks. Moreover, at each metasomatic front between two zones, the acidity of the percolating solution diminishes causing a corresponding decrease in the solubility of alumina producing partial deposition.

In this manner, replacement must be accompanied by gradual enrichment of the successive metasomatic zones with alumina displaced from the zones of total leaching. From

this point of view the regular replacement of essentially quartz rocks — the product of total leaching — by quartz-sericite rocks is quite natural. From incomplete analyses of quartz-sericite rocks it appears that lithium (Li_2O - 0.04-0.5%), rubidium (Rb_2O 0.01-0.03%), cesium (Cs_2O - about 0.01%) and boron are always present in them. The results of semi-quantitative spectral analyses show that strontium, zirconium, tin, gallium, lead, zinc, copper, and silver also are characteristic secondary elements of the quartz-sericite rocks.

According to the incomplete chemical analyses of the discussed rocks, and the results of semi-quantitative spectral analyses, the leading accessory elements in the tourmaline and quartz-tourmaline rocks are strontium, lithium, rubidium, cesium, zirconium, boron, with gallium, vanadium, and copper always present. The tourmalines in question differ from the high-temperature tourmalines of the granites in the absence of germanium and yttrium, higher content of vanadium, and the presence of nickel and cobalt.

Cassiterite in smaller quantities is to be found in the greisens and the association of metasomatic rocks referred to above. In the latter case, along with the cassiterite characteristic for the given association, the principal occurrence is superimposed cassiterite deposited in the subsequent quartz-cassiteritic phase of the hydrothermal stage.




FIGURE 5. Fragments of quartz-tourmaline rocks (right and below intersected and cemented by combed quartz containing cassiterite, sample 615. (3/4 natural size).




FIGURE 6. Replacement of needle-shaped tourmaline crystals by cassiterite (dark grey). Section 439. Without analyzer. 72 x.

The maximum quantity of cassiterite was deposited together with the combed quartz after the formation of the association of the metasomatic rocks.

This mineralization stage is distinctly separated from the preceeding one (quartz-tourmaline) by an interval of time and intense crushing. Usually the combed quartz containing the cassiterite contains a large quantity of fragments of quartz-sericite, quartz-

tourmaline, and tourmaline rocks (Figures 4 and 5). The cassiterite often surrounds and replaces the fragments of quartz-tourmaline and tourmaline rocks (Figure 5) and the individual tourmaline crystals (Figure 6). Numerous semi-quantitative spectral analyses of the cassiterites revealed the invariable presence in them of niobium in quantities ranging from 0.00 to 0.0n%. Frequently, lead may also be noted, and sometimes antimony.

The final stage of minero-genesis is the quartz-sulfide-carbonaceous stage, during which an important role is played by the carbonates represented mainly by calcite and siderite. Ordinarily, these minerals fill in the hollows between the previously deposited quartz crystals, or they form stringers penetrating the previously formed hydrothermal rocks. The low-temperature sulfides — pyrite, galenite, sphalerite, and chalcopyrite are developed in the veinlets along with the calcite and quartz.

In summing up all of the above discussion of this section, it is possible to draw the following basic conclusions.

1. The formation of mineralized zones is a long and complicated process. It consists of several phases and stages successively replacing each other. Thus, the greisen stage is replaced by a lower-temperature hydrothermal stage comprising several phases; quartz-tourmaline, quartz-cassiterite, quartz-sulfide, and quartz-sulfide-carbonate. These phases are usually separated by well-defined periods of inter-mineralization, and follow each other over a certain interval.

2. In the formative process of the mineralized zones an exceptionally important role is played by metasomatic processes, which are attributable to favorable structural circumstances and the high mobility of the metalliferous solutions which, as a rule, are rich in volatile constituents, particularly boron.

3. A definite regularity exists in the manifestation of the described types of mineralization. For example, the early mineralization stages are most extensively developed in and near the granodiorite massifs whereas the later stages are at some distance from the intrusions.

Relationship Between Mineralization and the Granitoids of the Myao-Chanskiy Intrusive Complex

The intrusives belonging to all three phases of the Myao-Chanskiy intrusive complex and the associated vein rocks are intersected and replaced by a variety of types of postmagmatic formations extensively developed in the investigated area. They are very heavily silicified, sericitized, and tourmalinized.

At the present time several facts have come to hand which makes it evident that the well-developed stanniferous and polymetallic mineralization in the Myao-Chanskiy region is related to the development of a composite igneous complex.

Among these facts one should point out the following.

1. The areal association of the mineralization with the intrusives of all three phases of the Myao-Chanskiy complex (Figure 1).

2. The horizontal zoning in the distribution of various types of post-magmatic formations. For example, the earlier and high-temperature minero-genetic stages are most widely developed in the intrusives and in their vicinity, while more recent and low-temperature types of mineralization occur away from them.

3. Comparable depths of formation of the intrusives and the ore bodies. According to our estimates, the formation of the intrusives in the Myao-Chanskiy region must have occurred at a depth of 1.5-2 kilometers. The various postmagmatic rocks were also formed at approximately the same depth.

4. Development of the same types of minerals in the later stages of cooling in the intrusives and in the various postmagmatic formations. Thus, among the third-phase granites of the Myao-Chanskiy igneous complex one finds tourmaline-bearing granites, the tourmaline being formed at the end of the actual magmatic stage. In addition, the second- and third-stage granites show signs of greisenization associated with autopenumatolysis. The greisenized granites and greisens, as well as quartz and muscovite, always contain tourmaline. Tourmalinization occurs in the exocontact aureoles of the intrusives belonging to all three phases. At the same time, tourmaline is the leading mineral of various postmagmatic formations so abundant in the Myao-Chanskiy region.

It is evident from the above that the magma which was responsible for the formation of the Myao-Chanskiy granitoid complex and the accompanying postmagmatic rocks was enriched with boron. The separation of boron began in the early stages of development of the magmatic chamber during the crystallization of granites, and the accumulation occurred later in connection with the action of postmagmatic processes.

5. The presence of cassiterite among the accessory minerals of the granitoids belonging to all three phases of the Myao-Chanskiy complex and the associated vein rocks.

6. Geochemical affinity of the granitoids of all three phases of the complex and the accompanying postmagmatic formations. The available factual material permits the singling out of a group of transient elements whose deposition began in the early stages of the development of the magmatic chamber during the crystallization of the granitoids of the Myao-Chanskiy intrusive complex. Accumulation took place later in connection with post-magmatic activity. These elements include: lithium, rubidium, cesium, copper, strontium, boron, tin, tungsten, gallium, lead, and zinc




FIGURE 7. Geochemical Diagram of the Granitoids in the Myao-Chanskiy Intrusive complex and the associated postmagmatic formations.

1 - trace - elements characteristic of the first-phase hybrid rocks; 2 - trace - elements characteristic of the second-phase granites; 3 - trace - elements characteristic of the third-phase granites; 4 - trace - elements characteristic of all postmagmatic formations; 5 - trace - elements characteristic only of high-temperature post-magmatic formations; 6 - trace - elements sometimes observable in granitoids.




FIGURE 8. Variation of cesium, rubidium, and lithium content in the granitoids of the Myao-Chanskiy igneous complex.

1 - Phase I: diorite-monzonites; 2 - Phase II: biotite and biotite hornblende granites; 3 - Phase III: biotite, alaskite, and tourmaline-bearing granites; 4 - high-temperature postmagmatic formations: greisens, sericite and quartz-sericite rocks; 5 - lower-temperature postmagmatic formations: quartz-tourmaline and tourmaline rocks.

Figure 7). Some of these elements, particularly boron, rubidium, and cesium, are present in the representatives of all three phases of the Myao-Chanskiy complex in quantities above Clarke values. Other metals - tin and lead - appear in amounts exceeding Clarke values only in the tourmaline-bearing granites of the third phase. It is interesting to note that the rubidium, rubidium, and cesium content increases gradually from first-phase to third-phase rocks and is maximum in the high-temperature postmagmatic formations: greisens and quartz-sericite rocks. In the lower temperature postmagmatic formations the proportion of the mentioned elements again decreases (Figure 8). The maximum content of boron and tin also increases from the first-phase rocks to the tourmaline-bearing granites of the third phase. For boron, the peak is reached in the tourmaline rocks, for tin - at a more recent stage coinciding with the formation of the combed quartz.

All of the factual material at our disposal - which is only partially discussed in this article - evidences that tin mineralization, widespread in the investigated area, is associated with the chamber responsible for the formation of the Myao-Chanskiy intrusive complex granitoids. The separation of metalliferous solutions from the magma occurred repeatedly, but the maximum amount was segregated after the formation of the tourmaline-bearing granites of the third phase and represents a derivative of the residual magmatic melt saturated with volatile constituents and ore elements.

A close genetic relationship can be established in the Myao-Chanskiy region between the relatively high-temperature ore manifestations of the greisen and cassiterite-quartz types and the relatively lower temperature cassiterite-

quartz-sulfide and sulfide ore manifestations. Therefore, if the genetic association of the greisens and tin mineralization with the granitoids of the Myao-Chanskiy complex appears to be beyond doubt, then the cassiterite-quartz-sulfide and sulfide mineralizations must also be related to the same chamber as parts of a single process.

SUMMARY

Studies of the granitoids in the Myao-Chanskiy igneous complex showed that they are specialized, and confirmed our conclusions formulated earlier for the case of the Prikhankayskiy region [9] concerning the existence of specialized intrusions. Comparison of the specialized intrusions of different age in different structure-facies zones revealed that, along with common features, they also possess a number of specific petrochemical characteristics. Thus, for example, the Upper Paleozoic specialized intrusions of the Grodekovo complex (Prikhankayskiy region) confined within the limits of the Khanayskiy tectonic massif and the Upper Cretaceous granitoids of the Myao-Chanskiy intrusive complex (Komsomol'skiy district of Khabarovsk Krai) located in the zone of Cretaceous folding,⁹ in addition to these common features, possess a number of specific petrochemical characteristics. Both the former and the latter belong to the multi-phase type, and in both cases there was a gradual increase in oxidation and potassium content in the rocks from the earliest phases to the more recent ones. Quite characteristic of both is the extensive

⁹Towards the Cenozoic the entire western part of lower Priamur'ye, including the Komsomol'sk district, became stable and approximated platform conditions.

development of hybridization processes occurring both in the upper and lower structural stages. The leading accessory minerals of the Upper Paleozoic granitoids in the Grodekovo intrusive complex are zircon, apatite, rutile, fluorite, cassiterite, and magnetite. In the Upper Cretaceous stanniferous granitoids of the Myao-Chanskiy intrusive complex the prevalent accessory minerals are zircon, apatite, fluorite, cassiterite, tourmaline, magnetite, molybdenum, and fergusonite. Characteristic of the former are such accessory elements as cesium, beryllium, lithium, zirconium, uranium, fluorine, boron, tin, gallium, and lead (Figure 9); and for the Myao-Chanskiy igneous complex these are: cesium, rubidium, zirconium, lithium, zirconium, gallium, copper, and lead. Typical of the third-phase granites of this complex are also molybdenum, uranium, and rare earths, and niobium.

Thus, the stanniferous granitoids of the Myao-Chanskiy igneous complex — along with a number of such common petrochemical features as abundance of potassium, tin, zircon, lithium, and rubidium — possess a number of peculiarities which distinguish them from the Upper Paleozoic granitoids of the Grodekovo intrusive complex. These peculiarities include:

1. A distinctly subordinate role for fluorine and a far greater importance for boron. In the granitoids of the Myao-Chanskiy intrusive complex and in the associated postmagmatic formations, fluorine has a very limited distribution, whereas in the granitoids of the Grodekovo igneous complex it is a dominant element.
2. The role of beryllium is distinctly subordinate. In the granitoids of the Myao-Chanskiy intrusive complex beryllium rarely occurs and always in quantities far below Burke values.
3. Cesium is always present. Cesium in small quantities occurs in the granitoids of the three phases of the Myao-Chanskiy intrusive complex, but attains its maximum in the high-temperature postmagmatic formations (Figure 7), and is absent in the low-temperature facies.
4. The presence among the accessory minerals of the third-phase granites of the Myao-Chanskiy intrusive complex of fergusonite which does not occur in the granitoids of the Grodekovo intrusive complex, and other intrusions.

All the materials discussed herein also testify to the major role of petrochemical criteria in establishing the character of the relationship between mineralization and magmatism. Moreover, comparison of specialized

intrusions of different age in different structure-facies zones show that, along with a number of common features, they possess certain specific petrochemical characteristics.

The above conclusions must be verified in other regions.

REFERENCES

1. Abdullayev, Kh.M., *Geneticheskaya svyaz' orudeneniya s granitoidnymi intruziyami. [THE GENETIC RELATIONSHIP BETWEEN MINERALIZATION AND GRANITOID INTRUSIONS]*: Gosgeolizdat, 1954.
2. Afanas'yev, G.D., *Problema granitoidov i nekotoryye voprosy svyazannoy s nimi metallogenii. Magmatizm i svyaz' s nim poleznykh iskopayemykh. [THE PROBLEM OF GRANITOIDS AND CERTAIN QUESTIONS OF RELATED METALLOGENY. MAGMATISM AND ITS RELATIONSHIP TO COMMERCIAL MINERALS]*: Trudy I-go vsesoyuznogo petrograficheskogo soveshchaniya. Izd-vo Akad. Nauk SSSR, 1953.
3. Betekhtin, A.G., *O geneticheskoy svyazi gidrotermal'nykh obrazovaniy s intruziyami. V kn. "Osnovnyye problemy v uchenii o magmatogennykh rudnykh mestorozhdeniyakh". [THE GENETIC RELATIONSHIP OF HYDROTHERMAL FORMATIONS TO INTRUSIVES. From the book: "THE BASIC PROBLEMS IN THE STUDY OF MAGMATIC ORE DEPOSITS"]*, Izd-vo Akad. Nauk SSSR, 1953.
4. Vinogradov, A.P., *Zakonomernosti raspredeleniya khimicheskikh elementov v zemnoy kore. [REGULARITIES OF DISTRIBUTION OF CHEMICAL ELEMENTS IN THE EARTH'S CRUST]*: Geokhimiya, No. 1, 1956.
5. Vol'fson, F.I., *Problemy izucheniya gidrotermal'nykh mestorozhdeniy. [PROBLEMS OF INVESTIGATING HYDROTHERMAL DEPOSITS]*: Izd-vo Akad. Nauk SSSR, 1953.
6. Gotman, Ya.D., *K voprosy ob olovonosnykh intruziyakh Primor'ya. [THE PROBLEM OF TIN-BEARING INTRUSIONS IN PRIMOR'YE]*: Izv. Akad. Nauk SSSR, ser. geol., No. 6, 1943.
7. Korzhinskiy, D.S., *Obshchiye svoystva infil'tratsionnoy metasomaticheskoy zonal'nosti. [THE GENERAL PROPERTIES OF INFILTRATIVE METASOMATIC ZONING]*: Dokl. Akad. Nauk SSSR, 78, t. No. 1, 1959.

8. Korzhinskiy, D.S., Zavisimost' aktivnosti komponentov ot kislotnosti rastvora i posledovatel'nosti reaktsii pri poslemagmaticheskikh protsessakh. [COMPONENT ACTIVITY AS A FUNCTION OF SOLUTION ACIDITY AND ORDER OF REACTION IN POST-MAGMATIC PROCESSES]: Geokhimiya, No. 7, 1956.
9. Rub, M.G., Granitoidy prikhankayskogo rayona i osnovnyye cherty ikh metalonosti. [THE GRANITOIDS OF THE PRIKHANKAYSKIY REGION AND THE PRINCIPAL INDICATIONS OF THE PRESENCE IN THEM OF METALS]: Izv. Akad. Nauk SSSR, ser. geol., No. 4, 1955.
10. Favorskaya, M.A., O svyazi olovyanogo orudneniya s magmatizmom. [THE RELATIONSHIP BETWEEN TIN-MINERALIZATION AND MAGMATISM]: Izv. Akad. Nauk SSSR, ser. geol., No. 4, 1955.

Institute of Mining Geology, Petrography,
Mineralogy, and Geochemistry, U.S.S.R.
Academy of Sciences,
Moscow

Received, January 20, 1960

MAGNETIC-FRACTIONAL-MINERALOGICAL STUDIES OF ROCKS¹

by
F. N. Yefimov

I presented a report on the first results obtained in making a magnetic-fractional-mineralogical analysis (MFMA) of various rocks and ores, mainly from the Ukrainian Shield, and partly from the Kusrk Magnetic anomaly, at the Second All-Union Petrographic conference held in May 1958 in Tashkent.

Investigations were carried out during the second half of 1958 and in 1959 to improve the method and verify the effectiveness of using these analyses to study the composition and the magnetizability of crystalline rocks exposed by drilling in Western Bashkiria and Eastern Turtaria. At the same time the improved MFMA method was used to investigate certain types of magmatic and metamorphic rocks from the Ukrainian Shield, the Chernigov shales, the Urals, and ores from the Caucasus and Eastern Siberia; such as gabbros, olivine gabbro, gabbro-amphibolites, gabbro-anorthosites, gabbro-diabases, basalts, diabases, pyroxenites, diorites, andesites, quartz diorites, granodiorites, granosyenites, syenites, dioritic granites, hybrid granites — amphiboles, amphibole-pyroxenes, and pyroxenes, charnockites, amphibolites, skarn-type rocks, granulite-gneisses, gneisses (biotitic, amphibole-biotite, amphibole-biotite, garnet-biotite, garnet-biotite-sillimanite, cordierite, and others), magnetitic and hematite-magnetite gneisses, magnetitic hornfels and schists, magnetitic websterites and ores, pyrrhotite and hematite ores.

This diversified assemblage of rocks and ores was required in order to determine the general nature of the relationships between the MFMA parameters and the composition of the geologic formations of different origin and metamorphism.

Altogether about 500 analyses were made. Obviously, the number of samples investigated for us is not sufficient to define concretely the revealed mineralogical and physical

regularities with respect to each of the enumerated varieties of rocks. However, the data at our disposal, and the considerable number and diversity of rock samples analyzed by us during the first development of MFMA, provide sufficient grounds, even now, to recommend the magnetic-fractional-mineralogical analysis for widespread use by geologic and geophysical organizations. As MFMA is gradually put to use in solving various geologic survey problems, the areas of its effective application will become more precisely evident. At the same time new factual material will be collected characterizing the details and peculiarities of the mineralogical and physical relationships between rocks of the same type, even if they are from widely separated areas and of different ages.

The newly developed analytical method was designated as a magnetic-fractional-mineralogical analysis because it is based on the utilization of the magnetic properties mostly of ferro-magnetic minerals occurring in nature, on the peculiarities of the distribution of these minerals in the separation process of the rock-powder batch and its segregation into magnetic and "non-magnetic" fractions.

The MFMA diagram shown in Figure 1 gives a good idea of the nature and sequence of operations involved in the analysis. The principal operations are:

1. Determination of the magnetic susceptibility (χ) of the original rock powder with grains measuring less than 0.25 mm.
2. Separation of the initial batch (1-2 g) of rock powder, with grains of less than 0.15 mm, in the field of a permanent electromagnet by the wet process method under rigidly fixed and identical conditions for all analyzed rocks ($H = \sim 900$ e).
3. Weighing of the magnetic fraction with a 0.1 mg accuracy and determination of parameter F_p equivalent to the weight percentage of ferro-magnetic minerals in the powder.

¹Magnitno Fraktsionno-Mineralogicheskoye issledovanie Gornykh Porod. pp. 24 - 36.

$$J = \kappa H + J_n.$$

i. e., the force attracting the particles to the poles of a permanent electromagnet depends not only on the susceptibility of the ferromagnetic material, but also on the magnitude of its natural residual magnetization. It is obvious that those particles which are more powerfully attracted to the poles apart from equal susceptibility values, also have some residual magnetization.

Under the selected separation conditions the overwhelming proportion of the relatively large ferromagnetic separates get into the magnetic fraction. The small ferromagnetic separates which are intimately intergrown with their silicates (pyroxenes, amphiboles, biotite, olivine, garnet), or ores (ilmenite, hematite, pyrite) minerals migrate into the "non-magnetic" fraction. Fine ferromagnetic particles settle into the "non-magnetic" fraction, first of all, because of their relatively weak magnetization - the coarser particles always have a higher κ - value than the fine ones. In the second place, this happens because their magnetic gravitation to the poles of the electromagnet turns out to be inferior to the gravity of the carrying silicate or ore minerals.

Depending on the composition of the investigated rocks, ferromagnetic materials with a definite composition, morphology, and in given quantities, will get into the "non-magnetic" and magnetic fractions. In accordance therewith, definite values of F_p (weight percentage of ferromagnetic material) and κ_n , which constitute the basic initial parameters of magnetic-fractional-mineralogical analysis, will be obtained for each variety of rock.

Among the initial MFMA parameters one should also include parameter κ . This parameter is measured under the specific conditions of the proposed analysis and characterizes the quantity, composition, and quality of the ferromagnetic materials contained in the initial batch of the investigated rock.

The ratios of the above three initial parameters make it possible to determine three very important additional parameters, namely:

$$\text{parameter } M = \frac{\kappa}{\Phi c},$$

$$\text{parameter } H = \frac{\kappa_n}{\kappa},$$

$$\text{parameter } O = MH = \frac{\kappa_n}{\Phi c}.$$

In addition to the enumerated parameters the MFMA complex also has three known very important parameters - natural residual




FIGURE 1. Diagram of the successive steps in magnetic-fractional-mineralogical analysis.

4. Determination of the magnetic susceptibility of the "non-magnetic" fraction (κ_n) characterizing the contents and quality of the ferromagnetic minerals not attracted to the poles of the electromagnet under given separation conditions, and determination of the magnetizability of the paramagnetic minerals of the rock.

It is known that the magnetization intensity of ferromagnetic material (J) is proportional to its magnetic susceptibility (κ) and the intensity of the magnetizing field (H)

$$J = \kappa H.$$

Considering the fact that separation proceeds at a single and constant value of H , the magnetization intensity is directly proportional to the degree of susceptibility.

In the general case, a ferromagnetic material possesses not only inductive, but natural residual magnetization J_n , as well. Then

magnetization (J_n), parameter $Q = \frac{J_n}{\bar{J}}$, and the volume weight of the rocks (σ).

If required, the number of physical parameters to be used may be increased by introducing such quantities, for instance, as specific gravity, total porosity, the Curie point. However, a larger number of parameters cannot substantially affect the effectiveness of MFMA, which is basically determined by the six specific parameters and density.

A study of sections and polished sections is required to interpret the physical and geological nature of MFMA parameters. Chemical, X-ray diffraction, spectral and other types of analyses also may be made on a moderate scale for the same purpose.

Of particular importance in a correct interpretation of MFMA data is the mineralogical study of the ferromagnetic minerals, not only of ferruginous formations, but for any types of rocks.

In studying polished sections of rocks using different values for MFMA parameters, it has been possible to establish the fact that ferromagnetic minerals in rocks are represented predominantly by magnetite and its varieties: titanomagnetite, magnesioferrite, and sometimes maghemite, muschketowite, and certain others, as well as by pyrrhotite. Magnetite and its variants are exceedingly widespread in nature and may be encountered in some concentration or other in almost all geologic formations of acid, alkali, basic, or ultra-basic composition of magmatic or metamorphic origin. Pyrrhotitic mineralization is far less abundant in nature and is usually found in rocks affected by hydrothermal alteration.

The described ferromagnetic minerals occur in rocks in various forms of separates diversely associated with other rock-forming minerals, and with varied composition. One may, for instance, observe primary magnetites belonging to several generations, and magnetites of secondary origin separated as a result of geochemical transformations of silicates or other minerals caused by metamorphic processes. Allotriomorphic forms are characteristic of the primary precipitations of magnetite. This is because this mineral was precipitated after many silicate rock-forming minerals and filled their intergranular spaces. The secondary precipitates of magnetite are concentrated within the minerals which are responsible for their formation. These are considerably finer grained than the primary magnetites.

We are not yet in a position to provide a more or less conclusive classification of the forms of the primary and secondary magnetitic separates since the data available on this matter are still insufficient. In particular, it

is still difficult to explain the nature of the fine deposits of magnetite which can be found more or less evenly distributed in certain types of basic rocks. Are they a more recent generation of primary precipitates, or do they represent a special form of secondary magnetitic deposition? There still is no differentiated characteristic of the secondary magnetitic separates formed as a result of the opacitization, amphibolization, biotitization, and chloritization of silicate minerals.

However, the data at our disposal convince us that all these questions will be answered in the process of further detailed rock investigations using MFMA. Moreover, these data make it possible, even today, to define the over-all character of the relationship between the MFMA parameter values and the forms of the magnetitic separates in rocks having different composition, metamorphism, and genesis.

It has also been established that pure separates of magnetite occur rather unfrequently. Often they contain inclusions of other ore minerals. More frequent are the inclusions of ilmenite and rutile. In titanomagnetite sometimes contains hematite inclusions. When isomorphous admixtures of magnesium, chromium and other metal oxides are present in magnetite, the mineral is referred to as magnesioferrite, chromomagnetite, etc. Here, as in the case of titanomagnetite, the ferromagnetic properties are retained by the magnetite, while the isomorphous admixtures of other metal oxides play a passive role, i.e. having the significance of paramagnetic inclusions. In speaking of passivity, we have in mind the very low susceptibility of paramagnetic inclusions. But the role of the latter can not be defined as passive when one considers the natural residual magnetization of some types of basic rocks. Under definite conditions the xenolith plates may play the role of energy barriers in the process of natural demagnetization of rocks.

It is a well-known fact that magnetite and its varieties oxidize (martitize) in supergene zones, i.e. turn into hematite, and subsequently also into hydrous ferric oxides (limonite). It is also known that in a reducing atmosphere hematite may turn into magnetite (muschketowite). In a number of cases, for example, in special investigations of iron-ore formations the processes of magnetite martitization and hematite muschketowitization may present a matter of independent practical and scientific interest and be studied with MFMA parameters. When, however, rocks become the object of investigation, the processes of martitization not only fail to help, but actually hamper, and at times render impossible, the exposure of natural connections between general mineralization and ferromagnetic mineralization. To avoid gross errors in interpreting the nature of MFMA parameters, samples containing deep-seated martitization of magnetite should be discarded before examination.

Special attention should be given to maghemite. Until recently most investigators considered this mineral as a ferromagnetic variety of hematite (5, 8). MFMA studies of several maghemite-bearing samples of magnetite from the Kezhma deposits, a maghemite specimen and a few hematitic and magnetitic ore samples from deposits located elsewhere in the Soviet Union, lead us to conclude that maghemite is actually a form of oxidized magnetite. The spinel-type cubic structure of magnetite is preserved in maghemite and the difference between the two minerals is that a part of divalent iron ions is replaced in maghemite by trivalent ions. The quantitative ratio between the di- and trivalent iron ions accounts for the fact that maghemite displays lower susceptibility and higher residual magnetization than magnetite. Maghemite is, probably, widely disseminated in nature and is present in small quantities not only in some types of hematite and magnetite ores, but also in rocks with different compositions. This explains the anomalously high residual magnetization values of these formations. This question will be answered conclusively in the process of subsequent investigations of various types of rocks and ores in which MFMA is used. Because of these maghemite inclusions and because their optical properties closely resemble those of hematite, the magnetic properties of maghemite may be attributed to hematite erroneously.

Mineragraphic studies of rocks with varying MFMA parameters made it possible to further clarify the confusing problem of the nature and character of the magnetization of rock-forming minerals. It is well known that many textbooks, monographs, and handbooks include lists of minerals giving the values of their magnetic susceptibility, natural residual magnetization, and other magnetic parameters. Such ores as ilmenite, hematite, even limonite, are described as ferromagnetic minerals (10). Almost all ferrous silicate minerals are shown as possessing a wide range of χ values varying from a few 10^{-6} to $10^{-4} - 10^{-3}$ CGS μ . Actually, however, the mentioned and all other iron-bearing silicate minerals (olivine, pyroxene, amphiboles, garnets, biotite, etc.) are paramagnetic. The ferromagnetic properties frequently discovered during the physical measurements of such minerals are attributable to the minute inclusions of magnetite and other ferromagnetic materials, whose presence, as a rule, disregarded.

Petrographic analyses which are included in the MFMA complex permit evaluations to be made of the general mineralogical composition of the investigated rocks, and to better the degree and character of the metamorphism, as well as the peculiarities of distribution and morphology of the ore formations.

In practice geophysical organizations en-

gaged in studying the physical properties of rocks are as a rule, known to be rather unconcerned about such matters as determination of elementary composition and the nomenclature of the investigated geologic formations. As a result cases frequently occur when physical properties are inconsistent with the names of the rocks. This situation has developed because of a lack of proper theoretical and methodical investigations of the natural geological and physical relationships.

The magnetic-fractional-mineralogical analysis promises to put this matter in better order, because this analysis provides for close coordination of magnetic and mineralogical studies. As the MFMA was being improved in the research process, the physical and petrographic methods of rock investigation gradually became more and more interdependent. If at the early development stages of the analysis physical and geologic characteristics were merely compared, during the later stages they were used as mutual checks, and to determine reasons for the noted inconsistencies between the magnetic and mineralogical characteristics. In most cases repeated measurements revealed erroneous determinations and the inconsistencies could thus be eliminated. Whenever the repeated observations failed to remove these inconsistencies, they served as the basis for a search for new geologic and physical relationships still unreflected in the determined regularities.

It is well known that a single-value determination of a rock's name with a microscope is not an easy task, and is often difficult, particularly if the rock is a hybrid or metasomatic formation. It is recommended that MFMA parameters be used for a more precise determination of rock compositions.

Let us now revert to the MFMA results of the rocks investigated by us. The symbols in Figure 2 designate the areas, or zones, of distribution of the rock samples depending on the magnitude of their magnetic susceptibility (χ) and parameter F_p . In order to avoid obscuring the drawing, only three lines of identical M values corresponding to 100, 1000, and 10,000 were drawn. As may be seen, each zone has a perfectly definite position within the area bounded by lines $M = 100$ and 10,000, and a very definite, though generalized, designation based on the elementary composition of the rock sample, is assembled therein.

In the middle section of the area there are three zones: hybrid granites, gabbroids, and effusive analogs of gabbroids. The granitoids have the highest values of parameter $M = 1000 - 1600$, the lowest ($M = 200 - 500$) are characteristic of basalts. The intermediate values of this parameter correspond to gabbroids.




FIGURE 2. Magnetic susceptibility of crystalline rocks in Volga-Uralian region, the Ukrainian Shield, and other areas as a function of the percentage of ferromagnetic minerals in them.

1 - zone of hybrid granitoids; 2 - zone of granodiorite-gneisses; 3 - zone of gabbroids; 4 - zone of basalts; 5 - zone of altered gabbroids; 6 - zone of biotite, biotite-amphibole-chlorite, garnetiferous biotite gneisses; 7 - zone of garnet gneisses and metasomatic skarn-type rocks; 8 - zone of amphibolites and amphibole-chlorite schists; 9 - zone of magnetite quartzites, hornfels, schists, and ores; 10 - pyrrhotite ores; 11 - hematite ores.

Is this distribution of samples of acid and basic magmatic rocks accidental? No, it conforms strictly to established law and is due to the fact that magnetite, free of inclusions, appears as the chief ferromagnetic material in granodiorites, granosyenites, gabbroids, and pyroxene granites which make up the zone of hybrid granitoids. When titanomagnetite occurs in the rocks along with magnetite the value of parameter M diminishes. Ferromagnetic minerals form large allotropic bodies occupying the intergranular spaces in the samples of hybrid granitoids. In the gabbro, olivine gabbro, and gabbro-diabases of the intermediate zone, the ferromagnetic component is titanomagnetite, i.e. magnetite in which ilmenite inclusions comprise from 10 to 50% and more of the total volume of the large primary anhedral ferro-magnetic aggregates. In addition, secondary magnetite

is present in all gabbroids — as well as ultra-basic rocks (pyroxenites and peridotites) — in large quantities as fine punctate inclusions associated with the grains of pyroxenes and amphiboles. In basalts and andesites, as well as in basaltic diabases, the ferromagnetic mineral is titanomagnetite in which the predominant component is not magnetite, but ilmenite. At the same time, ferromagnetic deposits in the basic effusives are very small in size (on the order of a few units to tens of microns). Because of this their parameter M values are the lowest.

In this manner it is possible to make a genetic classification of acid and basic magmatic rocks on the basis of parameters χ , F_p , and M . Rocks of average composition — diorites — occupy an intermediate position between the granitoids and the gabbroids.




FIGURE 3. The χ to χ_n ratio for crystalline rocks from the same regions. (Designations in this and the following diagrams are the same as in Figure 2).

In the upper right section of the diagram there is a zone of ferruginous formations. The highest parameter M values, up to 5000-6000, are manifested by ores and quartzites made up of primary magnetite deposits forming massive aggregates or ore bands. The lowest M values characterize pyrrhotite ores. The intermediate values of this parameter pertain to amphibole magnetite hornfels and schists which, in addition to primary magnetite, also contain secondary magnetite associated with silicate intercalations or minerals. Certain types of these rocks also contain titanomagnetite deposits along with those of magnetite. The samples of hematite ores represent the zones of altered gabbroids and granitoids and have lower values for parameter $M = 400-1500$.

It is known that pyrrhotite ores may possess ferromagnetic and paramagnetic properties (1). The five samples of pyrrhotite ores from different deposits in Eastern Siberia and the Kola Peninsula examined by us turned out to be ferromagnetic, but having a wide range of parameter M values equal to 300-1000. Mineragraphic investigation of the pyrrhotite ores showed that the high values of this parameter characterize magnetic pyrites containing inclusions of magnetite. Samples without magnetite inclusions had low values for parameter $M \approx 300$. Consequently, it is also possible to construct a genetic classification for ferruginous formations with the aid of parameters χ , F_p , and M .

Several zones are located in the lower left section of the diagram. These are the zones

of altered gabbroids, altered granitoids, gneisses skarn-type rocks, and amphibolites.

The zone of altered gabbroids includes gabbro-amphibolites and amphibolites; the zone of altered granitoids consists of granodiorite-gneisses; the zone of gneisses incorporates biotite, garnetiferous biotite, sillimanite-biotite, and cordierite facies; the zone of amphibolites supposedly consists of para-amphibolites, amphibole-chlorite schists and diabases. Characteristic of all these rocks is a negligible quantity (in gabbro-amphibolites), or total absence (in gneisses and amphibolites) of primary magnetite. In view of this fact, weak magnetization of these rocks is, to a considerable extent, or entirely, due to the thin intercalations of secondary magnetite. This is the reason that small values of parameter M (less than 1000) were registered for the bulk of these samples. Only in weakly magnetic samples containing very low ferromagnetic mineralization parameter M was found to be sharply increased to 5000-7000 and higher.

The zone of skarn-type formations contains concentrated metamorphic rocks enriched by garnet. Typical of these rocks is pyrrhotitic mineralization and a corresponding low value of parameter $M = 200-300$.

Figure 3 shows the distribution character of the zones from which the examined samples were taken and its relationship to the ratio of parameters χ , χ_n , and H . Since the values of one and the same parameter (susceptibility) are plotted in this and the preceding diagrams




FIGURE 4. Relationships of χ and σ values in crystalline rocks from the same regions.

Along the χ -axis, the distinctions and the community of the zone configurations and dimensions are determined by the ratio of parameters F_p and χn .

In Figure 3 all the zones are located in the lower right half of the diagram between the diagonal for $H = 100\%$ and the line of $H = 0.1\%$. As can be seen, the lowest values of parameter H pertain to the rocks containing the smallest relative quantity of secondary magnetite (magnetite ores, quartzites and hornfels, hybrid granites, gabbro-diabases). The highest values characterize metamorphic rocks containing negligible concentrations of dispersed secondary magnetite, effusive analogs of gabbroids containing small segregates of primary ferromagnetic material (titanomagnetite), and the hematite ores containing relict magnetite. It is readily noticeable that the zones of gabbroids, granitoids, and iron-ore formations occupy a larger area in Figure 3 than in Figure 2, and the zone of granodiorite-gneisses appears at the same time to be even more strictly confined to its proper location between the hybrid granites and the gneisses.

Thus, a graphic construction clearly illustrates the fact that parameters χn and H characterize not a random but the basic properties of rocks predetermined by their origin and metamorphism. If the value of parameter M is affected most by the composition and quality of the ferromagnetic material (concentration and character of deposition of paramagnetic minerals), then the value of parameter H depends in the first instance on the size of the ferromagnetic grains and the manner of aggregation of the ferromagnetic components with the rock-forming minerals. Parameter H is higher, as the size of the magnetite deposits decreases and as the grouping of these deposits with the silicate minerals is closer. Such a picture is observable, in particular, in basalts and basaltic diabases.

Distinctions in rock characteristics in terms of parameter M and H values also may be illustrated by the difference in the configuration and position of the zone of skarn-type rocks. Thus, if in Figure 2 this zone occupies the extreme left position and is elongated along the line of $M = 200-400$, in Figure 3 this zone is transverse with respect to the H lines over a




FIGURE 5. Relationship of χ and J_n values in crystalline rocks from the same regions.

range of H values from 100% to ~ 20%. The different location of this zone is due to the fact that in Figure 2, parameter M mainly expressed the composition of ferromagnetite minerals (pyrrhotite) whereas in Figure 3 parameter H expresses the relation between the coarse-grained (pyrrhotite) and the fine-grained (secondary magnetite) ferromagnetic bodies. On the basis of the range of parameter H values it is possible to distinguish rocks with distinctly pronounced pyrrhotitic mineralization accompanied or unaccompanied by secondary magnetitic mineralization. Analogous relationships between primary and secondary ferromagnetic metallization may be observed in comparing the other zones indicated in the graphs.

Since the ferromagnetic composition (magnetite, titanomagnetite, magnetic pyrites, etc.) is mainly determined by the content and origin of the rocks containing them (magnetite for granitoids, titanomagnetite for gabbroids, pyrrhotite for several types of metasomatic rocks, etc.), parameter M can be utilized as an objective criterion for the genetic classification of rocks.

Parameter H, which reacts discriminately to the ratio of large and fine ferromagnetic segregations resulting from secondary geochemical transformations, may be used as an objective

criterion in determining the nature and degree of rock metamorphism.

The outlined regularity in M and H variations in rocks having different composition, origin, and metamorphism permits us to express the conviction that not only each type, but each variety of crystalline rocks will be given a perfectly definite and specific magnetic and mineralogical characteristic as a result of subsequent detailed MFMA analyses of geological formations.

MFMA parameters are capable of providing a precise definition of the orientation and depth of the metamorphic alterations of ortho- and para-rocks constitutionally similar but differing in origin and metamorphism. For this purpose it is necessary that investigations be continued to provide a more definite specification of the relationships between MFMA parameters and the composition of ferromagnetic segregates with due consideration of their morphology in rocks of different origin having distinctly expressed manifestations of such processes as opacitization, amphibolization, serpentization, and biotitization of ferruginous minerals which are responsible for the generation of a secondary and definite transformation of primary magnetite deposits.

In order to attain these ends use could be made

not only of magnetic parameters but also of density values. Figure 4 shows the relationships between parameters χ and σ with respect to the same rocks and zones as indicated in Figures 2 and 3.

It is noteworthy that the zones of concentration of rocks having specific composition, metamorphism, and genesis as outlined on the basis of magnetic characteristics have not only lost nothing in terms of specificity and definiteness of graphic expression with the introduction of density characteristics, but have, on the contrary, gained in overall vividness in certain cases. If desired it is possible, for example, to isolate the granite-gneisses and acid migmatites possessing the lowest density values ($\sigma = 2.5 - 2.65$) in the gneiss zone. The highest density ($\sigma = 2.70$ to 2.85) is in the gneisses of biotite-sillimanite and biotite-garnetiferous compositions. Biotite gneisses and diorite-gneisses occupy an intermediate position. Densities, clearly distinguishable from those of other varieties of hybrid granitoids are the grano-syenites ($\sigma = 2.5$ to 2.6). The most highly differentiated in terms of density and susceptibility are observable in iron-ore formations. If, on the basis of density values, one can distinguish the magnetitic ores, magnetitichornfels, magnetite quartzites, and magnetite schists, then, from the magnitude of susceptibility it becomes possible to isolate the magnetite, pyrrhotite, hematite-magnetite, magnetite-hematite, and hematite minerals.

A comparison of Figures 2, 3, and 4 shows that ferromagnetic mineralization is characterized by the basic MFMA magnetic parameters and is regularly related to the overall mineralogical content of the investigated rocks, expressed in density values. The ferruginous formations having magnetitic and pyrrhotitic composition have the highest ferromagnetic mineralization. As ferromagnetic concentration diminishes, the iron-ore formations retaining their high density values ($\sigma > 3$) gradually acquire a hematitic aspect.

Among igneous rocks the highest ferromagnetic mineralization occurs in olivine gabbro, gabbro-diabases, basalts, andesites, pyroxenites, diorites, and hybrid granites. All of these rocks, depending on their composition, contain predominantly primary magnetite or titanomagnetite, average-composition rocks - magnetite and titanomagnetite. Secondary magnetitic mineralization is of no significance in granitoids. In gabbroids it represents but a negligible fraction of the primary magnetite, but in pyroxenites and other ultrabasic rocks may play a substantial role.

In rocks subjected to various stages of metamorphism, primary ferromagnetic metallization gradually loses its importance

in giving way to secondary mineralization. In such highly altered and relatively heavy rocks as biotite-garnetiferous and sillimanite-garnetiferous gneisses or amphibolites, only secondary magnetite is to be found, and sometimes pyrrhotite. The presence of maghemite, in addition to secondary magnetite, is possible in certain varieties of gabbro-amphibolites. Consequently, ferromagnetic mineralization reflects the specific conditions of formation and subsequent mineralogical transformations of rocks.

MFMA parameters characterize the individual manifestations and the aggregate totality of ferromagnetic mineralization of rocks and constitute the new reliable criteria of the magnetic and mineralogical state of the substance. Their significance in the development of geological science is obvious, and their practical utilization absolutely indispensable.

In conclusion, we shall discuss Figure 5 in which the investigation of rocks is based on parameters J_n , χ and Q . This construction shows that natural residual magnetization is also related in a definite manner to element composition, i.e. to a definite composition and structural peculiarities of ferromagnetic minerals. In the general case parameter J_n depends on ferromagnetic concentrations: rocks enriched by ferromagnetic minerals possess higher values of residual magnetization. However, this general postulation calls for certain qualifying specifications. For example, rocks with different compositions - even if the ferromagnetic concentrations in them are identical - may have sharply differing values of parameter J_n and a wide range of parameter Q variations (from 0.02 to 20).

The highest values of parameter Q are displayed by magnetite ores having columnar structure and containing maghemite, olivine gabbro, and hypersthene gabbro-amphibolite (amphibolized norite) with oriented mineralization of secondary magnetite. Low values of this parameter are observable in oxidized varieties of constitutionally different rocks, including the martinitized types of basalt found in the Chernigov Boring - the youngest of all rocks examined by us (Devonian). Intermediate values of this parameter typify the hybrid granites, diorites, diorites and gabbro-diorites. Therefore, depending on the composition and content of ferromagnetic minerals, rocks may possess various values of parameter Q .

In speaking of the state of ferromagnetic minerals we mean not only the degree of oxidation but also the character of inclusions. Analysis of polished sections of samples having different values for parameters J_n and Q indicates that the samples containing titanomagnetite with decay structure in the form of a very

dense lattice of very thin ilmenite plates are characterized by higher values of the above parameters as compared to samples containing pure magnetite or titanomagnetite with ilmenite inclusions of irregular form, or in the shape of short and rare plates. It is obvious that parameters J_n and Q also may be utilized for rock characterization.

Thus, MFMA provides an opportunity to discover new, and to define more precisely, already known geological and physical regularities and relationships. As a result of investigations of various types of rocks applying this analysis, a scientifically and practically important postulation is being borne out: ferromagnetic mineralization is regularly related to general mineralization and reflects the peculiarities of origin and transformation of rocks. Rocks of varying composition, genesis, and metamorphism show that their ferromagnetic mineralization has specific characteristics, which are reflected in the variety of compositions, sizes, forms, and in the character of associations of the primary ferromagnetic minerals; in the presence or absence in them of inclusions of paramagnetic minerals varying by composition, size, and shape; in the presence or absence of secondary ferromagnetic mineralization; and in the diversity of its dimensions, shape, and association with different silicate minerals.

MFMA parameters react to the individual or composite manifestations of ferromagnetic mineralization and facilitate characterization of the elementary composition of rocks with consideration for their genetic and metamorphic peculiarities.

The results of rock investigations by means of MFMA open new opportunities for practical utilization of magnetic parameters. The obsolete idea that magnetic properties are of interest allegedly only to those geophysicists and geologists who deal with interpretations of magnetic surveys or exploration of iron deposits should be cast away. As shown above, the magnetic properties should interest, in a very practical way, all geologists, mineralogists, geochemists, and geophysicists engaged in studying the elementary composition of geologic formations.

In particular, MFMA may be recommended for purposes of classification of iron ore deposits, genetic classification of igneous and metamorphic rocks, for geologic mapping of crystalline rocks, metallogenic studies of individual areas and major regions, for elaboration of theoretical and methodological problems associated with the geologic interpretation of geophysical data and studies of magnetization of geologic bodies and minerals, as well as of methods of complex regional

geologic and geophysical investigations, for investigations of terrigenous strata containing beds with different but sufficiently intensive ferromagnetic mineralization.

The MFMA method should play a leading role in the work of laboratories dealing with systematic studies of physical properties and elementary compositions of rocks and ores.

REFERENCES

1. Yeliseyev, E. N., and A. P. Denisov, Rentgenometricheskoye issledovaniye pirrotina. [X-RAY EXAMINATION OF PYRRHOTITE]: Vestn. Leningr. un-ta, ser. geol. i geograf., No. 18, vyp. 3, 1957.
2. Yefimov, F. N., Magnitno-fraktsionno-mineralogicheskiy analiz. [MAGNETIC-FRACTIONAL-MINERALOGICAL ANALYSIS]: Geologiya nefi, No. 1, 1958.
3. Malyshev, I. I., Zakonomernosti obrazovaniya i razmeshcheniya mestorozhdeniy titanovykh rud. [MECHANISM OF FORMATION AND EMPLACEMENT OF TITANIUM ORE DEPOSITS]: Gostoptekhizdat, 1957.
4. Nagata, T., Magnetizm gornyykh porod. [MAGNETISM OF ROCKS]: Izd-vo IL, 1956.
5. Ovchinnikov, L. N., O maggemite. [ON MAGHEMITE]: Tr. Gorno-geol. in-ta, Ural'skiy fil. Akad. Nauk SSSR, vyp. 20, 1953.
6. Ovchinnikov, L. N., Nekotoryye voprosy magmatogennogo rudoobrazovaniya. [CERTAIN PROBLEMS OF MAGMA ORE FORMATION]: Izv. Akad. Nauk SSSR, ser. geol., No. 4, 1959.
7. Pavlov, N. V., and I. I. Chuprynina, O magnomagnetitakh kak indikatorakh glubinnosti orudeneniya. [MAGNESIO-FERRITES AS INDICATORS OF THE DEPTH OF METALLIZATION]: Dokl. Akad. Nauk SSSR, t. 104, No. 2, 1955.
8. Pavlov, N. V., O maggemite v magnetitovykh rudakh Kezhemskogo mestorozhdeniya. [THE MAGHEMITE AND MAGNETITIC ORES OF THE KEZHMA DEPOSITS]: Tr. Mineralog. muzeya Akad. Nauk SSSR, vyp. 8, 1957.
9. Sal'dau, E. P., Izmeneniye razmerov yacheyki pri okislenii iotsita v magnetit i maggemit. [ALTERATION OF THE CELL DIMENSIONS AS A RESULT OF JOSEITE OXIDATION INTO MAGNETITE]

AND MAGHEMITE]: Zap. Vses. mineralog. o-va, ch. 86, No. 3, 1957.

10. Khramov, A. N., Paleomagnitnaya korrelyatsiya osadochnykh tolshch. [PALEOMAGNETIC CORRELATION OF SEDIMENTARY STRATA]: Tr. Vses. n. -i. geol. -razv. in-ta, vyp. 116, 1958.

All-Union Scientific Research Institute
Petroleum Exploration
(VNIGNI)
of the Ministry of Geology and
Conservation of National Resources
of the U. S. S. R.,
Moscow

Received, June 6, 1960

THE TECTONICS OF THE NORTHERN PARTS OF PATOMSK UPLAND¹

by
S. V. Ruzhentsev, and Chang Bu-Chung

The geologic structure of the Patomsk upland long ago attracted the attention of geologists. It is mainly under the influence of V. A. Obruchev's writings that a theory has been developed concerning the presence of a smooth arc of folding within the northern confines of the upland. M. M. Tetyaev (15) explained the bend in the fold structure by the presence of a major anticlinorium which pitched in a northeasterly direction and was responsible for the prevalence of northeasterly trends which determine the closure of the folds. A criticism of M. M. Tetyaev's views will be found in Ye. V. Pavlovsky's paper [12], in which it was shown Tetyaev's constructions are an artificial combination of a variety of Precambrian and Lower Paleozoic structures. Considerable attention is devoted to the northern outskirts of the Patomsk upland in N. M. Chumakov's works [17, 18], in which the Patomsk folded zone is shown to be complicated by two transverse uplifts (Urinskiy geanticline and Modinskaya anticline) which are contemporaneous with the Patomsk zone structures.

Up to this time there is still a considerable divergence of opinion as to the nature of the historical development and mechanism of formation of the geologic structures observable on the outskirts of the Patomsk upland. L. I. Salop [14] identifies the Proterozoic Baykal-Patomsk foredeep as separated from the inner Baykal geosyncline by the Baykal-Vitim uplift which, in the Upper-Proterozoic, was an erosion area from which detrital materials were transported into the adjacent troughs. A generally similar view is held by N. M. Chumakov [17], who distinguishes the Priбайkal foredeep as being at the junction of the Baykal geosyncline and the Siberian platform.

N. S. Zaytsev [7] is inclined to consider only the Precambrian trough as a foredeep, relating the Lower Cambrian deposits to a

special relict structure. In the opinion of P. Ye. Offman [11], the Angara-Lena trough is a system of two strictly platform-type synclises. In advancing his concept of a "pericraton subsidence zone", Ye. V. Pavlovskiy has recently given an original evaluation of the Angara-Lena trough. The specifics of such a zone consist, first of all, of an exceedingly long period of development lasting for many geologic periods.

The stratigraphy of the Precambrian and Lower Paleozoic deposits in the northern and northeastern limits of the Patomsk upland is sufficiently well studied [1-6, 8, 9, 12, 14, 18]. Consequently, we shall restrict ourselves to a brief enumeration of the series (starting from the top).

I. The Patomsk complex (Upper Proterozoic, Riphean, Sinian): Ballaganakhian, Mariinskan, Bol'shepatomskian, Barakunian, Valyukhtinian series, Kullekinian horizon, Nikol'skian and Alyanch-Kholychian series.

II. Lower Cambrian: Zherbinian, Tinnovian varicolored series (Aldanian stage): El'gyanian, Tolbachanian, Olekminian, Charskian series (Lena stage); Upper Cambrian: Verkholenian series.

III. Ordovician - Lower Silurian: Ust'kutian series (0); Krivolukian and Makarovskian series (0₂₋₃); Meikian series (S₁).

From data obtained during the 1958-1959 operations we shall try to give below the development characteristic of the region's principal structures for the different stages of its geologic history.

1. Patomsk and Urinsk Upper Proterozoic (Riphean and Sinian) Troughs.

The thick carbonate-terrestrial deposits in the lower and middle parts of the Patomsk complex accumulated along the northern offshoots of the Baykal-Patomsk upland where they filled the large Patomsk trough which is traceable from the basin of Zhuya River to the

¹О Тектонике северных частей Патомского Нагорья. pp. 37-48.




FIGURE 1. Sketch of the location of the principal Upper Proterozoic structures in the northern fringe of the Patomsk trough.

1 - uplifted erosion zones, 2 - accumulation zone of the thick carbonate-terrestrial strata of the Patomsk complex.

middle course of the Lena. Another trough (Mariinsk), approximately perpendicular to the former, extends into the basin of the Dzherba and Ura rivers from the southwest to the northeast (Figure 1). Both troughs are filled with Upper Proterozoic deposits (Ballaganakh-Tn, Mariinsk, Bol'shepatomskian, Barakunian, and Valyukhtinian series).

In the deposits of the investigated series we identify two types of sections (Figure 2): the northern (lower courses of the Mal. and Bol. Patom, Ura, and Lena Rivers), and the southern (upper reaches of the Mal. Patom River). An extensive development of coarse terrestrial deposits including powerful boulder-conglomerate series (Bol'shepatomskian series), characteristic of the northern sections of the Mariinskian and Bol'shepatomskian series. In the upper-course basin of the Mal. Patom (southern series) predominant formations are carbonate (Mariinskian series) or carbonate-argillaceous deposits. The distinctions between the northern and southern sections were heavily eroded during the period of deposition of the Barakunian and Valyukhtinian series. In the lower reaches of Mal. Patom along the Ura and Dzherbedyanka Rivers, carbonate and carbonate-argillaceous deposits interstratified with quartz- and quartz-feldspathic sandstones and thick beds of underwater slump breccia are developed. Prevalent in the south (Mal. Patom River headwaters) are the same carbonate and carbonate-argillaceous deposits.

Here, a reduction of sandy material is noted in a number of horizons (the middle section of Barakunian series), and there is no breccia.

It is obvious, therefore, that there was a variation in facies in the deposits of the lower and middle parts of the Upper Proterozoic in the Patom complex. This variation is expressed by the occurrence of the coarsest sediments in the northern part of the trough adjacent to the platform, whereas the fine sediments appear in the southern inner part of the trough.

A conclusion may be drawn to the effect that the clastic material which fills the Patomsk trough came from the north, i.e. from the intensively eroded platform regions. The finer sediments, however, are exposed in the upper reaches of Mal. Patom River, closer to the Baykal-Vitim uplift (according to L.I. Salop). Thus, the position taken by this author concerning the erosion of the uplift, which in his opinion represented the source of the terrigenous material filling the Patomsk trough, appears to be unsupported.

The trough under consideration has a number of characteristic features: linearity, and a specific position between the platform and the inner parts of the Baykal geosyncline characterized by a widespread distribution of volcanic rocks. In approaching the platform, the terrigenous material becomes even




FIGURE 2. Facies profile across the Patomsk Trough.

I - Ballaganakhian series, II - Mariinskian series, III - Bol'shepatomskian series, IV - Barakunian series; V - Valyukhtinian series, VI - Kullekian horizon; VII - Nikol'skian series; VIII - Alyanch-Kholychian series. 1 - conglomerates, 2 - sandstones, 3 - clay shales and cherty carbonaceous schists, 4 - limestones, 5 - dolomites, 6 - marls, 7 - carbonate and carbonate-argillaceous slump-breccia, 8 - tuffs. Vertical scale - 1:100,000.

coarser and increases in volume toward the edge of the platform which is the obvious source area. These features, and an almost total lack of effusive formations, permits us to consider the Upper Proterozoic Patomsk trough to be a typical miogeosyncline.

The deposits filling the Urinski trough differ almost not at all from those in the Patomsk miogeosyncline. This makes it possible to regard it as a peculiar offshoot of this miogeosyncline projected onto the platform. The first to analyze the morphological characteristics of similar structures was N.S. Shatskiy [20]. He has identified the transverse marginal systems genetically and structurally most closely associated with geosynclinal depressions.

Toward the end of the Valyukhtinian series sedimentation, the troughs widened and their shapes were reformed. The nature of sedimentation in them was changed, and as a result, a new independent structure can be identified: the Nyuya trough.

2. The Lower Paleozoic Nyuya Trough

The trough in question today consists of a system of two basins: the Berezovsk and Nyuya-Dzherba basins separated by the Urinsk uplift. Both basins are filled with a thick (4300-4400 m) complex of deposits forming the Nikol'skian

and Alyanch-Kholychian series belonging to Late Precambrian, Lower and Upper Cambrian Lower Silurian. More than 3000 m of this formation consists of carbonate strata including the Nikol'skian and Alyanch-Kholychian series and Lower Cambrian deposits.

Of great interest is the distribution of thicknesses and facies of Upper Cambrian, Ordovician and Silurian deposits in the Berezovsk basin and the Urinsk uplift (Figure 3). The thickness of Verkholenian series along the lower course of the Bol. Patom River amounts to 350 m according to N.M. Chumakov's data [18], and 314-322 m in the central part of the trough, according to R.F. Gugol' [2]. The uniform thickness of the Ordovician formations is striking. For example, the Ust'kut series in the central section of the trough reaches 220 m in thickness [2], but around the western limb of the Urinsk uplift it is 205 to 220 m thick. The thickness of the Krivolutsian series is 178 m in the first case, and 210-240 m in the second.

The Verkholenian series in the central part of the Berezovsk basin, according to R.F. Gugol' is represented by a stratum of reddish-brown and greenish clay marls containing subordinate intercalated dolomites and limestones. Bands of coarse-grained siltstones are also encountered. According to the data of a recent survey (G.S. Borushko, M.L. Kokaulin), the sections of the Verkholenian series on the western and eastern




FIGURE 3. Facies profile of the Upper Cambrian and Ordovician deposits of the Nyuya-Dzherba and Berezovsk basins.

Horizontal scale - 1:2,000,000,
vertical scale - 1:5,000.

I - Upper-Cambrian, II - Lower Ordovician, III - Middle Upper Ordovician; 1 - sandstones, siltstones; 2 - argillites, 3 - dolomites, limestones, 4 - marls.

slopes of the Urinsk uplift are basically similar. Predominantly developed here are clay marls, argillites, dolomites and limestones, while terrigenous rocks are present in very negligible quantities.

The Ust'kut series in the central part of Berezovsk basin, according to N.S. Zaytsev and N.V. Pokrovskaya's data [6], primarily consists of dolomitized sandy limestones, alternating with polymictic sandstones, siltstones, argillites, and marls. On the eastern limb of the Urinsk uplift the Ust'kutian deposits are represented by dolomitized, sometimes sandy, algal limestones interstratified with argillites and marls. The sandstones occupy a strictly subordinate position. A generally analogous sequence occurs on the western slope of the uplift, the only difference being that sandstones are absent here altogether.

Thus, we observe that in approaching the Urinsk uplift, whose amplitude exceeds 8 km, from the central part of Berezovsk basin we find almost no facies changes. In certain series (particularly well manifested in the Ust'kut, and to a lesser degree, in that of the Verkholenian series) one notes the smaller role played by terrigenous material as one draws closer to this uplift.

The deduction is obvious. In the Lower Silurian, the Nyuya trough constitutes a major single structure free of internal uplifts which

would have separated it into individual basins.

It is interesting to trace the mode of transition between the deposits in the lower and middle parts of the Patomsk complex, on the one hand, and the upper section of the Patomsk complex and the Lower Cambrian formations, on the other. As studies by V.N. Makhayev, L.I. Salop, V.K. Golovenko, and N.M. Chumakov showed, the Valyukhtian schists are related by gradual transition to the Nikol'skian series, whereas the deposits of the Alyanch-Kholychian series are related to the Zherbinian sandstone series of the Lower Cambrian. Under these circumstances, the only method which we could use successfully in our attempt to characterize the Nyuya trough through the various stages of its existence, is the formational method.

During the time when the Upper Proterozoic Patomsk miogeosyncline was in existence the Aldan shield was uplifted and served as an erosion zone. It was not until the end of the Valyukhtinian age that it was partially covered by the sea and a thin bed of quartz and quartz-feldspathic sandstones containing individual intercalations of dolomite was deposited within its limits [5].

The following three formations have been identified by us in the Patomsk miogeosyncline: the lower formation (Ballaganakhian and Mariinian series), composed of alternating quartz and quartz-feldspathic sandstones,

clay shales and, less frequently, limestones; the middle formation (Bol'shepatomskian series) made up by conglomerates, sandstones, cherty coals, and clay shales; and the upper formation (Barakunian and Valyukhtinian series) represented by interbanded limestones, dolomites, clay shales, quartz sandstones and thick beds of carbonate and carbonate-clay breccias.

By the end of Upper Proterozoic (Nikol'skian and Alyanch-Kholychian series) a diversified 900-1200-m complex of deposits had accumulated in the northern and northeastern outskirts of the Patomsk uplift. The chief characteristic of the strata under discussion is the marked prevalence of various carbonate rocks, including marls, dolomitic marls, fine-grained — sometimes algal and oolitic, sometimes clastic — limestones alternating with sandy limestones. These deposits are classified by us as the lower carbonate formation of the Nyuya trough.

The deposits of Zherbinian series, represented by uniform strata of quartz sandstones containing individual bands of quartz siltstones, are identified by us as the terrigenous formation of the Nyuya trough. The deposits of this formation are 250-300 m thick and, apparently, already resemble a platform formation.

Higher in the column there is a diversified complex of light-colored, aphanitic limestones, limestone shales, dolomites, dark bituminous limestones, interbanded with argillites. This complex we also classify as a terrigenous-carbonate formation of the Nyuya trough (it corresponds to the Lower Cambrian Tinnovian series). The thickness of this formation is about 500 m.

The upper terrigenous carbonate formations in the Nyuya trough correspond to the Lena stage deposits (El'gyanian, Tolbachanian, Olekminian and Charskian series). As a whole, it is distinguished by its considerable facies uniformity and is represented by brown, dark-gray limestones, often oncolitic or stromatolitic, alternating with light-colored dolomites. The formation attains a thickness of 1000-1100 m.

Thus, in the period of the Patomsk miogeosyncline's existence, carbonate-terrigenous strata of great thickness were deposited under conditions of intense downwarping. The detrital material was supplied from platform regions which were subjected to intense erosion.

Beginning with the end of Upper Proterozoic, new formations appeared (the lower carbonate, terrigene, and terrigene-carbonate) which marked the disappearance of the Patomsk miogeosyncline and the development of an

independent trough (Nyuya). This depression is filled with a thick (up to 3000 m) series of mostly carbonate rocks among which terrigenous and argillaceous formations sometimes played a substantial role. The deposits of the lower carbonate, terrigeneous, and terrigeneous-carbonate formations have many features in common with the Patomsk miogeosyncline deposits (great thicknesses, variegated lithological composition, etc.). Therefore, their identification as formations characteristic for the independent trough must be considered as rather conditional and mainly due to their intermediate position between the platform rocks (variegated and carbonate formations) and the miogeosyncline deposits.

A widening of the trough took place during the same period, with the resultant downwarping involving vast areas of the Aldan shield where a relatively thin (400-500 m) dolomite series was formed. Both in terms of thickness of deposits and in composition, the rocks of both formations are drastically different.

The Lower Cambrian variegated formation in the Nyuya trough still indicates that there must have been considerable downwarping in this depression where the deposits are generally twice to three times as thick as those of the Aldan shield. At the same time, one should note a certain leveling out of structural distinctions between the variegated and the older formations.

A still greater lessening of the differences between the deposits of both tectonic zones must have taken place while the upper carbonate formations were being deposited at the end of the Lower Cambrian epoch. The upper carbonate formation in the Nyuya trough acquired the features of typical platform structures. The Nyuya trough area and the Aldan shield differed little from one another at that time as far as tectonic conditions were concerned. The difference was only in the greater degree of warping in the Nyuya depression. As a result, the Nyuya deposits are one- and one-half times thicker than those of the Aldan shield.

On the basis of the enumerated data, the general sequence of changes affecting the formations can be formulated as follows. There was a gradual lessening of the distinctions between the formations of the Patomsk miogeosyncline and the inherited Nyuya trough, on the one hand, and those of the Aldan shield, on the other. This happened against a background of regeneration of the Patomsk miogeosyncline into a special structure of platform type.

Thus, the development of the investigated region in the Riphean (Upper Proterozoic) and Lower Cambrian can be reduced to the



FIGURE 4. Schematic geologic map of the northern fringe of the Patomsk Upland (compiled from materials produced by G.S. Borushko, V.K. Golovenko, L.L. Kokaulin, S.V. Ruzhetsk, Chang-Bu-Chung, N. M. Chumakov).

1 - Ballaganakhian, Mariinskian, Bol'shepatomian series; 2 - Barakunian series; 3 - Valyukhtinian series; 4 - Nikol'skian, Alyanch-Kholychian series; 5 - Zherbinian series; 6 - Tinnovian and the variegated series; 7 - Lenian stage; 8 - Upper Cambrian, Ordovician, Lower Silurian; 9 - Deso-Cenozoic deposits; 10 - ruptured zones.

formation of the Patomsk miogeosyncline which, starting at the end of Upper Proterozoic, gradually but rapidly, changed into a peculiar syncline, marginal with respect to the Baykal geosyncline.

In the light of this discussion it is apparent that the zone of pericratonall subsidence transcends the concept of the Angaro-Lena depression as argued by Ye. V. Pavlovskiy.

It is true that the Upper Proterozoic Patomsk complex is structurally inseparable from the overlying sedimentary series of the entire Lower Paleozoic. Nevertheless, there exists, in our opinion, a sufficiently sharp boundary between the major portion of the Patomsk complex and the Nikol'skian series. This is manifested in the sequence of formations, as well as in the reconstruction of the shape of the Lower Cambrian trough as compared to the Upper Pro-

terozoic trough. A sufficiently distinct structural boundary also can be traced along the border between the Lower and Upper Cambrian.

The following sequence of formations is characteristic of the formation peculiarities of the various Upper-Proterozoic - Lower-Paleozoic stratigraphic complexes. The lower and middle beds of the Patomsk complex represent the strata of the Patomsk miogeosyncline. The late Precambrian deposits of the Nikol'skian and Alyanch-Kholychian series, and the Lower Cambrian deposits are, apparently, already formations of the platform type. However, judging by their thickness they are not comparable to typical platform-type formations. Finally, the Upper Cambrian (Lower Silurian) deposits are to be classed as typical platform formations. Thus, two historical stages can be specified for the Upper Proterozoic - Lower Paleozoic and the boundary between them is distinct in all respects. These periods are the period of the Patomsk miogeosynclines existence and the period of the Nyuya trough formation. Structurally, the Nyuya trough is far from being uniform, and the history of its development must, in turn, be divided into two stages: the Lower Cambrian and the Upper Cambrian - Lower Silurian. The discussed material is a good illustration for the theory that the zone of pericratonall subsidences is a complex of distinctly pronounced depressions which have a similar structural plan and which were formed as a result of inherited and protracted development from structures of the geosynclinal type to structures of strictly platform type.

3. The Folded Zone of the Baykal Caledonides.

Sedimentation in the Nyuya trough came to an end at the boundary between the Lower and Upper Silurian. By that time there appeared along the northern fringe of the Patomsk upland a wide zone of linear folds which folded the Precambrian and Upper Paleozoic (including the Lower Silurian) formations. There is considerable literature dealing with the age of these dislocations and we shall, therefore, refrain from discussing this problem in greater detail. We shall merely point out that the theory concerning the Caledonian age of the structures appears preferable to us since there is ample evidence pointing to the conformity of Lower and Upper Cambrian deposits [3, 4, 8, 9, 12, 16] and no proof - other than the isolated case described by Ya. A. Shalek [19] - of any major discordances at the Lower-Upper Cambrian contact. In a few instances, an unconformity was noted only with respect to the overlying Verkhnelenian deposits on the beds of the Charskian series [10, 17], but the displacements here are very insignificant.

It is far more difficult to pass a more accurate judgement on the precise time of origin




FIGURE 5. Geologic diagram of the middle course of the Mal. Patom basin.

1 - Bol'shepatomskian series; 2 - Barakunian series; 3 - Valykhinian series; 4 - Nikol'skian series; 5 - Alyanch-Kholychian series; 6 - Zherbinian series; 7 - Tinnovian series; 8 - variegated series; 9 - Lenian stage; 10 - Jurassic.

for the linear folds. The fact is, that it has been definitely established in a number of places in the northern outskirts of the Patomsk upland that sheets, comparable with the traps of the Tungus basin synclines, have been dislocated in perfect conformity with the country rocks. This indicates in a way that the northern marginal structures are younger. For the time being, there are no data affording a single-value solution of the age problem with respect to the formation here of linear dislocations. In view of this fact we have chosen, in this work, to hold to the generally accepted view about their Caledonian origin. However, in the light of the foregoing discussions it appears to be entirely possible to assign these dislocations to the Upper Paleozoic. Such a point of view was actually expressed in the works of D.K. Zegerbart and Z.M. Starostina [8] and Ye.V. Pavlovskiy [12].

4. The Urinsk Hercynian Uplift.

The Urinsk uplift extends from the lower course of the Bol. Patom River for about 100 km in a northeasterly direction. In the upper reaches of the Ura basin, it is covered by the Mesozoic deposits of the Vilyuy syncline (Figure 4).

The uplift under consideration represents a system of simple, usually box-shaped anti-

clines, and flat synclines, often almost equidimensional in plan. The most commonly occurring structural types are anticlines with flat and wide arches and flexure-shaped limbs, and synclines with slightly warped bottoms. Narrow flexures having steeply dipping common limbs, separate wide areas with gently sloping layers — this is the typical feature of transverse uplift. Similar structures are widespread in the marginal parts of the Berezovsk and Nyuya-Dzherba troughs (Turuktinsk, Nyuya, Solyankinsk, Khadar, etc.). A study of these flexures helps to clarify the mechanism of structural formation in the Urinsk uplift.

Many cases have been noted where these flexures either turn into major faults along the strike (Dzherbin-Patom flexure — Kuduktinsk fault; Sulyankinsk flexure — Machinsk fault), or are complicated by minor ruptured zones having small amplitudes.

A conclusion may, therefore, be drawn to the effect that the flexure bends are associated with the ruptured zones in the closest possible manner. It is obvious, that such flexures must be considered as fractures transformed in the sedimentary mantle. The planes of these fractures are sometimes exposed at the present surface. The tops of the anticlines and the trough bottoms constitute more or less large



FIGURE 6. Geologic sketch of the Kudutka River basin.

1 - Valyukhtinian series, 2 - Nikol'skian series;
3 - Alyanch-Kholychian series; 4 - Zherbinian
series; 5 - Tinnovian series, 6 - Variegated
series; 7 - Lenian stage; 8 - traps.

blocks located at different elevations and limited by faults.

Of special interest is the intersection node of the Patomsk folded zone and the Urinsk uplift. The extreme southern structure of the Urinsk uplift is the Zhedaysk anticline. In the Mal. Patom's lower-course basin the linear folds of the Patomsk zone (the Kanskaya, Kurachanskaya, and Chepelyakhskaya folds) join almost at a right angle against the anticline's eastern limb. At the point of their junction with the Zhedaysk anticline, the eastern limb of the latter is disrupted by small-amplitude synclines. The following regularity is observed in the distribution of the Zhedaysk anticline's eastern limb structures: the narrow synclines which complicates the limb of the Zhedaysk fold are located along the strike of the Patomsk zone synclines, whereas the large wide noses revealing the ancient deposits of the Patomsk complex correspond to the anticlines of this zone.

By way of an example let us consider the Kanskaya fold which is an asymmetrical syncline, 5 - 6 km wide, extending for tens of kilometers and similar in every respect to the typical Patomsk zone structures. Its shape changes sharply only in the basin of Dolganakh stream, where its centroclinal terminus appears to be truncated (Figure 5), whereas in plan it has the outline of a somewhat deformed rectangle. Further west of

the stream valley in continuation of the Kanskaya syncline there is the narrow (about 1.5 km wide) minor-amplitude Lower Orlakon syncline.

The mode of articulation of the Lower Orlakon and Kanskaya structures is interesting. Essentially the Lower Orlakon syncline is only a narrow and very sharp constriction of the Kanskaya fold. It looks as though a sharp closure in the Kanskaya syncline occurs in the basin of the Dolganakh. Consequently, if one is to consider both folds as a unit, their structure can be represented as a system of two synclines: western (small amplitude) and the eastern (a far greater amplitude), separated by a sharp bend of layers (40° - 60° dip to the east-southeast). The azimuth of this slope coincides with the axis of the Kanskaya syncline. The above bend is typical not only for the region being discussed. It has been traced to the north along the Mal. Patom valley and is identified by us as the Malo-Patom flexure, along which the western end of the Kanskaya syncline was shifted upward causing the formation of the Lower Orlakon fold.

Another, no less interesting, transition area from the zone of linear folds to the structures of the Urinsk uplift is the basin of the Kudutka River. Here one distinctly observes the meridionally striking fault (Kudutka) which cuts off the Kudutka left-bank linear folds and bounds, on the west, the series of brachy-structures generally oriented parallel to this

fault (Figure 6). The Kudukta fault is traceable for only 25-30 km. It passes northward from the Bol. Patom River valley into the already discussed Dzherbinsk-Patomsk flexure which bounds the Urinsk uplift on the west. West of the Kudukta fault Lenian-stage deposits and those of the variegated series are prevalent. To the east occur the Alyanch-Kholychian, Zherbinian and Tinnovian series, i. e., the zone of brachy-structures was uplifted (approximately 1000 m) along the fault relative to the zone of linear folds.

Deformation and truncation of the linear folds is also observable south of the Zhedaysk anticline (Kovalinskaya anticline, the Upper Orlakon syncline).

Analysis of the cited examples makes it apparent that the transformation of the original shape of the Patomsk zone linear folds is associated with the Urinsk uplift: 1) the Patomsk zone linear folds and the Urinsk uplift are structures of different ages; 2) the Urinsk uplift was formed later than the linear folds of the Patomsk zone.

The peculiarity of the geologic structure in the basins of the lower reaches of the Bol. and Mal. Patom Rivers, and of the Ura and Dzherba Rivers consists in the widespread occurrence here of sheet traps which ideally repeat the shape of the folds of the country rocks. The youngest deposits — metamorphosed by traps — are Lower Silurian limestones. This, apparently, permits the traps of the described territory to be tentatively correlated with the well-known traps of the Tungus basin.

On the basis of the relation of these intrusions to the principal ruptured zones of the Urinsk uplift an attempt may be made to draw certain conclusions concerning the time of its origin.

The Kudukta fault is one of the basic ruptured zones separating the Urinsk uplift from the linear folds of the Patomsk zone. On the north this fault passes into the large Dzherbinsk-Patomsk flexure. Both the Kudukta fault and the above flexure are the most important dislocations bounding the Urinsk uplift. This is the reason that the problem of structural relationships between the Kudukta fault and the traps is of special interest. Two conclusions could be made: 1) the basic magma does not follow the fault fractures as it is intruded; 2) the Kudukta fault truncates the traps in shifting two parts of the same intrusive sill by 1000 m in a vertical direction.

From the above it is evident that the Kudukta fault, and the west, were finally formed after the intrusion of the traps. Apparently, this position, — with but very slight reservation, can be applied to all other flexures and fractures in the Urinsk uplift, since we have no

reason to consider them as being of different age than the Kudukta fault. At the same time, so far we know of no fracture, of which there is an abundance within the limits of the Urinsk uplift, which could be utilized by trap bodies. Hence, the uplift in question must have been fully formed during the period when synclines existed in the Tunguss basin.

Briefly, the history of the geologic development of the northern fringe areas of the Patomsk upland can be summarized follows: in the Upper Proterozoic (Riphean, Sinian) and Lower Paleozoic, there existed here a number of downwarps (the Patomsk miogeosyncline, Nyuya trough) characterized by distinctly expressed inherited development. We are inclined to share Ye. V. Pavlovskiy's opinion that the depressions referred to above should be identified as a single major structure — a zone of pericratonnal subsidences.

The development of the troughs in question came to an end toward the end of the Lower Silurian when a wide zone of linear folds was developed around the Patomsk upland from the north in the form of a smooth arc. The Urinsk uplift was formed in the Upper Paleozoic and its structure was superimposed on the linear folds of the Patomsk zone.

REFERENCES

1. Arkhangel'skaya, N. A., V. V. Grigor'yev, and K. K. Zelenov, *Fatsii nizhn-ekembriyskikh otlozheniy yuzhnoy i zapadnoy okrain Sibirskoy platformy*. [THE FACIES OF THE LOWER CAMBRIAN DEPOSITS ON THE SOUTHERN AND WESTERN EDGES OF THE SIBERIAN PLATFORM]: Tr. Geol. in-ta Akad. Nauk SSSR, vyp. 33, 1960.
2. Gugol', R. F., *Normal'nyy razrez kembriyskikh otlozheniy Del'geyskoy razvedochnoy ploschadi po dannym glubokogo bureniya*. V kn. "Voprosy geologii i neftegazonosnosti Yakutsk. ASSR". [NORMAL SECTION OF CAMBRIAN DEPOSITS IN THE DELGEYSKAYA PROSPECTING AREA FROM DEEP BORING DATA. From the book, "PROBLEMS OF GEOLOGY AND OIL AND GAS POTENTIALITY OF YAKUTIA"]]: Gostoptekhizdat, 1958.
3. Gurari, F. G., *K stratigrafii kembriya yugo-vostoka Sibirskoy platformy*. [THE GEOLOGY OF BAYKAL-PATOMSK UPLAND]: Izv. Akad. Nauk SSSR, ser. geol., No. 4, 1945.
4. Dombrovskiy, V. V., *Geologiya Baikalo-Patomskogo nagor'ya*. [THE GEOLOGY OF BAYKAL-PATOMSK UPLAND]:

- Tr. Vost.-Sib. geol. upr., vyp. 26, 1940.
5. Zhuravleva, Z. A., V. A. Komar, and N. M. Chumakov, Stratigrafi-cheskiye sootnosheniya patomskogo kompleksa s osadochnymi otlozheniyami zapadnogo i severnogo sklonov Aldan-skogo shchita. [STRATIGRAPHIC CORRELATION OF THE PATOMSK COMPLEX WITH THE SEDIMENTARY DEPOSITS OF THE WESTERN AND EASTERN SLOPES OF THE ALDAN SHIELD]: Dokl. Akad. Nauk SSSR, t. 128, No. 5, 1959.
 6. Zaytsev, N. S., and N. V. Pokrovskaya, Stratigrafiya i tektonika nizhnepaleozoyskikh otlozheniy rayona srednego techeniya r. Leny. V kn. "Ocherki po geologii Sibiri". [STRATIGRAPHY AND TECTONICS OF LOWER-PALEOZOIC DEPOSITS IN THE MIDDLE COURSE REGION OF THE LENA RIVER. From the book; "OUTLINES OF THE GEOLOGY OF SIBERIA"]: Vyp. 15, Izd-vo Akad. Nauk SSSR, 1950.
 7. Zaytsev, N. S., O tektonike yuzhnoy chasti Sibirskoy platformy. V kn. "Voprosy geologii Azii". [THE TECTONICS OF THE SOUTHERN PART OF THE SIBERIAN PLATFORM. From the book; "Problems of the Geology of Asia"]: T. 1. Izd-vo Akad. Nauk SSSR, 1954.
 8. Zegebart, D. K., and Z. M. Starostina, K voprosy geologicheskoy istorii severovostochnoy chasti Leno-Yeniseyskogo polya i severnoy chasti Baykal'skoy skladchatoy zony. [THE PROBLEM OF THE GEOLOGIC HISTORY OF THE NORTHEASTERN PART OF THE LENA-YENISEY AREA AND THE NORTHERN PART OF THE BAYKAL FOLDED ZONE]: Probl. sov. geologii, No. 4, 1935.
 9. Kolyun, M. N., Stratigrafo-litodogicheskii ocherk verkholenskoy svity kembriya spednego i verkhnego techeniya r. Leny Materialy. [STRATIGRAPHIC AND LITHOLOGIC OUTLINE OF THE UPPER-LENA CAMBRIAN SERIES AND THE UPPER REACHES OF THE LENA RIVER]: Vses. n.-i. geol. in-ta, nov. ser., vyp. 24, 1959.
 10. Naslov, V. P., Sushchestvuyet li pereryv mezhdru srednim kembriyem i verkholenskoy svitoy v Vostochnoy Sibiri? [IS THERE A BREAK BETWEEN THE MIDDLE CAMBRIAN AND THE UPPER-LENA SERIES IN EASTERN SIBERIA?]: Izv. Akad. Nauk SSSR, ser. geol., No. 3, 1952.
 11. Offman, P. Ye., Tektonika i vulkanicheskiye trubki tsentral'noy chasti Sibirskoy platformy. V kn. "Tektonika SSSR". [THE TECTONICS AND THE VOLCANIC PIPES OF THE CENTRAL PART OF THE SIBERIAN PLATFORM. From the book; "Tectonics of the USSR"]: T. 4. Izd-vo Akad. Nauk SSSR, 1959.
 12. Pavlovskiy, Ye. V., Geologicheskaya istoriya i geologicheskaya struktura Baykal'skoy gornoy oblasti. [THE GEOLOGIC HISTORY AND GEOLOGIC STRUCTURE OF THE BAYKAL UPLAND]: Tr. In-ta geol. nauk Akad. Nauk SSSR, vyp. 99, ser. geol., No. 31, 1948.
 13. Pavlovskiy, Ye. V., Zony perikratonnykh. Opuskanii-platformennyye struktury pervogo poryadka. [ZONES OF PERICRATONAL SUBSIDENCES - FIRST ORDER PLATFORM STRUCTURES]: Izv. Akad. Nauk, SSSR, ser. geol., No. 12, 1959.
 14. Salop, L. I., Dokembriy Baykal'skoy gornoy oblasti. V kn. "Tez. dokl. na Mezhdruvedomstv. Soveshch. po razrabotke unifits. stratigraf. skhem Sibiri". 1956 g. [THE PRECAMBRIAN OF THE BAYKAL UPLAND. From the book; "Outlines of the Report at the Inter-Departmental Conference on the Development of a Unified Stratigraphic Map of Siberia"]: Izd-vo Akad. Nauk, 1958.
 15. Tetyayev, M. M., Geotektonika SSSR. GONTI [THE GEOTECTONICS OF THE USSR]: 1938.
 16. Frolova, N. V., Geologicheskii ocherk sredney chasti Leno-Kirengskogo mezhdru-Rech'ya. V kn. "Ocherki po geologii Sibiri". [GEOLOGICAL OUTLINE OF THE CENTRAL PART OF THE LENA-KIRENGA INTERFLUVE. From the book; "Outlines of the Geology of Siberia"]: Vyp. 18. Izd-vo Akad. Nauk SSSR, 1955.
 17. Chumakov, N. M., Novyye dannyye o geologicheskoy stroenii yugo-zapadnogo obramleniya Vilyuyskoy vpadiny. [NEW DATA ON THE GEOLOGIC STRUCTURE OF THE SOUTHWESTERN FRINGES OF THE VILYUY DEPRESSION]: Izv. vysshe. uch. zavedniy. Neft' i gaz, No. 3, 1958.
 18. Chumakov, N. M., Stratigrafiya i Tektonika yugo-zapadnoy chasti Vilyuyskoy vpadiny. V kn. "Tektonika SSSR", t. 4. [THE STRATIGRAPHY AND TECTONICS OF THE SOUTHWESTERN PART OF THE VILYUY DEPRESSION. From the book; "The Tectonics of the USSR"]: Izd-vo Akad. Nauk SSSR, 1959.

19. Shalek, Ye.A., Uglovoye nesoglasiye mezhdu verkhnim nizhnim kembriyem na zapadnom sklone Severo-Baykal'skogo nagor'ya. [THE DISCONFORMITY BETWEEN THE UPPER AND LOWER CAMBRIAN ON THE WESTERN SLOPE OF THE NORTH BAYKAL UPLAND]: Byul. nauchno-tekhn. inform. M-vo geol. i okhrany nedr SSSR, No. 4, 1958.
20. Shatskiy, N.S., O strukturnykh svyaz-yakh platform so skladchatymi geosinklinal'nymi oblastyami. Sravnitel'naya tektonika drevnikh platform. [THE

STRUCTURAL RELATIONSHIPS BETWEEN PLATFORMS AND FOLDED GEOSYNCLINAL AREAS. COMPARATIVE TECTONICS OF ANCIENT PLATFORMS]: Izv. Akad. Nauk SSSR, ser. geol., No. 5, 1947.

Geological Institute of the
U.S.S.R. Academy of Sciences,
Moscow

Received June 17, 1960

AGE AND STRATIFICATION SEQUENCE OF DEPOSITS IN THE UPPER PART OF THE KAR-TAU SERIES IN THE SOUTHERN URALS¹

by
Yu. R. Bekker

The article describes the stratification sequence of the Min'yar and Ukian deposits of the Karatau series of the Bashkir geanticline. The data given permit these deposits, and consequently the entire Karatau series, to be considered as a part of the Proterozoic formations.

The upper deposits of the Karatau series in the Southern Urals are still inadequately studied. Meanwhile, a detailed investigation of these formations provides a wealth of material with which to consider a number of cardinal problems relative to the geology of the Urals.

1. The Min'yar Series

In the section of the ancient Ural formations the Min'yar series lies between the overlying Ukian and the underlying Inzer deposits. In the Southern Urals this complex was studied by O. P. Goryainova, E. A. Fal'kova [5, 6], H. Dingel'shtedt [8], K. A. L'vov [11], A. I. Li [15], A. P. Tyazhev and S. M. Domrachev [16].

However, "attempts to divide the Min'yar formations into a set of beds traceable over considerable areas have failed so far" [16]. This fact is attributable to the view which was generally held until recently that Min'yar deposits are lithologically uniform, pure carbonate sediments in which terrigenous rocks play no significant part. In the meantime, investigations of these rocks showed that key horizons, covering large areas, can be identified in them.

I distinguish two formations in the Min'yar deposits: the Minki and the B'yanki strata (Figure 1).

The deposits of the lower Minki stratum, though poorly exposed, are apparently widespread in the Southern Urals. Outcrops of these rocks can be observed along the Yurezan'

River near the railroad siding at Minki (from which the name of the stratum was derived), along the Sim River in the vicinity of the railroad siding at B'yanki, along the Revat River west of Urman-Revat settlement, along the Belaya River near the village of Muradymovo, and in a number of other places.

In the basin of Sim River, near the settlement of Min'yar, a well exposed section of the Minki stratum reveals two beds (from top to bottom): 1) a bed of clearly stratified limestones with thin platy structure and dolomites; and 2) a bed of algal dolomites.

In the lower section of the Min'yar series the lower beds grade into the Inzer deposits without exhibiting signs of erosion. This type of transition is traceable near the Kal'tyarau River (Belaya River basin) as described in the following section.

The Inzer Series: 1. A bed of alternating greenish-grey, less frequently bordeaux-colored (red) sandy shales containing subordinate intercalations of grey siltstones. The proportion of clay shales increases in the upper section of the series. Noteworthy is the considerable amount of chlorite and almost total lack of glauconite — a mineral which is very typical of these rocks in the western sections of the Inzer series. The shales are plicated and more than 50 meters thick.

The Min'yar Series: 2. Pinkish, coarsely-crystalline limestones containing greenish clay flucans and quartzite veinlets are superposed sharply upon the Inzer shales described above. The azimuth of dip is $90^\circ < 34^\circ$. The thickness of the limestones is 20 cm.

3. A bed of interstratified white, greenish-grey thin-platy-structure limestones containing subordinate intercalations of greenish-grey

¹Vozrast i posledovatel'nost' naplastovaniya i slozheniy verkhney chasti Karatauskoy Serii vzhnogo Urala. pp. 49 - 60.




FIGURE 1. Stratigraphic column of the upper section of the Karatau series in the Southern Urals.

1 - dolomites, limestones with sandstone lenses; 2 - algal dolomites; 3 - detrital limestones, dolomites, siltstones, argillites; 4 - algal dolomites, limestones with medium platy structure; 5 - layered dolomites, flinty limestones; 6 - coarse platy limestones, dolomites; 7 - siltstones, argillites; 8 - algal dolomites; 9 - thinly-layered dolomites and limestones; 10 - glauconite sandstones, argillites.

marls and argillites. The clay content diminishes upward along the section. The thickness of this bed is 3.4 m.

4. Medium platy-structure limestones, light-grey at the bottom and grey, above. Veinlets of quartzite, sutures and styolitic structures are often observed in the limestones, which exceed 10 m in thickness.

Similar contacts between the Inzer and Min'yar series is also typical of other sections (Min'yar settlement), where minor ferrugination may sometimes be observed in the near-contact zone.

Peculiar to the lower layer deposits of the Minki stratum is the presence of sutured and styolitic surfaces filled with a ferruginized

pelitic substance. Occasionally thin veinlets of calcite are observed in these rocks. In sections having particularly densely distributed calcite veinlets the rock acquires a pseudo-conglomerate aspect.

The mineralogical composition of the Min'yar series carbonate rocks was studied by the method proposed by L. G. Berg, which was given the name of thermal-volumetric-quantitative phase analysis. The principles involved in this method are discussed in detail in L. G. Berg's works and in an article written jointly by T. Ye. Biryukova, I. V. Yevseyeva, I. I. Ivanova, Ye. P. Levando, and O. I. Nekrasova [2].

The results of the thermal-volumetric-quantitative phase analysis,² in conjunction with the data obtained from microscopic examination involving color reactions, make it possible to identify two varieties of carbonate rocks in the Minki stratum: dolomitic limestones and dolomites.

The dolomitic limestones are grey and dark-grey in color and contain dolomite (up to 25%) and an insignificant amount of insoluble residue (2.4%). Thermal analysis revealed the presence of siderite and rhodochrosite occurring as admixtures usually not exceeding 1%. The presence of (0.01-0.03%) strontium in these deposits was detected by spectral analyses.

The Minki stratum dolomites are brown-grey in color and manifest an uneven tubercle-like bedding plane. The dolomites are more macrocrystalline than the limestones. Veinlets of calcite are observed in them. This, probably, explains the rather high content of calcite established by quantitative thermal-volumetric analysis in these rocks. The proportion of magnesite in the Minki dolomites ordinarily does not exceed 7%.

Carbonate rocks in the top layer contain cylindrically shaped stromatoliths having a maximum diameter of from 2 to 3 cm. They are almost at right angles to the bedding plane of the rock and are "enveloped" by a clayey substance containing a high percentage of glauconite. The algal layer rocks, which seem to be less abundant than the underlying deposits, terminate the Minki sequence of the Min'yar series. The apparent thickness of the Minki stratum in many sections does not exceed 100 m, with the top algal layer usually accounting for some 5-8 meters.

The B'yanki stratum is the thickest and most exposed member of the Min'yar series, covering a considerable area in the Southern Urals from the basin of the Ay River in the north to the basin of the Belaya River in the south. In the stratigraphic section of this stratum near the B'yanki siding on the Sim River the following beds (from bottom up) can be identified

² This analysis of the carbonate rocks of the Min'yar series was made by T. Ye. Biryukova.

1) argillites and siltstones; 2) massive dolomites; 3) dolomites containing chert; 4) limestones with medium platy structure and dolomites.

The lower layer of the B'yanki formation is exposed only in the basin of the Sim River. Here, in the vicinity of the B'yanki railroad siding, overlying the Minki limestones are greenish-grey argillites interbedded with light-grey, thin-plated quartzose siltstones containing glauconite. The argillites predominate in the lower part of the section, the siltstones, in the top.

The greenish-grey argillites of this formation contain small quantities of quartz terrigenous material and large round grains of authigenic glauconite. According to the data of thermal, electron-diffraction, and electron-microscopic investigations, argillite fraction particles measuring less than 0.001 mm occur mainly in the hydromicaceous component. The heating curve of the argillites is characterized by three endothermic reactions at 130°, 630°, and 940° temperatures and one exothermic reaction at 1000°. Electron-diffraction studies produced the following unit-cell parameters for the argillites in question: $a = 519$, $b = 8.97$, $c = 20.0$ kX, $\beta = 95^\circ$. A two-layer period and a rather low degree of structural perfection characterize the hydromicas under consideration. Electron-diffraction analyses have revealed the presence of elongated, sharply delineated scales of glauconite and fragments of kaolinite contaminating these argillites.

The siltstones are light grey in color, are of the thin-layered quartz variety and contain large, whitish grains of glauconite. Apart from the prevalent quartz grains, particles of plagioclases and glauconite. Apart from the prevalent quartz grains, particles of plagioclases and quartzites were found in the clastic material of these rocks. The clastic material is fairly well sorted out much less rounded. The accessories are zircon, chromite, olivine, and magnetite. The cement is mixed: carbonate, quartzose, and occasionally sericitic.

The observed thickness of the terrigenous layer in this section is 35 m.

Rocks of this formation were found in the estuarine zone of the Min'yar River in the same position of the stratigraphic column as the Min'yar series. Here, in the lower sections of this stratum was found an intercalation of ferruginized oölitic limestones containing significant amounts of terrigenous quartz material. The oölitic are composed of carbonaceous ferruginized material and are 5 mm in diameter.

The presence of oölitic in the carbonate

rocks of the Min'yar series is a matter of decided interest. According to L. Cayeux's [25] and D.V. Nalivkin's [14] data oölitic have not been known to occur in Precambrian carbonate rocks. However, in the Karatau series on the Western slope of the Southern Urals, oölitic limestones are fairly conspicuous. Apart from the mentioned stratum, oölitic limestones also are found in the deposits of the Subinzer series overlying the Ukian formations (Figure 2).




FIGURE 2. Oölitic from the Ukian deposits in the Inzer basin.

(Nicol prisms are parallel; 32x).

It should be noted that intercalations of "roe-type" (oölitic) limestones occur in the Subinzer series, in which the oölitic are associated with spherulites.

Thus, conditions favorable to the formation of oölitic repeatedly occurred, not only in the Paleozoic, but also in the Proterozoic.

In contrast to the discussed, usually poorly exposed terrigenous deposits, the massive dolomites overlying the layer of the B'yanki formation produce spectacular cliffs. These dolomites usually occur in the sections of the Min'yar series in the Ay, Yurezan', Sim, Inzer, Zilim, and Belaya rivers basins. The massive dolomites are brownish-grey in color and mottled with reddish spots. Dolomites with conchoidal structure and limestones with thin, platy structure occur in them as intercalations. Stromatolites of the *Collenia* genus in certain places constitute the rock-forming organisms in these deposits. Individual sections show thin bands of dolomitic breccia subordinate to the massive dolomites.

According to the data of quantitative thermal-volumetric phase analysis these rocks are mostly composed of dolomite (84.7%), with calcite, siderite, and smithsonite occurring as admixtures. The dolomite is macrocrystalline, individual crystals being as long as 2 mm. The proportion of insoluble residue in the dolomites is about 4%. The thickness of the massive dolomite bed does not exceed 200-250 m in many sections.

Upward in the section, the massive dolomites gradually change into a bed of dolomites containing flint. These dolomites make a well-exposed key horizon traceable in the majority of the Min'yar series sections in the Southern Urals. Bedrock outcrops of this horizon can be observed along the Ay River, in the area of Vanyashkino and Sikiyaz-Tamak villages; along the Yurezan' River, near Vyazovaya station and the settlement of Ust'-Katav; along the Sim River, near the Min'yar and Sim factories; along the Ray and Meneyka Rivers (Inzer River basin), and in several other localities.

The carbonate rocks containing the flinty concretions are represented by grey, light-grey dolomites and very rarely, by limestones.

The dolomites are microcrystalline, locally recrystallized, contain numerous calcite veins, and a small proportion of insoluble residue (3.4%). A small quantity of siderite was revealed in these rocks by quantitative-thermal-volumetric phase analysis.

A study of the Min'yar series flints permitted us to identify the nodule and vein forms of silica segregation. The former is particularly characteristic of the rocks of the discussed horizon. Among the concretionary siliceous formations in this horizon, one can distinguish layer-type, lenticular concretions, and flint nodules.

The thickness of the siliceous concretion layer is very irregular. In places it attains 20-30 cm, but 30-40 m further along the strike it drops to 2-3 cm and even to zero - in flint lenses. Occasionally one layer splits into two or three separate thinner layers. At individual outcrops, however, it was possible to trace the flint layers extending without substantial variation in thickness for 80 - 100 m along the strike.

The flint concretions are, probably, the most commonly occurring form of silica deposits in the Min'yar series. Their shapes are very diversified. Most cross-sections are ellipsoidal and spherical and, on the bedding planes, look like intricately uncoiled, knotty rope. Siliceous concretions of cylindrical form (about 10 cm in diameter) also are found in a vertical position.

The concretions are usually grey, sometimes dark-grey and unevenly colored, and consist of cryptocrystalline silica. Individual sections reveal the preferential distribution of flints, lighter varieties gravitating toward the lower, the darker - towards the upper, parts of the horizon under consideration. In places, the relationship between the color of the flints and the enclosing rocks is apparent. For example, light-color flints frequently predominate in light-grey dolomites. However, both light and dark flints are found in the dark-grey carbonate rocks. Beds of siliceous substance containing a significant amount of carbonate material are found in places where the concretions are in contact with the enclosing rocks. Often the groundmass of the cryptocrystalline flint contains individual idiomorphic rhombohedra of dolomite. Widely distributed throughout the concretions are veinlets containing quartz in the center and chalcedony on the periphery.

Carbonate rocks containing flint concretions often display well pronounced layers. This makes it possible to study the relationship between stratification and the shapes of the siliceous concretions. We made such a detailed study of cliff outcrops in the basins of the Yurezan' and Sim Rivers. These investigations disclosed that in the majority of cases (55-65%) the concretions clearly followed the contours of the layers. Only a small part of the concretions (10-15%) cut across the interbedded layers. For concretions embedded in thick unstratified dolomites, such a relationship could not be established. There were instances where the flint concretions were on the uneven bedding plane of the dolomites. As may be seen in Figure 3, the flint rocks faithfully repeat the irregularities of the bedding and exhibit the same wavy upper surface.

The data on the relationship between the shape of the flint concretions and stratification attest to the fact that most flint concretions in this case was formed concurrently with sedimentation.

During the lithification process there was a subsequent shift of the silica and a few of the veinlets and siliceous concretions which now cut across the layers, were formed.

It should be noted that syngenetic and diagenetic concretions can not be distinguished either macroscopically or microscopically. Their identical composition once again indicates that their formation is associated with a redistribution of an identical portion of silica.

Small stringers of flint are observed in the silica bed itself and in a considerably smaller amount in the overlying and underlying layers. Consequently, they have no important stratigraphic significance.




FIGURE 3. Relationship of the shape of flint bodies to the stratification of the carbonate rocks of B'yanki series.

1 - stratified limestones; 2 - flint bodies.

Because there are no organic remains having silica skeletons in the Min'yar deposits, it must be assumed that the silica here is of inorganic origin and must have been transported into the marine basin of this period from the adjacent uplands.

In concluding the review of the lithological characteristics of this horizon, it is important to emphasize the fact that the silica concretions in Min'yar series occupy a definite stratigraphic position. This is indicated both by direct stratigraphic study and by the data on the lithology and the conditions under which these concretions were formed. The dolomites containing layered and lenticular concretionary segregations of silica in the Min'yar series constitute a reliable key horizon permitting the identification of sub-siliceous, siliceous, and supra-siliceous layers even when the exposure is poor.

In the upper part of the Min'yar series in the Yurezan' and Sim river basins dark-grey argillal and medium-platy and algal dolomites and limestones were found. Here C.N. Domrachev discovered some problematical remains of *Fistulipora*. The thickness of these deposits does not exceed 50 m and varies considerably from section to section because of the ungressive deposition of the overlying Ukian series.

II. The Ukian Series

The terrigenous-carbonate rocks of the Ukian formation are unconformably overlain by the upper bed of the Min'yar series, discussed earlier. First, these deposits were differentiated and described layer by layer by S.M. Domrachev near the Min'yar works and on the Uka River (Sim basin). The Ukian series is located in these sections between the usually characterized Devonian deposits and

the Min'yar rocks. The Ashinian deposits, which ordinarily occur in this interval of the Paleozoic sequence are absent here. In view of this fact, S.M. Domrachev assumed that the series he identified was a facies analog of the Ashinian deposits. This point of view was later shared by A.I. Olii [17] and a few other geologists of the Southern Urals Geologic Administration. The same views were also reflected in a collated study by S.G. Sarkisyan and G.I. Teodorovich [19].

My geologic survey in the Sim river basin and adjacent regions permitted me to establish the extensive regional distribution of the Ukian deposits in many of the sections of the ancient formations in the Southern Urals, for example, in the oil pipeline trench on the western limb of Suleymanovskaya anticline, at the watershed of the Yurezan' and Minka Rivers, along the Yurezan' River near the settlement of Ust'-Katav and Limonovka village, along the Sim River in the vicinity of the Sim works; further south along the Rau River (Inzer basin), Basa, Meneyka, Revat - west of the settlement of Urman-Revat, along the Belaya River near Krivaya Luka.

Two beds, interconnected by gradual transgressions, can be distinguished in the majority of the Uka deposit sections.

The lower bed consists of a rather diversified series of rocks including grey and light-grey siltstones, green argillites, clastic and oolitic limestones, ferruginized dolomites, and ash-grey marls. The Ukian sequence usually begins either with conglomerates containing pebbles of the subjacent carbonate rocks, or with ferruginized siltstones.

The siltstone clastic material is represented by well-rounded but poorly sorted quartz grains (80-90%) containing an insignificant admixture

Table 1

Number in sequence	Name of minerals	Content of minerals, in %				Number in sequence	Name of minerals	Content of minerals, in %			
		Min'yar station	Sim station	Ust'-Katav	Average			Min'yar station	Sim station	Ust'-Katav	Average
1	Pyrite	—	—	2,1	0,7	13	Monoclinic pyroxene group	—	—	—	—
2	Martite	—	0,3	—	0,1						
3	Hematite	90,4	92,5	—	61,0			1,3	1,8	0,4	1,2
4	Limonite	RT ¹	RT	54,1	18,0	14	Olivine	1,0	1,2	4,3	2,2
5	Ilmenite	RT	0,3	2,1	0,8	15	Garnet	1,7	0,9	3,0	1,9
6	Rutile	0,3	—	0,4	0,2	16	Epidote	RT	—	—	RT
7	Anatase	RT	—	—	RT	17	Tourmaline	1,0	0,3	RT	0,4
8	Leucoxene	—	—	1,3	0,4	18	Monazite	RT	—	RT	RT
9	Chromite	RT	0,9	8,5	3,1	19	Glaucanite	RT	RT	RT	RT
10	Chromopicitote	RT	0,3	0,8	0,4	20	Mica	0,3	—	0,4	0,2
11	Magnetite	RT	RT	RT	RT		Output of concentrate in grams per cu. m. of rock ²				
12	Zircon	4,0	1,5	22,6	9,4			27,4	26,6	8,0	—

¹ RT - rare² Synthetic concentrates were obtained by crusing lump samples weighing about 10 kg and subsequent washing out of heavy fraction minerals.

NOTE: Comma represents decimal point.

of plagioclase particles. Glaucanite usually is the principal component of Ukian siltstones. The rock cement is basal. Its composition is mixed: sericitic and carbonaceous.

The following heavy fraction minerals were in the siltstones: pyrite, martite, hematite, limonite, ilmenite, rutile, anatase, leucoxene, chromite, chromopicitote, magnetite, zircon, monoclinic pyroxene, olivine, garnet, epidote, tourmaline, monazite, glaucanite, mica.³ The percentage values of these minerals are shown in Table 1.

Noteworthy is the increasing content of accessory minerals in the basic and ultrabasic rocks (olivine, chromite, chromopicitote) eastward from the Min'yar works (1%) and Sima settlement (2.4%) toward the Ust'-Katavskiy works (13.6%). Such percentage variations in these components, apparently, indicate that the clastic material in the Ukian deposits was transported from the island uplands in the east which consisted of basic and ultrabasic rocks.

The detrital sediments in the clastic limestones contain pebbles of microcrystalline dolomites, pelitomorphic micro- and macro-crystalline limestones, and silt-size quartz grains. Carbonate rock pebbles in the clastic limestones

attain a size of 5-8 cm. The amount of insoluble residue in these rocks, according to data from a single determination, is 5.7%. Calcite, with occasional strong ferruginization, is the cement in the clastic limestones.

The deposits of the lower formation gradually merge into the basically carbonate overlying rocks which complete the Ukian series section. Good outcrops of these rocks can be observed along the Basa River near the settlement of Kulmas, as well as in the region of the Ust'-Katav and Min'yar works. Distinguishable among the carbonate rocks in the upper section of the Ukian series are light-grey dolomites and grey limestones. Apart from dolomite (79.4%) and secondary calcite (15.9%), the presence of magnesite (2.2%) and siderite (0.4%) was revealed in the light-grey varieties of these carbonate rocks by quantitative-thermal-volume phase analysis.

The grey limestones may be more correctly referred to as dolomitic limestones since, according to thermal analyses, they were found to contain a significant amount of dolomite (26.2%).

Lenses of light-grey friable feldspathic sandstones occur in the upper formation of the Ukian series as seen in individual sections along the Sim and Yurezan' Rivers. The mode of deposition of these rocks is very unusual. The sandstones usually fill 1-1.5-m wide pockets in the carbonate rocks. This deposition of sandstones apparently, results from the moving of terrigenous material in connection with the processes

³Determinations made by the Mineralogical Laboratory of the All-Union Scientific-Research Geologic Institute's (VSEGEI) central expedition.

of ancient karst formation. This theory seems especially probable because occasionally undisturbed, primary, sheet-like or lenticular bedding is observed in the sandstones.

A comparative study of the Ukian, Min'yar, and Ashinian deposits shows that the composition and the organic remains in the Ukian formations display a number of features common to the Min'yar rocks and rather sharply different from the deposits of the Ashinian series. In view of this fact, the supposition concerning the coetaneousness of Ashinian and Ukian deposits must be based on a detailed stratigraphic analysis of the sections.

Investigations carried out in this direction indicate that the Ukian formations lie, with a small break, on the flint and alga-B'yanki carbonate rocks of the Min'yar series and in the basins of the Yurezan' and Inzer Rivers are covered by Ashinian series, and in the basin of the Sim River — by Devonian deposits. Of particular interest are the sequence relationships between the Ashinian and Ukian deposits as they can be observed in the outcrops near the settlement of Ust'-Katav on the Yurezan' River. A generalized description of this section is given below (up from the bottom).

Min'yar series. 1. Massive grey dolomites belonging to the upper part of the Min'yar series containing globular algal formations 5-7 cm in diameter. Total thickness of the bed — about 10 m.

2. Dark-grey and grey limestones, with medium platy structure, and dolomites, forming a 15-meter layer.

Ukian series. 3. Clastic limestone containing gravel and the flattened pebbles of the underlying carbonate rocks. Total thickness, 16 cm.

4. Dark-grey ferruginized detrital or organic limestones, mottled, reddish, locally dark-grey dolomites alternating with grey siltstones and argillites. A layer-by-layer description of this 120-130 m section was prepared on the basis of small bedrock outcrops and mine workings.

5. Massive algal limestones and dolomites, the upper part containing lenses of feldspathic-quartz sandstones which, on cursory inspection, are hardly distinguishable from the enclosing dolomites. Thickness, 85 m.

Ashinian series. 6. Conglomerate with dark-grey flint pebbles (70 to 75%), dolomite (10 to 15%), quartzite and quartz (15 to 20%). The clastic material in the conglomerate is fairly well rounded. The average roundness factor, determined by Wadell's method, is 0.48 for the flint pebbles. The layer is 10 cm thick.

7. Green-grey argillites, containing intercalated siltstones. The observed thickness is about 15 m. Higher, after a thin meadow are interval, on the slope there are exposures of Uryukian gravel conglomerates belonging to the Ashinian series which contain quartz, feldspar, and flint pebbles.

Thus, in the section near Ust'-Katav the Ashinian deposits overlap transgressively the Ukian series. A similar stratigraphic position is occupied by the Ukian series in other sections of the Bashkir anticlinorium (basin of the Inzer and Bas Rivers). In these sections obolitic limestones are occasionally observed in the Ukian series. In these limestones L.V. Khmelevskaya [21] found the remains of annelides in the Sim River basin.

In some places the Ashinian formations also lie unconformably on the Min'yar deposits. For example, in the section near Sikiyaz-Tamak village on the Ay River, Ashinian sandstones overlay the flint dolomites of the B'yanki formations of the Min'yar series.

The above data on the occurrence, composition, and stratigraphic position of Ukian rocks permit us to reject as unsubstantiated the assumption that the Ashinian and Ukian deposits are coetaneous. In retaining the previous designation, the author of this article interprets the stratigraphic position of Ukian deposits in a different way, considering them as an independent formation in the Karatau series of the Southern Urals. The deposits of Ukian series begin a new cycle of sedimentation.

The Min'yar and Ukian deposits were formed in a fairly wide shallow-water marine basin which formerly covered a considerable area of the western slope of the Urals. These deposits were identified both on the western and eastern slopes of the Bashkir geanticline, with the eastern sections of the Min'yar deposits of the western fringe of the Bashkir anticlinorium characterized by great thickness and absence of noticeable facies alterations. This fact justifies the assumption that the marine Min'yar deposits covered the area of the present-day Bashkir anticlinorium and were later eroded (Figure 4).

East of the Min'yar basin, in the recent Ural-Tau zone, there were a number of base-leveled, possibly insular, uplifts. The erosion of these structures was responsible for the formation of a considerable proportion of the clastic material contained in the Min'yar and Ukian deposits.

III. Age of the Karatau Series

The age of the Karatau series, in particular of the Min'yar and Ukian deposits, is still one of the most debatable and essential problems of




FIGURE 4. Schematic Lithological Map of the Min'yar an

FIGURE 4. Schematic Lithological-Facies Map of the Min'yar and Ukan deposits in Bashkir geanticline.

(Scale, 1:2,500,000).

1 - distribution area of shallow-water marine deposits; 2 - area of presumed distribution of marine deposits; 3 - Base-leveled peneplain; 4 - the numerator indicates the section number, the denominator - the thickness of deposits.

Ural geology. Different authors assign the Min'yar series to the Proterozoic [3, 4, 23, 24], to the Sinian [20], and to the Cambrian [5, 6, 13]. The Ukan series was considered by S.M. Domrachev [7] as contemporaneous with the Lower Devonian and Eifelian deposits [22]. N.G. Chochia [22] correlated these formation with the Silurian, and the author of this article tentatively assigned them to the Lower Cambrian.

With the view to finding a more definite solution of these problems I have paid particular attention in my field operations to the layer-by-layer collection of fauna, gathering samples for spore analysis, and to the determination of the absolute age of the rocks by the argon method in terms of glauconites. Despite thorough search I failed to collect any identifiable faunal remains in the Min'yar and Ukan deposits.

A fairly large collection of assembled stromatolites was sent to A.G. Vologdin, but has not yet been monographically classified. According to A.G. Vologdin's preliminary report, the collection has revealed well-preserved products of Cyanophycean life activity, similar to the analogous formations found in the Cambrian and

Proterozoic of the Siberian platform, but still unrecorded in the literature.

Spores were identified both in the Min'yar and in the Ukan deposits.

In the argillites of the terrigenous horizon of the B'yanki stratum (in the vicinity of the settlement of Min'yar), I.N. Solomina was able to identify the following spores: *Mycteroligotritiles marmoratus* Tim., *Bothroligotritiles exasperatus* Tim., *Trachyoligotritiles incrassatus* (Naum.) Tim., *T. nevelensis* Tim., *T. planus* Tim. The complex of these spores differs from the Inzer series spores by the lack of the characteristic *Tiloligotritiles asper* Tim forms.

In the Ukan argillites, I.N. Solomina discovered a more diversified complex of spores: *Leiologotritiles mimitissimus* (Naum.) Tim., *L. crassus* (Naum.) Tim.; *Mycteroligotritiles marmoratus* Tim., *Trachyoligotritiles mimitus* (Naum.) Tim., *T. hyalinus* (Naum.) Tim., *T. incrassatus* (Naum.) Tim., *T. nevelensis* Tim., *T. arillatus* Tim.

According to I.N. Solomina's data, the spore complex of the Ukan stratum - judging by the

Table 2

No. in series	No. of sample	Stratigraphic horizon	Location	$\frac{Ar^{40}}{K^{40}}$	Age in millions of years
1	322	Min'yar series, lower part of Minki stratum	Sim River basin, Min'yar settlement	0,0520	760
2	158	Ukian series	Basa River basin, Kulmas settlement	0,0405	616
3	222	Top part of the Ukian series	Revat River basin, Urman-Revat settlement	0,0407	618

NOTE: Comma represents decimal point.

presence of *Trachyoligotriletes hyalinus* (Naum.) Tim., — is younger than the B'yankian stratum in the Min'yar series. B.V. Timofeyev and I.N. Solomina believe the spore complexes of the Min'yar and Ukian deposits to be characteristic of the Sinian formations.

Glaucanite — the most widespread syngenetic mineral of the Inzer, Min'yar, and Ukian deposits of the Karatau series — is usually found in the form of large oval grains of aggregate structure in the terrigenous, and often in the carbonate deposits. The size of its grains in terrigenous rocks is ordinarily several times that of the quartz fragments. These data, as well as the mode of glauconite occurrence in the section, cause us to reject the theory about possible widespread distribution of redeposited glauconite in these sediments.

The results of absolute age determination by the argon method on the basis of glauconites are incorporated in Table 2. The determination was made by N.I. Poleyeva and G.A. Kazakov [18].

The aforementioned data make it impossible to assign the Min'yar series, and consequently, also the subjacent deposits of the Karatau series, to the Cambrian and Ordovician.

The views concerning the Devonian and Silurian age of the Ukian deposits are totally groundless. The transgressive overlap of the Ashinian deposits on the rocks of the Ukian series, and data obtained by the argon method, attest to the much older origin of the Ukian formations.

The available factual material allows us, in concurrence with the views held by M.I. Garanya, N.S. Shatskiy and B.S. Sokolov, to consider the rocks of the Karatau series as belonging to the Late Precambrian.

The absolute age of the glauconite from the Inzer deposits, according to data resulting from several determinations, amounts to about

900 million years. Thus, it is possible to assume that the bottom beds of the Karatau series are not younger than 1 to 1.1 billion years, while the duration of Sinian sedimentation in the Urals should be estimated as not less than 400-500 million years. These, to a certain extent, unexpected results preclude the possibility of regarding the Sinian deposits as being formed within the limits of a single, independent period of the Paleozoic. In duration, the Sinian was five to six times longer than the longest Paleozoic period, the Cambrian, and can be compared only with the entire Paleozoic group.

These data and other arguments advanced in the studies of N.S. Shatskiy [23], B.M. Keller [9], Lee-Sy-Huang [10] and other authors, make it possible to consider the Sinian as corresponding to an independent group or even larger unit of the Precambrian.

REFERENCES

1. Bekker, Yu.P., O stratigraficheskom polozhenii ukskikh otlozheniy na Yuzhnom Urale. [THE STRATIGRAPHIC POSITION OF UKIAN DEPOSITS IN THE SOUTHERN URALS]: Dokl. Akad. Nauk SSSR, t. 122, No. 5, 1958.
2. Biryukova, T.Ye., I.V. Yevseyeva, I.I. Ivanova, Ye.P. Levando, and O.I. Nekrasova, K voprosu opredeleniya fazovogo sostava karbonatnykh Porod po metodu L.G. Berga (predvaritel'noye soobshcheniye). [THE PROBLEM OF PHASE COMPOSITION DETERMINATION FOR CARBONATE ROCKS BY L.G. BERG'S METHOD (PRELIMINARY REPORT)]: Tr. Vses. n.-i. geol. in-ta, nov. ser., vyp. 11 "Materialy po litologii," 1956.
3. Garan', M.I., Dokembriyskiye otlozheniya zapadnogo sklona Yuzhnogo Urala i svyazannyye s nimi poleznyye iskopayemye. [PRECAMBRIAN DEPOSITS ON THE

- WESTERN SLOPES OF THE SOUTHERN URALS AND THE ASSOCIATED MINERAL RESOURCES]: Tr. 17 ses. Mezhd. geol. kongr., 1939.
4. Garan', M.I., O vozraste v usloviyakh obrazovaniya drevnikh svit zapadnogo sklona Yuzhnogo Urala. [THE AGE AND DEPOSITION CONDITIONS OF THE ANCIENT SERIES ON THE WESTERN SLOPE OF THE SOUTHERN URALS]: Gosgeolizdat, 1946.
 5. Goryainova, O.P., and E.A. Fal'kova, Geologicheskiye issledovaniya v basseynе rek. B. Inzer i Zilim. [GEOLOGIC INVESTIGATIONS IN THE BOL. INZER AND ZILIM RIVER BASINS]: Tr. Vses. geol. razved. ob'yed., vyp. 272, 1933.
 6. Goryainova, O.P., and E.A. Fal'kova, Geologicheskiy ocherk Inzerskikh i Komorozovo-Zigazinskikh zhelezorydnykh mestorozhdeniy. [GEOLOGICAL OUTLINE ON THE INZER AND KOMOROZOVO-ZIGAZINSK IRON ORE DEPOSITS]: Tr. Bashkirsk. geol. upr., vyp. 9, 1940.
 7. Domrachev, S.M., Devon khrebta Kara-Tau i privileyushchikh rayonov Yuzhnogo Urala. [THE DEVONIAN OF THE KARATAU RANGE AND THE ADJACENT SOUTH URAL REGIONS]: Tr. Vses. n.-i. geol. - razv. in-ta, nov. ser., vyp. 61, 1952.
 8. Dingel'shtedt, N.N., K voprosy u geologicheskoy storoyenii Yuzhnogo Urala. [THE PROBLEM OF THE GEOLOGIC STRUCTURE OF THE SOUTHERN URALS]: Tr. Vses. geol. - razv. ob'yed. vyp. 243, 1932.
 9. Keller, B.M., Rifeyskiye otlozheniya krayevykh progibov Russkoy platformy. [RIPHEAN DEPOSITS IN THE FORE-DEEPS OF THE RUSSIAN PLATFORM]: Tr. In-ta geol. nauk Akad. Nauk SSSR, geol. ser., No. 37, 1952.
 10. Lee-Sy-Huang, Geologiya Kitaya. [THE GEOLOGY OF CHINA]: Izd-vo in. lit., 1952.
 11. L'vov, K.A., Dokembriyskiye i nizhnepaleozoyskiye otlozheniya Urala. [THE PRECAMBRIAN AND LOWER PALEOZOIC DEPOSITS IN THE URALS]: Ob'yasnitel'naya zapiska k geol. karte Urala m. 1:500,000, GGU, 1939.
 12. L'vov, K.A., O drevnikh otlozheniyakh Urala, ikh vozraste i stratigrafii. [THE ANCIENT DEPOSITS OF THE URALS, THEIR AGE AND STRATIGRAPHY]: Sov. geologiya, sb. 55, 1957.
 13. L'vov, K.A., and A.I. Olli, Ob otnoshenii srednego devona k ashinskoy svite i o vozraste nemykh tolshch zapadnogo sklona Urala. [THE RELATIONSHIP BETWEEN MIDDLE DEVONIAN AND THE ASHINIAN SERIES AND THE AGE OF THE BARREN FORMATIONS ON THE WESTERN SLOPES OF THE URALS]: Zap. Ros. mineralog. o-va, ch. 64, vyp. 2, 1935.
 14. Nalivkin, D.V., Ucheniye o fatsiyakh. Geograficheskiye usloviya obrazovaniya osadkov. [THE DOCTRINE OF FACIES. GEOGRAPHIC CONDITIONS OF SEDIMENTATION]: Izd-vo Akad. Nauk SSSR, 1956.
 15. Olli, A.I., Geologicheskiye issledovaniya v rayone srednego techeniya rek Lemeza i Inzer. [GEOLOGIC INVESTIGATIONS IN THE MIDDLE-COURSE REGIONS OF THE LEMEZA AND INZER RIVERS]: Tr. Bashkirsk. geol. upr., vyp. 1, 1936.
 16. Olli, A.I., Drevneyshiye otlozheniya zapadnogo sklona Urala. [THE OLDEST DEPOSITS ON THE WESTERN SLOPE OF THE URALS]: Izd-vo Saratovsk. un-ta, 1948.
 17. Olli, A.I., Eshche raz o vozraste ashinskoy svity na Urale i vzaimootnosheniiye s dokazannym paleozoyom. [AGAIN ON THE AGE OF THE ASHINIAN SERIES IN THE URALS AND ITS RELATION TO PROVEN PALEOZOIC FORMATIONS]: Sov. geologiya, sb. 45, 1955.
 18. Polevaya, N.I., G.A. Murina, and G.A. Kazakov, Opredeleniye absolyutnogo vozrasta osadochnykh porod po glaukonitam. [ABSOLUTE AGE DETERMINATION OF SEDIMENTARY ROCKS ON THE BASIS OF GLAUCONITES]: Sov. geologiya, No. 7, 1960.
 19. Sarkisyan, S.G., and G.I. Teodorovich, Osnoviyye cherty paleogeografii devonskoy epokhi Uralo-Volzhskey oblasti. [THE PRINCIPAL FEATURES OF THE DEVONIAN EPOCH'S PALEOGEOGRAPHY IN THE URAL-VOLGA REGION]: Izd-vo Akad. Nauk SSSR, 1955.
 20. Sokolov, B.S., O vozraste drevneyshego osadochnogo pokrova Russkoy platformy. [THE AGE OF THE OLDEST SEDIMENTARY MANTLE OF THE RUSSIAN PLATFORM]: Izv. Akad. Nauk SSSR, ser. geol., No. 5, 1952.
 21. Khmelevskaya, L.V., Fossilizirovannyye annelidy iz paleozoya khr. Kara-Tau doliny r. Sim Bashkirskoy ASSR. [FOSSILIZED ANNELIDS FROM THE PALEOZOIC OF THE KARATAU RANGE IN THE

SIM RIVER VALLEY OF THE BASHKIR ASSR]: Tr. Neft. n. -i. geol. -razv. in-ta, ser., vyp. 34, 1949.

22. Chochia, N.G., Geologicheskoye stroyeniye Kulvo-Visherskogo kraya. [GEOLOGIC STRUCTURE OF KOLVO-VYSHERSKIY TERRITORY]: Gostoptekhizdat, 1955.
23. Shatskiy, N.S., O granitse mezhdupaleozoyem i proterozoyem i o rifeyskikh otlozheniyakh Russkoy platformy. [THE CONTACT BETWEEN THE PALEOZOIC AND PROTEROZOIC AND ON THE RIPHEAN DEPOSITS OF THE RUSSIAN PLATFORM]: Izv. Akad. Nauk SSSR, ser. geol., No. 5, 1952.
24. Shatskiy, N.S., Ocherki tektoniki Volgo-

Ural'skoy neftenosnoy oblasti i smezhnykh chastey zapadnogo sklona Yuzhnogo Urala. [ESSAYS ON THE TECTONICS OF THE VOLGA-URAL OIL-BEARING REGION AND THE ADJACENT AREAS ON THE WESTERN SLOPES OF THE SOUTHERN URALS]: Izd-vo Mosk. o-va ispyt. prirody, 1945.

25. Cayeux, L., Les roches sédimentaires de France; roches carbonatées (calcaires et dolomies). Mém. Carte Géol. France, 1935.

All-Union Scientific-Research Geologic Institute (VSEGEI) of the Ministry of Geology and Conservation of Mineral Resources of U. S. S. R., Leningrad

Received July 27, 1959

OUTLINE OF THE STRATIGRAPHY AND TECTONICS OF THE TAS-KHAYAKHTAKH RANGE¹

by
N. A. Bogdanov

INTRODUCTION

The Tas-Khayakhtakh Range extends for more than 300 km on a line from the upper reaches of the Indigirka River to the valley of the Selennyakh River, covering a 50-80 km wide belt, and forming the main divide between the basis of the Yana and Indigirka Rivers. Paleozoic formations are well developed within its confines. A study of these deposits was initially made in 1933 by V. A. Fedortsev [14], and in 1935-1939 by A. P. Atlasov [1], K. K. Demokidov [6], V. I. Fagutov and V. K. Lezhoyev. As a result of these surveys, Upper Cambrian, Ordovician, Silurian, Devonian, and Lower Carboniferous deposits were identified, as well as those of the Permian-Triassic and Jurassic. In 1954-1958, geologists of the Yana Regional Geologic Administration, L. K. Dubovikov, Yu. T. Krezhevskikh, R. N. Titov, and others made geologic surveys of the Tas-Khayakhtakh Range. The results of this work were correlated by L. K. Dubovikov and V. K. Lezhoyev [7], as well as by A. A. Nikolayev [11], I. Ya. Nekrasov and V. I. D'yachenko [10], and A. V. Zimkin [8]. Nonetheless, a number of questions concerning Paleozoic stratigraphy remained unresolved and required additional study. Nor was a detailed description of the tectonic structure of the area available.

In recent years the author of this article has carried out geologic investigations in the southern and central parts of the Range, including the valleys of the Umba, Uchugey-Yuryakh, Biyergichek, Kuranakh-Yuryakh, Dogdo, on the upper reaches of the Solonya River, and other areas. These operations enabled him to prepare a more differentiated, stratigraphic chart of the region's Lower-Paleozoic formations, and to describe in greater detail the peculiarities of the tectonic structure of Tas-Khayakhtakh Range.

Our data on the stratigraphy of Upper Ordovician and Lower Silurian formations were recently published, in part [2].

In this paper we shall deal with the stratigraphy of the entire Paleozoic complex.

During field survey, a paleontological collection was assembled. The specimens were identified by A. M. Obut, graptolites; Yu. I. Tesakov and V. N. Dubatolov, tabulate corals; V. A. Vostokova, gastropods, Z. G. Balashov, nautiloids; N. A. Flerova, stromatopores; R. Ye. Alekseyeva, Devonian brachiopods. The author wishes to express his profound gratitude to them for this work.

Two large tectonic structures in the Paleozoic formations can be identified the Tas-Khayakhtakh Range: the Tas-Khayakhtakh marginal uplift of the Kolyma central massif [12] and the Chibagalakh anticlinorium (Figure 1). They differ in composition and in thickness of component formations and in the pattern and intensity of tectonic dislocations.

Stratigraphy

Paleozoic deposits make up the greater part of the Tas-Khayakhtakh Range. Distinguishable among them are the Ordovician, Silurian, Devonian, and Lower Carboniferous formations interrelated by gradual transitions without any traces of discontinuities or breaks in sedimentation. Resting upon them with sharp angular unconformity are the effusive-sedimentary formations of the Upper Paleozoic and Mesozoic.

Ordovician. Ordovician rocks are abundant in the area discussed. Within the confines of the Tas-Khayakhtakh marginal uplift they are represented by the thick strata of carbonate formations, which includes the Uchugeyan, Usunian, Tagan'yinian, and Kharkindzhinian formations. The first belongs to the Lower, the Usunian and Taganinian — the Middle, and the Kharkindzhinian to the Middle and Upper Ordovician.

¹Ocherk stratigrafii i tektoniki Khrebtu Tas-Khayakhtakh. pp. 61 - 76.




FIGURE 1. Geologic-structural map of the Tas-Khayakhtakh range. Compiled by N.A. Bogdanov using materials of Yu.A. Grebennikov, L.K. Dubovnikov, Yu.T. Krezhevskikh, V.K. Lezhoyev, R.N. Titov, and I.P. Shlykov.

A - Marginal uplifts of the Kolyman central massif: 1 - Precambrian, 2 - Lower Ordovician, 3 - Middle and Upper Ordovician, 4 - Silurian, 5 - Devonian and Lower Carboniferous; B: 6 - Lower and Middle Paleozoic of the Chibagalakh anticlinorium; C: 7 - Middle and Upper Triassic and Jurassic of the In'yali-Debinsk synclinorium; 8 - Mesozoic-Cenozoic deposits of the Zyryanskaya depression, 9 - Upper Paleozoic and Mesozoic of the superimposed basins of the marginal uplifts, 10 - Mesozoic granitoids, 11 - structural lines, 12 - axes of antichinal and synclinal structures, 13 - faults, 14 - Chimalgino-Chibagalakh fault zone, 15 - Datinskaya fault zone. Figures on the map indicate the inner structures of the Tas-Khayakhtakh marginal uplift: 1 - Soloniya anticline, 2 - Vertinskaya syncline, 3 - Yrgandzh anticline, 4 - Kuranakh syncline, 5 - Uchukey anticline, 6 - Berelekh syncline, 7 - Oysordakh anticline, 8 - Dogo graben, 9 - Umba graben-syncline.

Rocks of the Uchukeyan formation are exposed in the valleys of the Uchukey-Yuryakh, Usuna, and Tagan'ya Rivers. They are represented by strata of black clay shales and calcareous shales, alternating with gray clayey and arenaceous limestones. The observable thickness of the formation is 1200 m. In the top part of the section gastropods were collected from the limestones which, in the opinion of V. A. Vostokova appear to be Lower Ordovician. Considering the fact that everywhere the rocks of this formation are conformably overlain by Llanvirnian rocks, we shall tentatively assign them to the Lower Ordovician.

The deposits of the Usunian formation were studied in the valleys of Usun, Uchukey-Yuryakh, and Yrgandzha. Lithologically they differ sharply from the subjacent formations and are represented by greenish-yellow calcareous-marly and marlaceous slates alternating in the upper part of the section with layers and lenses of light-grey medium-laminated limestones. In the Uchukey-Yuryakh and Tagan'ya valleys the formation is about 600 m thick. Further north, in the basin of Yrgandzha River, its apparent thickness increases sharply, attaining 1000-1200 m. The age problem of the Usunian formations deserves special attention, since until recently they were considered as belonging to the Upper Cambrian. This view was based on isolated finds of brachiopods in the Yrgandzha River valley (7, 14) some of which, in the opinion of O. N. Andreyeva (oral statement), were erroneously identified as Upper Cambrian *Billinasella* sp. In 1958 L. K. Dubovikov and in 1959 I myself succeeded in collecting from the same rocks in this formation an assemblage of the trilobites, *Asphidae*, the brachiopods, *Strophomena* sp., and the nautiloids, *Tofangoceras* sp., typical of the Middle Ordovician (Llanvirnian stage). Moreover, in sections along the Uchukey-Yuryakh and Tagan'ya Rivers, the gastropods, *Maclurites* sp., were found in the deposits of the Usunian formation. This also implies that the surrounding rocks are of Middle Ordovician age. All these findings justify the assertion that there are no outcrops of Upper Cambrian rocks within the limits of the Tas-Khayakhtakh uplift.

The formations of the Usunian formation in the Uchukey-Yuryakh, Oyugordakh, and Yrgandzha river valleys gradually merge with the overlying rocks of the Taganian formation. They are represented by grey massive and average stratified limestones penetrated by calcite stringers. Higher in the section, the massive limestones give way to thinly laminated rocks containing rare intercalations of black clay shales and clay limestones. The formation attains a thickness of 1600 m. A collection was made from the Taganian limestones of the brachiopods, *Strophomena* sp., *Rafine-*

squina sp., *Ortidae*; the trilobites, *Asaphidae*; the gastropods, *Lesueurilla* sp., *Pararaphistoma* sp., *Maclurites* sp.; the nautiloids, *Sactorthoceras* sp., and *Geisonoceras* sp., which are characteristic of the Middle Ordovician. On the basis of this collection it is possible to suppose that, the Tanganian formation corresponds in age to the lower and middle parts of the Middle Ordovician Llandeylo stage. The fact that the Tanganian formation is related to the underlying Usunian formation by gradual transition (an alternation of layers belonging to both formations is observable in the transition zone), points even more convincingly to the Ordovician age of the latter.

Kharkindzhinian formation deposits cap the Ordovician section in the Tas-Khayakhtakh uplift and outcrop in the valleys of the Uchukey-Yuryakh, Boryllakha, and Artykh-Yuryakh Rivers. We have described these formations [2]. At the base of the formation one observes black calcareous-argillaceous and argillaceous slates alternating with thin layers of grey platy limestones. The over-all thickness of the Kharkindzhinian formation is 180-200 m. The slates contained the graptolites, *Climacograptus bicornis* (Hall), *Dicellograptus divaricatus* Hall, *Diplograptus multidentatus* Elles and Wood, *Pseudoclimacograptus aff. scharenbergi* (Lapw.), *Orthograptus* sp., *Dicarnograptus* sp., *Dicellograptus* sp., *Dictyonema* sp., and other species whose ages date this formation as of Upper Llandeylo-Ashgill age.

The total observable thickness of the Ordovician within the limits of Tas-Khayakhtakh uplift is about 3900-4100 m.

The Ordovician deposits of the Chibagalakh anticlinorium differ radically lithologically from the coetaneous formations of Tas-Khayakhtakh uplift. They are intensively dislocated and metamorphosed and belong to the Kuntuchnyan formation.² The rocks of this formation are best exposed along the Umba and Kuntuchnya Rivers. Here, the lower half of the section is made up of green muscovite-chlorite and chlorite-quartz schists alternating with interbedded schistose quartz porphyrites and their tuffs (200 m thick), and marmorized limestones. The upper section of the formation is composed of quartz-muscovite, quartz-feldspathic, and quartz-chlorite schists. The total apparent thickness of the Kuntuchnyan formation is approximately 1600 m. Paleontologically, these deposits are poorly characterized. The lower part of the formation contains the stromatopores, *Beatricea* sp., known to occur in the formations of Middle and Upper Ordovician in North America. Considering the

²L. K. Dubovnikov and V. K. Lezhoyev [7] assigned the Kuntuchnyan formations to the Proterozoic.

the fact that the Kuntuchnyan rocks concordantly underlie the coral limestones of the Lower Silurian, we assign them to the Ordovician.

Silurian. The Silurian deposits were studied over the entire area of the region under review. In the Tas-Khayakhtakh uplift they are represented mainly by dolomites and dolomitic limestones. Identifiable in this stage are the Omulevian (Lower Silurian), Umbian (Lower Ludlow), and Datnian (Upper Ludlow - Lower Devonian) formations.

The rocks of the Omulevian formation are exposed in the valleys of the Uchukey-Yuryakh, Boryllakha, and Artykh-Yuryakh Rivers and lie conformably everywhere on the Kharkindzhinian formations [2]. At the base of the Omulevian formation one finds a bed of black clay shales containing thin bands of limestones. Higher in the section they are replaced by limestones containing grey, medium-platy, which at the very top alternate with clay shales. The formation is 220-250 m thick. The slates of the Omulevian formation have yielded the graptolites, *Monograptus* sp., *M. lobiferus* (McCoy), *M. inornatus* (Harkness), *M. ex gr. flemingi* (Salter), *Pseudoclimacograptus nughesi* (Nicholson), *Petalograptus ex gr. palmeus* (Barr.), *Pernorograptus revolutus* (Kurck.), *Rastrites longispinus* Perner, etc., which put the formation in the Llandoveryan - Venlockian age group.

The overlying Umbian series grades into the Omulevian formation. It outcrops along the right tributaries of the Dogdo River, Datna I and Dadyn'ya streams, as well as in the upper reaches of the Umba, Boryllakha, and Solonya. The lower part of the Umbian section consists of dark-grey, massive dolomitic limestones showing thin bands. Higher up they give way to white and light-grey dolomites alternating with bands of yellowish-green and red marl slates and pink dolomites having a thickness of 50-100 meters. The top portion of the section consists of grey, variously laminated dolomitic limestones. In the western and northern parts of the uplift the top sections of Umbian formation contains a 70-100-m bed of coarse- and fine-pebble, brecciated conglomerates. The pebbles and brecciated debris in them originate from the limestones and dolomites in the bottom of the formation. The layer of brecciated conglomerates gradually pinches out in the southeasterly direction. The over-all thickness of the rocks in this formation is 1700-1800 m. The following tabulate corals species were found in the Umbian deposits: *Favosites* cf. *niagarensis* Hall, *Fav. sp.*, as well as stromatopores and gastropods. These fossils date this formation as Lower Ludlow.

Exposures of the Datnian formation deposits occur in the valleys of the Kuranakh-Yuryakh,

and Nakhatta Rivers, as well as along the tributaries of the Dogdo - Datna I, Dadynya, and Khobochalo streams. Everywhere they rest conformably on Umbian rocks. In the Kuranakh-Yuryakh and Nakhatta valleys the base of formation consists of dark-grey argillaceous limestones which, further up the section, change to laminated dolomitic limestones. Layers of organic-clastic detritus are found interbedded in them. The top section of the formation is composed of massive grey limestones interbedded with layers of coral limestones, which usually do not exceed 25-40 m in thickness. The total thickness of the deposits of this formation in these sections amounts to 1000-1100 m. In the western part of the uplift the Datnian sequence exhibits a facies change. Here, the base of the formation consists of grey coral limestones which alternate with layers of medium-laminated dolomitic limestones in the upper parts of the section. The total thickness of the formation decreases to 550-650 m along the course of the Datna I.

The age of the Datnian formations is, in the main, determined on the basis of corals, which, in addition to the Upper Ludlow species, includes *Favosites forbesi* M. Edw. and Haime, *Fav. forbesi* M. Edw. and Haime var. *mammillatus* Tchern., *Fav. aff. coreaniformis* Sok. *Squameofavosites* sp., *Dictyofavosites aff. tchernajaensis* Dubat. One also finds the Lower Devonian specimens: *Striatopora tschichatschewi* Peetz., *Parastriatopora rzhonitskajae* Dubat. and *Pseudomicroplasma* sp. This is the basis for believing that the Datnian series is of Upper Ludlow - Lower Devonian age.

The over-all thickness of the Silurian deposits in the eastern section of the uplift reaches 3000-3100 m and decreases to 2300-2500 m in the western part.

In the Chibagalakh geanticline the Silurian sequence is changed. In the upper reaches of the Umba and Tukhana Rivers beds of metamorphosed limestones and slates of the Tukchanian formation lie conformably on the Kutuchnyan metamorphic schists. The bottom of the former is predominately quartz sandstones and marmorized limestones interstratified with quartz-sericitic schists. The upper part of the section consists mainly of marmorized limestones containing thin bands and lenses of phyllite and quartz-sericitic schists. The Tukchanian formation are about 2000 meters thick. The following fossils were collected from the base of the formation in the marmorized limestones: *Favosites* sp., *Palaeofavosites longispinus* Tchern., *Mesofavosites* sp., *Syriogopora* sp., *Catenipora aff. parallelus* Schmit., *C. aff. gothlandicus* Yabe, *C. vulgaris* Tchern., the most of which are typical of the Llandoveryan stage of the Lower Silurian. Discovered in the middle part of the section were the tabulate corals, *Favosites forbesi*, *M. Edw.* and *Haime*

var. *coreanicus* (Ozaki) and Fav. sp. which characterize the age of the overlying formations as belonging to the Wenlock-Ludlow stages. On the basis of the foregoing data we consider the age of the Tukchanian formation to be Silurian.

Devonian. All three Devonian formations, interrelated by gradational transition, are identifiable within the limits of the Tas-Khayakhtakh uplift. In recent years they were described by L. K. Dubovikov [7], and studies in detail by R. Ye. Alekseyeva and V. N. Dubatolov, whose materials, together with our own observations, are utilized in this article.

Lower Devonian deposits outcrop along the banks of the Dadyn'ya, Khobochalo, and Kuranakh-Yuryakh Rivers. In the latter's valley they occur as dolomite beds of average thickness and dolomitic limestones whose strata contain individual beds of coral limestones attaining 20 to 25 m in thickness. The Lower Devonian deposits here are about 600-650 m thick. Further west the sequence of the coetaneous deposits is somewhat modified. Along the tributaries of the Dogbo River and in the valleys of the Khobochalo and Dadyn'ya streams the base of the section is made up of grey, banded, coral limestones. Higher up, they give way to grey and dark-grey dolomites, which contain the tabulate corals: *Syringopora abdita* vern., *S. aff. borealis* Tchern., and the brachiopods, *Delthyris tenuisnuosus* Khod., *D. ex. gr. tiro* Barr., etc. The over-all thickness of this section is about 350-400 m.

Eifelian deposits are exposed in the Dogdo River basin. They are represented by black calcareous clay shales and slates alternating with thin bands of dark-grey shaly limestones containing the brachiopods, *Gypidula aff. ivdelensis* Khod., *Carinata raris* Ržon., *Spinatrypa ex gr. aspera* Schloth., *Atrypa ex gr. reticularis* L., *Eospirifer irbitensis* Tchern.; the tabulate corals; *Pachyfavosites* sp., *Alveolitella* sp., *Favosites aff. goldfussi* Orb., *Fav. robustus* Lec., *Syringopora cf. crispa* Schluter, etc.; as well as specimens of trilobites, crinoids, and tetracorals, characteristic of the Eifelian stage of the Middle Devonian. These deposits were found to vary in thickness from 250 to 500-600 m.

Givetian deposits were studied along the Khobochalo, Sebechan, and Dadyn'ya streams. They are represented by grey and light-grey dolomitic limestones and interbedded dolomites containing the tabulate corals; *Scoliopora ex gr. denticulata* (M. Edw. and Haime), *Thamnopora ex gr. polyforata* (Schlothem), *Alveolitella* sp., *Cladopora* sp., *Syringoporella* sp., *Clathrodictyon* sp., and the brachiopods; *Stringocephalus burtini* Defr. In the top of the sequence the dolomites alternate with individual

lenses and non-uniformly striking bands of red and greenish-yellow marls, calcareous-marly slates and anhydrites, varying in thickness from 10-15 to 200-250 m (the right-headwaters of the Tuostakh River, [7]). The total thickness of the Givetian stage deposits is about 450-500 m.

Upper-Devonian deposits are represented by grey and light-grey, medium-laminated dolomitic limestones. They were encountered in the valleys of the Emtachan and Dadyn'ya Rivers. However, these sections fail to display the youngest formations of the Upper Devonian. In the dolomites of the Emtachan valley, L. K. Dubovikov [7] succeeded in collecting specimens of *Nalivkinella* sp., *Thamnopora nalivkina* Tchern., *Th. reticulata* Blainv., *Th. sp.*, *Alveolitella* (?) aff. *karmakensis* Tchern., which give evidence of the Upper Devonian age of these formations. The observed thickness of the Upper Devonian deposits attains 100-110 m. The over-all apparent thickness of the Devonian in the Tas-Khayakhtakh uplift is about 1500-1800 m.

The Biyergichekian formation is a formation of the Lower and Middle Paleozoic in the general area of the Chibagalakh anticlinorium.³ The Biyergichekian rocks are less metamorphosed than are the Tukchanian and Kuntuchnyan formations, and are in contact with the latter along the faults. The Biyergichekian rocks are best exposed in the valley of the Biyergichek River and along its tributaries. The basal strata of the formation here consist of reddish-brown quartz sandstones with thin bands of phyllite slates. The middle of the formation consists of quartz-like sandstones and phyllite slates, and the upper part is entirely composed of black and dark-grey phyllite slates. The total observed thickness of these deposits is approximately 2000 m. The age of the Biyergichekian formation has not been definitely established.

Carboniferous. Carboniferous deposits were investigated only in the central portion of the Tas-Khayakhtakh uplift along the tributaries of the Dogdo River in the valleys of the Uklin and Dzhabul'denga. Their relation to the subjacent rocks, so far, have not been determined. The formations of this age are represented by grey, laminated limestones interstratified with black argillaceous and calcareous-argillaceous shales. In the limestones L. K. Dubovikov and R. Ye. Alekseyeva found the brachiopods, *Spirifer cf.*

³Earlier L. K. Dubovikov and V. K. Lezhoyev [7] assigned this formation to the Sinian. According to the new data produced by V. K. Lezhoyev and G. N. Loginova (verbal report) the Biyergichekian formation should be considered as a stratigraphic equivalent of the Kuntuchnyan formation and attributed to the Ordovician.

ornacensis Kon., *Linoproductus* cf. *panderi* Ueberl., *Stereophrentis* ex. gr. *disjuncta* (Carr), and other fossils of the Tournaisian and the lower part of the Visean stages. The apparent thickness of the Carboniferous deposits is 50-400 m.

In summing up the above description of the Paleozoic sections in the Tas-Khayakhtakh Range, it seems important to dwell on some of their special features. Within the limits of the Tas-Khayakhtakh marginal uplift, just as in the uplift of the Cherskiy Range [4], the Paleozoic formations are represented chiefly by the rock of the carbonate series. The abruptly increasing thickness of the Upper Silurian and Devonian formations in the eastern part of the uplift (Kuranakh-Yuryakh and Berelekh River basins - Figure 2) is attributable to the fact that it was here that the central and most downwardwarped part of the large Paleozoic depression extending from the Omulyevskiy Mountains in the Southwestern part of the Kolyma central massif [3] and stretching out in the southwesterly direction, was located. A more complex Paleozoic structure exists within the confines of the Chibagalakh geanticline. The Ordovician rocks are represented here by terrigenous formations and marmorized limestones containing rare bands of schistose acid effusives, whereas in Tas-Khayakhtakh uplift, as mentioned earlier, limestones and marls predominate. The facies of the deposits of the Lower Silurian of these two structural zones are also different. If, in the marginal uplift they consist of a thin band of graptolite schists, in the Chibagalakh anticlinorium the Lower Silurian is represented by a thick stratum of coral limestones. It is to be regretted that it is difficult, at the present time, to compare the Biyergichekian formation with any known formation, since it has not yet been paleontologically defined. On the whole, the over-all composition of the Paleozoic rocks in the Chibagalakh anticlinorium suggests that the sedimentation conditions here were different from those in the adjacent Paleozoic trough in the southeastern part of the Kolyma central massif.

Upper Paleozoic and Mesozoic. The Upper Paleozoic and Mesozoic formation occur in the Dogdo graben and in the Umba graben-syncline. They are thoroughly described by L.K. Dubovikov and V.K. Lezhoyev [7], as well as by A.V. Zimkin [8], on the basis of materials collected by V.P. Fagutov, D.P. Vas'kovskiy, V.K. Lezhoyev, and L.K. Dubovikov. We shall refer to them very briefly in this paper.

In the Khobochalo stream basin in the northwestern part of the Dogdo graben the Silurian and Devonian rocks are transgressively and unconformably overlain by Lower Permian conglomerates and arenaceous limestones, whose observed thickness is 70-80 m. Further

south in the valley of the Krest-Yuryakh River, also lying unconformably on Middle Paleozoic limestones, there is a 250-300 m formation of shales, tuffaceous sandstones, diabase porphyrites and their tuffs containing Middle Triassic ammonites. The relationship between the Permian and Triassic deposits in the Tas-Khayakhtakh Range was not determined. The described Paleozoic rocks, as well as the Permian and Middle Triassic formations, are unconformably overlain by Upper Jurassic basic and acid effusives and their tuffs, containing shale and sandstone layers. It is mainly these formations which fill the Dogdo graben and the other graben-like structures in the Tas-Khayakhtakh Range. In the Kyra River valley, in the northern part of the Dogdo graben, in the formation of shales in the lower part of the section, one observes lenses of brown coal about 0.6 m thick. The total thickness of the Upper Jurassic deposits in the Dogdo graben is 2000 m. The younger rocks covering the Paleozoic and Mesozoic deposits are the Tertiary and Quaternary alluvial and deluvial formations which are widespread in the Zyryanskaya depression which bounds the Tas-Khayakhtakh uplift on the northeast.

Tectonics

As stated above, the principal tectonic elements of the Paleozoic structure in the Tas-Khayakhtakh Range are the Tas-Khayakhtakh marginal uplift of the Kolyma central massif, and the Chibagalakh anticlinorium (see Figure 1), which differ in the composition and thickness of the component rocks, as well as in the intensity and type of tectonic dislocations.

The Tas-Khayakhtakh uplift is the western extension of the Kolyma central massif which extends from the Indigirka River valley in the northwest, then in a near-meridional direction to the valley of the Selenyakh River. Its total length is 300 km with a maximum width ranging from 50 to 70 km. On the west, the Tas-Khayakhtakh uplift is bounded by the Chimalgino-Chibagalakh fault zone which separates it from Chibagalakh anticlinorium and the In'yali-Debinsk synclinorium. In the east, the uplift abuts on Zyryanskaya depression which is filled with Mesozoic and Cenozoic formations. The Lower and Middle Paleozoic rocks which comprise the uplift are contorted into large folds and are cut by numerous strike and diagonal faults which are responsible for its complex block-fold structure.

The following tectonic structures from east to west, trend in a northwesterly direction, can be identified within the limits of the marginal uplift: (Figure 1): Berelekh brachyanticline [6], Kuranakh syncline [4], Uchugey antiline [5], Yrgandzhi anticline [3], Oysordakh anticline [7], Vertinskaya syncline [2], and the Solon'ya anticline [1]. The two latter structures are separated

Chibagalakh anticlinorium
Umba and Biyergichek
River basins

Tas-Khayakhtakh Uplift
Dogdo River basin




FIGURE 2. Comparison of the stratigraphic columns of the Paleozoic in the Tas-Kyakhtakh Range.

1 - limestones, 2 - shales, 3 - marls, 4 - dolomites, 5 - coral limestones, 6 - metamorphic schists, 7 - schistose effuses, 8 - marmorized limestones, 9 - quartzites, 10 - fossil locations.

by the Dogdo graben [8] extending to the Umba graben-syncline. Both are filled with Upper Paleozoic and Mesozoic effusive-sedimentary formations.

The Berelekh brachyanticline extends from the Suordakh River valley to the Middle course of the Nakhata River for a distance of 50 km, and has a width of 25 to 30 km. This structure can be described as dome-shaped, and it is somewhat elongated in the meridional direction. The center of the brachyanticline consists of rocks of the Datnian formation (Figure 4). They form a gentle arch with an 8-15° pitch. Toward the limbs, the dip becomes steeper and reaches an angle of 25-30°. The eastern limb of the brachyanticline is truncated by a fault which separates it from the Zyryanskaya depression. In the fault zone the rocks are crumpled into minor folds tilted toward the east. The width of the folds in the fault zone does not exceed 0.5-0.7 km. The western limb of the Berelekh brachyanticline is also complicated by a longitudinal fault separating it from the Kuranakh syncline.

The Kuranakh syncline is traceable from the middle course of the Nakhata River to the upper Kyra River, a distance of more than 100 km. The syncline is 30 - 40 km wide. The eastern limb of the structure is everywhere truncated by faults separating it from the Zyryanskaya depression and the Berelekh brachyanticline. The Upper Silurian rocks comprising the western limb of the syncline dip in a northeasterly direction at an angle of 25-30° as a monocline which gradually flattens out toward the east. The syncline core is composed of nearly horizontal Datnian formations. Only gentle bending of the beds is observed in this part of the structure.

In the Nakhata River valley, following the trend of the Kuranakh syncline, and south of its trochoclinal closure, lies the Uchugey anticline.

The Uchugey anticline has an almost meridional strike and is traceable from the Nakhata River valley to the Umba River. It is 70 km long, and has a width of about 25 to 30 km. Here, we were able to investigate only the western part and the northern periclinal closure of the anticline. The eastern limb and the southern periclinal closure of the fold plunge step-by-step along the longitudinal faults under the formations of Zyryanskaya depression. The Ordovician rocks which make up the western portion of the core and the anticline limb form a monocline where the dip of the beds increases from east to west from 25-30° to 45-50°. In its western marginal section the monocline is complicated by faults and upthrusts with negligible throw. In this faulted zone, laminated limestones and shales are contorted into small flexures, while the more

massive rocks are usually slightly recrystallized and crushed. The pattern of the northern periclinal closure of the Uchugey anticline is more complex. The component Taganian formation limestones are contorted into a number of minor folds whose bends have a steep northward pitch. The dips of these folds vary from 25° to 40°.

The Yrgandzhi anticline is situated northwest of the Uchugey anticline, forming a step-like pattern with the latter. It is bounded on all sides by faults and extends in an almost meridional direction from the Sebechan River to the Boryllakh River for a distance of 35-40 km, and is 8-10 km wide. The bend of the structure exhibits a northward rise, and here in the Yrgandzha and Inarodzha River valleys the Ordovician rocks of the Usunian and Taganian formation in the core are exposed. They form a steep crest-like antinormal fold, whose limbs gradually flatten out to the west and east (see Figure 3). The dip of their beds changes from 50°-60° to 25°-30°. Further south, on the upper reaches of the Artykh-Yuryak River the anticline core has a less complex structure. Here, the Ordovician formations form a gentle arch with the strata dipping at an angle of 25°-30°. The limbs of the Yrgandzha anticline are composed of Silurian rocks and everywhere display a simple structure. The component rocks plunge at an angle of 10°-20° and are crumpled into small complex folds only as they reach the edges of the faults. The fault which bounds the Yrgandzhi anticline on the west is the border between it and the Mesozoic Dogdo graben and the Vertinskaya syncline.

The Vertinskaya syncline is composed of Upper Silurian and Devonian rocks. It is 10-12 km wide and extends for 40 km from the upper Kyra river on the north to the Inarodzha River on the south. On all sides it is surrounded by ruptured zones. On the whole, the Vertinskaya syncline has an oval shape. Its limbs are complicated by minor and larger folds. Thus, the rocks of the syncline's western limb in the valley of Datna I stream, in the ruptured zone which separates the syncline from the Dogdo graben, are contorted into small folds manifesting a dip of 70°-80°. In individual blocks and wedges it was possible to observe a westward overturning of folds in the direction of the graben. The eastern limb of the Datna syncline is much simpler in structure. The Umbian limestones, of which it is composed, form a monocline, which dips 40°-50°, complicated by a sloping box fold of secondary order. The syncline's crest rises in the northwest, and here the northern trochoclinal closure of the trough is replaced by the tectonic elements of the Oysordakh anticline.

The Oysordakh anticline is located in the northeastern part of the Tas-Khayaktakh uplift




FIGURE 3. Schematic geologic sections across the Tas-Khayakhtakh Range.

Tas-Khayakhtakh marginal uplift of the Kolyma central massif: 1 - Lower Ordovician, 2 - Middle and Upper Ordovician, 3 - Silurian, 4 - Devonian, 5 - Middle Triassic and Upper Jurassic.

Chibagalakh anticlinorium; 6 - Tukchian formation, 7 - Biyergichekian formation, 8 - Triassic of the In'yali-Debinsk synclinorium, 9 - Mesozoic granitoids, 10 - deep-seated faults, 11 - dislocated zones.

between two Mesozoic granite batoliths — the Khadaryninskiy [10] in the east, and the Tirekhtykhinskiy, in the west — and was described by P. N. Titov. The inner anticline structure is complicated by two discontinuous dislocations: the northwestern and the southeastern. The first is traceable along the entire Tas-Khayakhtakh uplift and separates it from the Zyryanskaya depression. The second is the western continuation of the faults which fringe the Polousenskoye uplift on the south. They cut the Oysordakh anticline into a number of separate tectonic blocks trending to the northwest. The anticline's core is composed of Ordovician and Silurian rocks, with Devonian formations on its limbs. The limbs have a 60° to 75° dip. In the zones adjacent to the faults, the Lower and Middle Paleozoic deposits are recrystallized and intruded by a large number of diabasic dikes.

The most complex structure occurs in the western part of the Tas-Khayakhtakh uplift where the Solon'ya anticline is situated. On the west it is in contact with the In'yali-Debinsk aynclinatorium along a fault, and is adjacent to the Dogdo graben on the east. It extends southeast - northwest for almost 80 km, maintaining a width varying from 10 to 12 km. The Solonya anticline is a narrow linear fold, its limbs and the northwestern periclinal closure complicated by faulting. Umbian rocks belonging to the Lower Ludlow form the core of the anticline. They are folded and crumpled into steep (dip of the beds, about 70°) symmetrical folds trending northwest, are complicated by faults and upthrows with a displacement of several hundred meters. The western limb of the Solonya anticline is composed of Upper Ludlow and Lower Devonian rocks and is complicated by narrow, linear, second order faults, showing bed dips attaining 75° - 80° . The Devonian limestones which make up the eastern limb of the structure form a monoclinical flexure, ruptured by numerous longitudinal faults. Only in the ruptured zone is the monocline complicated by minor folding.

The Umbian graben-syncline and the Dogdo graben are located in the central and southern parts of Tas-Khayakhtakh marginal uplift. They are the northernmost structures in the narrow belt of Mesozoic troughs traceable from Lake Darpir in the southeast, to the basin of the Kyra River on the northwest. The Mesozoic troughs extend along a system of major faults which delineate the central massif and are superimposed on the Paleozoic folds of the marginal uplifts (see Figure 3).

The Umbian graben-syncline is confined to the Chimalgino-Chibagalakh fault zone and strikes in almost latitudinal direction for some 35 km. Its maximum width is 5-8 km. The inner part of this structure is complicated. In cross-section, it is sharply asymmetrical

because of the general dip and occasional southwesterly overturn of the layers in the direction of the Chimalgino-Chibagalakh fault zone. The Umbian graben-syncline is not structurally uniform. In its central and northeastern parts, the Upper Jurassic effusives manifest a gentle 5° - 7° dip, like a monocline sloping in a south-southwesterly direction. The Middle Triassic shales and sandstones and Upper Jurassic effusives, which comprise the western section of the graben-syncline, abut on the fault zone and are crumpled into a series of small folds, sometimes tilted southward with a dip varying from 50 - 70° to 15 - 25° . At the contact with the faults the rocks are recrystallized and schistose and form dislocation breccia and slicken sides. Along a line from the east to the northwest, the graben-syncline narrows abruptly, and here it is connected with the Dogdo graben by a narrow (2-3 km in width) graben extending along the Datnian fault. The graben is composed of Middle Triassic and Upper Jurassic rocks forming an asymmetrical fold, the most downwarped part of which is coordinated with the Datnian fault. The Mesozoic formations are intensively cataclasticized in the fault zone.

As described earlier, the Dogdo graben is situated further to the north in the central part of the Tas-Khayakhtakh uplift. It extends in northwesterly direction from the upper reaches of the Goredomchan-Yuryak River to the latitudinal bend of the Dogdo River. Its width attains 30-35 km and its length totals 100 km. Permian rocks comprise the western limb, while the central and eastern sections are composed of Upper Jurassic deposits unconformably superposed on Middle Paleozoic rocks. On the northeast, the Dogdo graben is bounded by the Datnian fault. On the opposite, southwestern side, the graben is also truncated by faults, although of lesser importance. Asymmetrical structure is, on the whole, characteristic of this graben, just as it is for the Umbian graben-syncline. In its 20-25 km wide western part, the Permian and Upper Jurassic rocks have a relatively gentle, 3° - 6° eastward dip. They form a gentle monocline complicated by small gentle quaquaversal folds. The more folded, 5-8-km wide eastern part of the Dogdo graben exhibits more complex structure. The Upper Jurassic effusives are crumpled here into small folds, whose limbs dip at increasing angles in an easterly direction, from 10-15 to 25 - 30° , as the folds approach the Datnian fault.

Thus, the Tas-Khayakhtakh marginal uplift constitutes a complex block-and-fold structure consisting of a number of major Paleozoic folds and superimposed Mesozoic structures of the graben and graben-syncline type.

One should dwell a little longer on some of the general features of the uplift structure (see Figure 3). Large folds with limbs extending

for 25-30 km are developed in its eastern part. In the central parts of these folds the rocks are almost horizontal, dipping at an angle of 15-30° on the limbs. The latter are usually faulted, emphasizing the block-and-fold pattern of this territory. In the western part of the Tas-Khayakhtakh uplift, this type of folding gives way to linear structures traceable for 30-40 km, over a 5-7 km wide belt. The limbs of these folds dip steeply at a 50-70° angle, and in the cores of crest-like anticlines the beds are vertical. The width of this zone of linear folding does not exceed 25-30 km.

The foregoing data indicate that the degree of dislocation affecting the Paleozoic rocks in the Tas-Khayakhtakh uplift gradually increases from east to west away from the central part of the Kolyma central massif. Permian and Mesozoic rocks are almost horizontal within the limits of the uplift and are dislocated only in the fault zones.

The Chibagalakh anticlinorium is the eastern element of the Yana-Kolyma megasyncline [13] and is located on the southwestern slope of the Tas-Khayakhtakh Range. It covers an area 35-50 km wide and 150-200 km long stretching from the Khara-Sal valley to the lower course of the Chibagalakh anticlinorium borders on the Inyali-Debinsk synclinore which is filled with Triassic, Upper, Middle, and Lower Jurassic deposits aggregating 3000-4000 m in thickness [5]. In the north and northeast, the geanticline is bounded by the Chimalgino-Chibagalakh fault zone which separates it from the Tas-Khayakhtakh uplift of the Kolyma central massif. The Chibagalakh anticlinorium is formed by intensely dislocated Paleozoic rocks and is broken up by widespread longitudinal and diagonal faulting into a number of horsts and grabens (see Figure 1), which make it difficult to differentiate its structure as a whole.

The central part of the geanticline is composed of Ordovician metamorphic rocks, with Silurian deposits on its limbs.

The metamorphic schist and marmorized limestones of the Kuntuchnyan formation which make up the Chibagalakh anticlinorium's core are contorted into complex isoclinal folds whose limb spread out for 50-100 m. The beds are tilted to the northeast at an angle of 40-60°. In that part of the core which is directly contiguous to the Chimalgino-Chibagalakh fault zone, the dip of the isoclinal folds becomes steeper, attaining an angle of 75°-80°. Here, the folds rarely exceed 25-35 m in width.

The anticlinorium's eastern limb is formed by Tukchanian limestones and is cut by diagonal faults into separate blocks plunging in a step-like pattern to the east. The rocks are folded

here into recumbent folds overturned to the northeast at an angle of 50°-60°. The spread of their limbs does not exceed 50-150 m, and their axes have a steep pitch, so that in longitudinal section these folds have a box-like shape. The western limb of the Chibagalakh anticlinorium is composed of the Tukchanian and Biyergichekian formation and is separated from the core of the anticlinorium by major faults having latitudinal and northeastern strikes. There is less intense dislocation in this portion of the structure than in its core and eastern limb. The western limb is complicated by regular linear, both symmetrical and asymmetrical, folds which, as a rule, are overturned in the southeasterly direction (see Figure 3). Usually, asymmetrical folding can be observed in the zones of the minor faults which complicate the limbs of second and third order folds. On the middle course of the Khara-Sal River, the western limb of the Chibagalakh anticlinorium is in contact with the tectonic elements of the In'yali-Debinsk synclinore along the fault. The internal structure of this synclinore has been described by several authors [13, 5, 8].

As can be seen from the above brief description, the Chibagalakh anticlinorium is structurally far more complex than the Tas-Khayakhtakh marginal uplift of the Kolyma central massif. There is no isoclinal folding similar to that which complicates the structure of the Chibagalakh geanticline, anywhere within the limits of the marginal uplift. Moreover, there are almost no recumbent narrow folds in the marginal uplift. Common to these two major folded systems are the numerous faults responsible for their block-and-fold structural pattern.

The Chimalgino-Chibagalakh fault zone which we defined, separates, as previously mentioned, two different tectonic elements: the central massif, marginal uplift of the Tas-Khayakhtakh and the Chibagalakh geanticline, located within the confines of the geosynclinal zone. This zone extends from the middle course of the Chibagalakh River to the Boldymbinskiy batholith for 350-400 km in a northwesterly direction, and controls in the south, the narrow belt of Mesozoic effusives. Along the Chimalgino-Chibagalakh fault zone, following the contact between the Jurassic effusives and the Paleozoic rocks, there is a thick (up to 100 m thick) band of fault breccia and mylonites. The effusives are more highly crushed and altered here. The marmorized Paleozoic limestones and metamorphic schists are recrystallized and particularly intricately corrugated in the fault zone. The presence of fault breccia and mylonites in the Jurassic effusives and Paleozoic rocks indicates that that development of the Paleozoic formations was of very long duration. The Chimalgino-Chibagalakh fault zone apparently, must have

been formed in the early Paleozoic at the boundary between two large structures, and maintained its tectonic activity throughout the entire Paleozoic history of this region. This may, no doubt, explain the difference between the Paleozoic sections of the Tas-Khayakhtakh uplift and the Chibagalakh anticlinorium.

CONCLUSIONS

We shall now deal with a few basic problems relative to the development of the tectonic structures in the investigated area.

There are no outcrops of Pre-Ordovician formations within the limits of the Tas-Khayakhtakh Range, and we are, therefore, unable to dwell at the present time on the older stages of the region's tectonic history. As already mentioned, the formation of a depression filled mostly with carbonate deposits occurred within the confines of Tas-Khayakhtakh uplift — as well as over the entire southwestern area of the Kolyma central massif — in the period embracing the Lower Ordovician to the Lower Carboniferous, inclusively. Another structure-facies zone was located further west from this downwarp in the region of the Chibagalakh anticlinorium and, possibly, also over the greater part of the In'yali-Debinsk syncline. Thick terrigenous strata containing individual beds of acidic effusives and their tuffs, are characteristic of this zone. The boundary between these two areas ran along the Chimalgino-Chibagalakh fault zone.

There was no sedimentation in the Middle and Upper Carboniferous within the limits of the presently reviewed central section of the Mesozooids in Northeast Asia. At the same time, Permian formations are found deposited with a mark angular unconformity on the older Paleozoic deposits. On the basis of this fact, it is possible to postulate that the formation of the principal Paleozoic folded structures was terminated in the Pre-Permian epoch. The amount of dislocation affecting the Paleozoic rocks in the Tas-Khayakhtakh uplift gradually increases from east to west, and formations of this age are very complexly contorted within the limits of the Chibagalakh geanticline. It appears, that in the Middle and Upper Carboniferous the tectonic movements were most intensely manifested in the eastern part of the Yana-Kolyma megasyncline, gradually decreasing in the direction of the ancient core of the Kolyma central massif.

Throughout the entire Mesozoic history — that is, during the long period of time when the recent structures of the Yana-Kolyma megasyncline were in the process of formation — the Tas-Khayakhtakh uplift must have reacted to tectonic movements by forming individual horsts and grabens and by complicating the folded structure of its western marginal section. The

Datnian fractured zone, along which — like the faulted zone of Chimalgino-Chibagalakh — there occurred a discharge of effusives in the Middle Triassic and Upper Jurassic, probably must have also developed during this period. Thus, at this stage of development, the Tas-Khayakhtakh marginal uplift — as well as the entire Kolyma central massif — was a fixed elevation to the outlines of which the folded structures of the geosynclinal zone have adapted themselves in, so to speak, a streamlined pattern. The appearance of primarily block-type dislocations within the uplift in Mesozoic time is also indicated by the horizontal bedding, within its limits, of Upper Paleozoic and Mesozoic deposits. The latter are dislocated only in fault zones where the vertical movements appear to have continued into Post-Cretaceous time.

REFERENCES

1. Atlasov, I. P., *Geologicheskoye issledovaniye rayona Sennnyakho-Indigirskikh gor i yuzhnykh otrogov khr. Poluosnogo. [GEOLOGIC INVESTIGATION OF THE REGION OF THE SELENYAKHO-INDIGIRKA MOUNTAINS AND OF THE SOUTHERN SPURS OF POLUOSKIY RANGE]: Tr. Vses. n. -i. Arkt. in-ta, t. 99, 1938.*
2. Bogdanov, N. A., *Stratigrafiya verkhnego ordovika i nizhnego silura yuzhnoy chasti khrebt Tas-Khayakhtakh (khr. Cherskogo) [STRATIGRAPHY OF THE UPPER OROVICIAN AND LOWER SILURIAN IN THE SOUTHERN PART OF THE TAS-KHAYAKHTAKH RANGE (CHERSKIY RANGE)]: Dokl. Akad. Nauk SSSR, t. 127, No. 2, 1959.*
3. Bogdanov, N. A., *Stroyeniye paleozoy-skogo progiba yugo-zapadnoy chasti Kolymaskogo sredinnogo massiva. [STRUCTURE OF THE PALEOZOIC TROUGH IN THE SOUTHWEST PART OF THE KOLYMA CENTRAL MASSIF]: Dokl. Akad. Nauk SSSR, t. 132, No. 3, 1960.*
4. Bogdanov, N. A., and M. N. Chugayeva, *Paleozoyyskiye otlozheniya Omulevskikh gor. [THE PALEOZOIC DEPOSITS IN THE OMULEVSKIYE MOUNTAINS]: Izv. Akad. Nauk SSSR, ser. geol., No. 5, 1960.*
5. Gavrikov, S. I., *O tektonicheskom rayoni-rovanii basseyna verkhnego techeniya r. Indigirki. [TECTONIC ZONES OF THE UPPER REACHES OF THE INDIGIRKA RIVER BASIN]: Tr. Vses. n. -i. in-ta zolota i redk. metallov, vyp. 38, geologiya, 1958.*
6. Demokidov, K. K., *Geologicheskoye stroyeniye i poleznyye iskopayemyye severnoy chasti khrebt Tas-Khayakhtakh*

- i privileguyushchego k nemy pravoberezh'-
ya r. Yany. [GEOLOGIC STRUCTURE AND
AND INDUSTRIAL MINERALS IN THE
NORTHERN PART OF THE TAS-
KHAYAKHTAKH RANGE AND THE AD-
JACENT RIGHT BANK OF THE YANA
RIVER]: Tr. Vses. n.-i. Arkt. in-ta,
t. 142, 1939.
7. Dubovikov, L.K., and V.K. Lezhoyev,
Stratigrafiya paleozoyskikh i mezozoy-
skikh otlozheniy khrebt Taz-Khayakhtakh.
[STRATIGRAPHY OF THE PALEOZOIC
AND MESOZOIC DEPOSITS IN THE TAS-
KHAYAKHTAKH RANGE]: Tr. Mezhdoved.
soveshch. po razrab. unifits. stratigraf.
skhem. sev.-vost. SSSR, 1957. Dok-
lady, 1959.
8. Zimkin, A.V., Geologicheskoye stroyeniye
Severo-Vostochnoy Yakutii. [THE GEO-
LOGIC STRUCTURE OF NORTHWEST-
ERN YAKUTIYA]: Tr. Yakut. fil. Akad.
Nauk SSSR, ser. geol., sb. No. 3, 1959.
9. Matveyenko, V.T., and Ye.T. Shatalov,
Razryvnyye narusheniya, magmatizm
i orudneniye Severo-Vostoka
SSSR. V sb. "Zakonom. razmeshch.
poleznykh iskop.", [FAULTING, MAG-
MATISM, and MINERALIZATION IN
THE NORTHWESTERN PART OF THE
USSR. From the collection, "Regularities
in the Emplacement of Industrial Min-
erals"]: T. 1, 1958.
10. Nekrasov, I.Ya., and V.I. D'yachenko,
Osnovnyye cherty geologicheskogo stro-
yeniya i metallogenii Nizhne-Indigirskogo
rudnogo rayona. [THE MAIN FEATURES
OF THE GEOLOGIC STRUCTURE AND
METALLOGENY IN THE LOWER IN-
DIGIRKA MINING DISTRICT]: Sov.
geologiya, No. 1, 1960.
11. Nikolayev, A.A., Skhema stratigrafii
dokembriya, nizhnego i srednego paleo-
zoya Severo-Vostoka SSSR. [STRATI-
GRAPHIC MAP OF THE PRECAMBRIAN,
LOWER AND MIDDLE PALEOZOIC
FOR NORTHWEST USSR]: Tr. Mezhdoved.
soveshch. po razrab. unifits. stratigraf.
skhem. Sev.-Vost. SSSR, 1957 g. Dok-
lady, 1959.
12. Pushcharovskiy, Yu.M., O tektonike
Severo-Vostoka SSSR. [THE TECTONICS
OF NORTHWESTERN USSR]: Byul.
Mosk. o-va ispyt. prirody. Otd. geol.,
t. 31, No. 5, 1956.
13. Pushcharovskiy, Yu.M., Priverkhoyan-
skiy krayevoy progib i mezozoidy Severo-
Vostochnoy Azii. [THE PRIVERKHOY-
ANSK FOREDEEP AND THE MESOZOIDS
OF NORTHWEST ASIA]: Avtoreferat
Dokt. diss., M., 1958.
14. Fedortsev, V.A., Geologiya i poleznyye
iskopayemyye Yano-Indigirskogo vodora-
zdela. [GEOLOGY AND INDUSTRIAL
MINERALS OF THE YANA-INDIGIRKA
DIVIDE]: Tr. I geol.-razved. konf.
Glavsevmorputi v aprele 1935, g., t. 2,
1936.

Geological Institute of the
U.S.S.R. Academy of Sciences,
Moscow

Received June 3, 1960

THE FACIES, DYNAMIC PHASES AND FORMATIONS OF ALLUVIUM¹

by
I. P. Kartashov

Studies of alluvial deposits are of great practical importance for the exploration, prospecting, and mining of placer mineral deposits, for building dams, canals, roads, and other construction projects, and also for many purposes in other fields of the national economy. The considerable successes achieved in this research during recent years do not, however, offset very substantial shortcomings. We cannot, for example, fail to point out the fact that the genetic peculiarities of alluvium, which this article is devoted, are still very poorly studied. Yet, many of these features, while considerably affecting the lithology of alluvium, are also of great importance to stratigraphic and paleogeographic development.

The factual material upon which the positions taken in this paper were formulated is based on the data of numerous exploratory and prospecting operations conducted in alluvial gold placers in different parts of the Soviet Union.

Facies of Unconsolidated Deposits

Present-day genetic studies of unconsolidated deposits are no longer, as a rule, confined to mere classification into genetic types. "Facies" has now become the most commonly used designation for the genetic subdivisions in the succeeding order serving as the principal object of study. However, the definition of facies as a lithologically homogeneous complex of deposits [9, 13] is hardly applicable to continental, unconsolidated deposits distinguishable, as is well-known, by excessive variations of lithologic characteristics. A reference to S. A. Yakovlev's genetic classification of Quaternary deposits [13] can furnish a good illustration of this fact. None of the continental facies identified therein can satisfy the requirements of lithologic uniformity on which S. A. Yakovlev so emphatically insists [3, pp. 41-43].

The term unconsolidated sedimentary facies properly should be applied to complexes of rocks of the same genetic type formed under identical geomorphologic conditions. The relating of unconsolidated deposits to one and the same facies must, therefore, be determined by their coincidence with the same development stages or relief forms.

An overwhelming majority of investigators consider alluvium as an independent genetic type. Since the publication of Ye. V. Shantser's study [12] on the alluvia of plains rivers, it has become common to class alluvial deposits into three genetic complexes: channel, flood-plain, and oxbow complexes. Each of these complexes fully corresponds to the above definition of unconsolidated sedimentary facies. In S. A. Yakovlev's classification [13] these complexes are, indeed, called facies. Ye. V. Shantser actually considers channel, flood-plain, and oxbow alluvium as "groups of facies". He distinguishes, for example, in channel alluvia the "facies" of near-channel shoals, bars, race and perluvial "facies". However, such fine genetic subdivisions are principal objects of study only in specialized investigations which are still infrequent. Hence, it is hardly worth while to attach to them the exceptionally popular term "facies" narrowing, thereby, its significance.

Analysis of the factual material gathered in the process of numerous exploratory and prospecting operations conducted in alluvial gold placers makes it possible to maintain that the three-member system of alluvial facies structure, as developed by Ye. V. Shantser, requires certain amplifications. The necessity for such amplification becomes evident upon consideration of the structural characteristics of alluvium as related to various dynamic phases.

The Basic Characteristics of the Dynamic Phases of River Valleys and Alluvial Deposits

It is well known that fluvial alluvial deposits in various stages of development differ very substantially from each other. This was first pointed out by V. V. Lamakin [7] who has divided

¹ Fatsii, Dinamicheskiye fazy i svity allyuviya.
77 - 90.

alluvial deposits into three dynamic phases: "instrative", "perstrative", and "constrative". Regrettably, this structural scheme for alluvium is not as popular as that proposed by Ye. V. Shantser. Yet, it reflects the essential structural characteristics of alluvial layers, without a correct understanding of which it is almost impossible to speak about classification of alluvium into facies. The concept of the dynamic phases of alluvium is particularly important for studies of the formation of alluvial placers. However, V. V. Lomakin's position calls for more specific definitions.

The state of dynamic equilibrium in river valleys, according to V. V. Lomakin's interpretation [8], represents only the transition from downcutting to aggradation, while perstrative alluvium which corresponds to this state represents the transition from instrative to constrative alluvium. In reality both the corradng and the alluvium-accumulating rivers strive to achieve, and do achieve, a state of dynamic equilibrium which is maintained when the amounts of loose material supplied to the stream and carried away by it are well balanced. The state of dynamic equilibrium is, perhaps, more commonly observed than the processes of corrasion and aggradation even in rivers of mountain regions. This interpretation of the conditions governing the geomorphological equilibrium of river valleys may not be universally accepted yet, but it is sufficiently well substantiated in a number of relatively recent works of our own and foreign authors [3, 5, 6, 10, 14, 15].

V. V. Lamakin believes that the "floor of alluvium in a perstrative section is approximately at the level of the river bottom" [8, p. 169], and that "here all of the alluvial mass belonging to a given cycle of the river's activity remains within the sphere of its immediate influence" (Ibid, p. 167). The same opinion is shared by Ye. V. Shantser [12], S. A. Yakovlev [13], and many others. But prospecting operations in auriferous regions have long since established the fact that there is always an alluvial layer in the equilibrium sections of river valleys under the bottom of the river and is not affected by the activity of the water current even during maximum freshets. It is precisely to this alluvial layer that the richest of all alluvial placers - the valley placers - are confined. In determining the causes for the formation of this layer, Yu. A. Bilibin wrote: "It is difficult to assume that every mountainous country after a period of uplift causing a deepening of the valleys, could have suffered a subsidence necessary for these valleys to be filled with river sediments" [1, p. 141].

It should be noted also that without the formation of an alluvial layer, preventing the river channel from coming into contact with

the bedrocks, the process of corrasion generally never stops since, of course, there is no equilibrium of any kind between the power of the flow and the resistance of rocks when there is indefinitely continuing action by the stream upon its own bed.

The main reason for the formation of such an alluvial layer during the process of transition from corrasion to the state of dynamic equilibrium, is the widening of the valley and formation of a flood plain. This involves an extension of the channel's length and a decrease of gradient due to the formation of meanders, an increase in the dissection of the interfluvial slopes, a reduction in freshet flow rates, and other phenomena producing a single-valued effect on the balance of the unconsolidated material in the river.

V. V. Lamakin's theories concerning the nature of dynamic equilibrium in river valleys and the mode of perstrative alluvium deposition are, apparently, also related to his views concerning which different phases of alluvial deposits are believed to replace one another as the river passes from one stage of development to another. In fact, however, only the beginning of the corradng process leads to the elimination (sometimes incompletes) of the previously existing alluvium, and to its replacement by instrative alluvium. Both perstrative, and constrative, alluvia, as a rule, do not replace other phases and are deposited on top of them. Consequently, in "normal" alluvial sections of rivers in a state of equilibrium, one can usually distinctly discern the deposits of two phases: the instrative phase deposited directly on the bedrocks (on the subterrane), and the perstrative phase resting on the instrative alluvium (Figures 1 and 3). In valleys containing alluvial deposits of considerable thickness, formations of all three phases generally are observed. In certain cases the alternation of these phases, reflecting the history of the relief development and alluvium formation, may be very complex (Figure 4).

Instrative alluvium. The corrosion process coinciding with the accumulation of instrative alluvium is characterized by a negative balance of loose material. Rivers carry away more loose material than they receive from the corradng sections of river valleys. Therefore the accumulation of alluvium within their limits is only temporary in the majority of cases. Instrative flood plains are usually insignificant in width (Figure 2) and only in exceptional cases are commensurate with the flood plains of sections in equilibrium (Figure 1). All of them have thin layers of alluvium which are completely exposed to the effects of water action, are very rapidly transformed into low terraces at intermediate levels (Figure 1), and are subsequently destroyed. Corrosion sections are encountered where there are no




FIGURE 1. Intrastive flood plain of the Ten'ka River (right tributary of the Kolyma). Section of consolidated deposits along survey line 386, prospect holes 27-78.

1 - soil, 2 - peat, 3 - sandy-silt peat, 4 - loam and argillaceous sand, 5 - silt, 6 - sand, 7 - silty sand; 8 - pebbly-argillaceous deposits (loam or argillaceous sand containing pebbles and gravel); 9 - pebbly silt deposits (silt and silty sand with pebbles and gravel), 10 - pebbly arenaceous deposits (sand with pebbles and gravel), 11 - deposits of detritus and clay (clay, loam, or argillaceous sand with detritus and gruss), 12 - pebbly-detrital-muddy deposits (silt or silty sand with pebbles, detritus, gravel, and gruss), 13 - pebble (pebbles and gravel with silt, sand with detritus and gruss), 14 - pebbly-argillaceous deposits (pebbles and gravel with loam or clayey sand), 15 - boulder beds (boulders, pebbles and gravel with any kind of fine material), 16 - detrital deposits (detritus, and gruss with any kind of fine material), 17 - pebbly-detrital deposits (pebbles, detritus, gravel and gruss with any kind of fine material), 18 - deposits of boulders and detritus (boulders, pebbles, detritus, gravel, and gruss with any kind of fine material), 19 - icing deposits, 20 - bedrocks (sandy-clay shales of the Verkhoyansk complex), 21 - boundaries between formations, facies, and genetic types of deposits, 22 - the same boundaries but lithologically not expressed, 23 - boundaries between the lithological varieties of deposits belonging to the same facies, 24 - boundaries of zones of high concentration placer-gold. Figures over the surfaces of sections are the numbers of prospecting holes.




FIGURE 2. Instrative flood plain of the Bol'shoy Inyakan channel (right tributary of the Kolyma). Section of loose deposits along survey line 180, holes 75-103.

Symbols are the same as in Figure 1.

flood plains. Alluvium here is in a state of constant movement.

The feeble development of flood plains has the result that bottom-land and oxbow facies play practically no role in the formation of instrative alluvium. Their occurrence is rare and they are confined mainly to abnormally wide instrative flood plains (Figure 1). The instrative variety of bed facies is characterized by a number of specific features resulting from its formation during the downcutting process. In the first place, one should point out the presence of a considerable number of angular fragments, not rounded by the river, which get into instrative alluvium by rolling down the slopes that often abut on the channel, and from the bedrock bottom eroded by the river.

The second characteristic is the considerable coarseness of the material making up the instrative variety of the bed facies. In the bottom part of the same sections, instrative alluvium contains, as a rule, coarser pebbles than the overlying deposits of the bed facies of perstrative and constrative alluvium.

At the same time, instrative alluvium is distinguished by poor sorting, and clay particles may often be found in it along with coarse fragmentary material. Clayiness — "unelutriatedness" — is the third peculiarity of instrative bed-alluvium. Ye.V. Shantser [12] describes this phenomenon as the combining of bed and flood-plain alluvia into a single layer, and believes this to be a characteristic feature of mountain alluvium in general. It

should be mentioned here, that the other features, which according to Ye.V. Shantser are peculiar to mountain alluvium — lack of regular succession of facies and insignificant thickness — are characteristic precisely of the instrative alluvium of mountain rivers, and are not so very typical of the perstrative alluvium of these rivers.

The fourth peculiarity of instrative bed alluvium is the largest coarse accumulations of heavy minerals producing commercial placer deposits are localized in its lower horizons. These accumulations never are as large in perstrative and constrative alluvial deposits. This phenomenon is attributable to the low mobility of placer-forming minerals. Every corrosive action of the river leads to the reworking and elutriation of the previously formed alluvial deposits. The heavy minerals contributed to the alluvium by the slope deposits are not carried away during corrasion, and they concentrate in the instrative alluvium, gradually forming very considerable accumulations. Of course, this latter characteristic of instrative alluvium may be manifested only in the river valleys, whose slopes furnish a sufficient quantity of minerals to form placer deposits.

In those cases when the amplitude of corrasion is inferior to the thickness of the previously accumulated alluvium, the instrative deposits are found to be resting not on bed-rocks but on deposits of other phases. It goes without saying that such lithologic characteristics as high proportion of unrounded fragments, the generally greater coarseness of the




FIGURE 3. Perstrative flood plain of Budenny brook (left tributary of the Ten'ka). Section of unconsolidated deposits along line 68, holes 3-33.

Symbols are the same as in Figure 1.

material and its poor sorting, cannot be manifested in instrative alluvium produced by rewashing of deposits no longer possessing these characteristics. Accumulations of concentrated minerals constituting commercial placers may frequently be encountered in such kind of instrative alluvium. These are the, so called, placers with "a false bottom" (Figure 4). However, they are usually not as rich as those localized in instrative alluvium overlying true bedrock [1].

Perstrative alluvium is distinguishable from instrative deposits, and quite often also from the constrative variety, by the more rounded coarse material, total, or almost total, absence of angular fragments, and a high degree of sorting.

In comparing the structural peculiarities of perstrative alluvium in the rivers of gold mining mountains regions with the structural peculiarities of the plains river alluvium described in detail by Ye.V. Shantser [12], one cannot fail to note that there are no fundamental differences between these types of alluvium.

The facies structure of the fluvial perstrative alluvium of mountainous regions is distinguishable only through the somewhat more clearly pronounced prevalence of the bed facies over the flood-plain and oxbow facies. The latter often are completely absent in its sections, but, as may be seen from the accompanying illustrations, this absence is not at all mandatory or typical.

A very considerable contents of coarse material (gravel, pebbles, boulders) in the perstrative alluvium of mountain rivers, as compared with the alluvium of plains rivers, is a specific feature characterizing the lithologic peculiarities of the former. In the deposits of the bed facies this coarse material is, as a rule, predominant; in the oxbow-

facies deposits it may also be encountered rather frequently, although always in lesser amounts (Figure 3 and 4). The flood-plain facies contain an insignificant amount of coarse material only in rare cases, but these materials are more often represented by sandy rather than clay material.

Often, considerable concentrations of heavy minerals are accumulated in the bottom sections of bed facies deposits of perstrative alluvium. Sometimes they occur as placers of commercial significance, but they almost never attain the proportions of placers located at the bottom of instrative alluvial horizons.

The perstrative alluvium of mountain rivers is always entirely within the sphere of action of rivers in a state of dynamic equilibrium. It is precisely its "normal" thickness which can be determined as "the difference between the levels of an average flood and the bottom of average-depth reaches" [12, p. 213]. But, as stated earlier, perstrative alluvium does not lie directly on the bedrock. Consequently, the total "normal" thickness of alluvium in rivers in a state of equilibrium, at least in mountain regions, always exceeds this value.

The peculiarities of the facies structure and the lithologic characteristics of perstrative alluvium are due to the fact that it is formed by rewashing and reworking of the alluvium belonging to other phases by a river occupying an approximately identical vertical position throughout the entire stage of dynamic equilibrium. When this equilibrium is established after the process of alluvium accumulation, the deposition of perstrative alluvium upon the constrative phase requires no special explanation. When equilibrium ensues as a result of downcutting by the rivers the formation of perstrative alluvium is, evidently, preceded by a sharp increase in the thickness of instrative alluvium caused by the phenomena which accompany the transition from corrasion

om the valley. The subordinate, third-rate role played by the flood-plain facies in the structure of the majority of alluvial formations becomes, therefore, comprehensible, regardless of the tectonic conditions of their position".

This position agrees very well with the prospecting data on the structure of constrictive alluvium in auriferous regions. In the sections of constrictive alluvium in auriferous regions, the sections of constrictive alluvium, deposits of flood-plain and oxbow facies are usually found as isolated lenses. They generally occupy a subordinate position and do not differ in their lithological characteristics from the flood-plain and oxbow facies of perstrative alluvium.

The constrictive variety of bed facies predominating in these strata is formed under conditions of aggradation which hinder repeated reworking. Consequently, in terms of its lithologic characteristics, it more closely approximates the instrative than the perstrative alluvium. Thus, poor sorting is quite typical of it — the clayiness of the component pebble beds. Frequently this variety is distinguishable by the noticeable presence of unconsolidated detritus. However, when unconsolidated material is supplied to the build-up area from an up-river section and not from its slopes, this lithologic characteristic may be entirely absent. Being bedded in perstrative alluvium, constrictive bed alluvium usually is conspicuous by its greater coarseness of material. But this distinction may be diametrically opposite. Whenever accumulation is caused by an increased supply of unconsolidated material to the river, the constrictive alluvium is generally coarser than that of the underlying perstrative bed. Whenever aggradation begins as a result of diminishing force of the current, the coarseness of the material comprising the constrictive bed alluvium decreases in comparison with the material of perstrative alluvium. In certain cases the constrictive bed alluvium in mountain rivers is altogether devoid of coarse material and approximates the flat-land type of bed alluvium in respect to lithologic characteristics.

The heavy concentrate minerals usually are dispersed throughout the entire layer of constrictive bed alluvium and very rarely form insignificant concentrations which, as a rule, have no commercial importance.

Subterranean and Cover Facies of Alluvium

Even a first acquaintance with the materials resulting from prospecting operations carried out in gold placers shows that there are no fundamental distinctions between the alluvia of mountain rivers and those of plains regions. Those difference that actually exist consist

mainly in that all mountain river facies are represented by coarser clastic material.

At the same time the differences between instrative, perstrative, and constrictive alluvium — even when they are relatively poorly reflected in lithological characteristics — are very important for a correct understanding of the history of relief formation, alluvial deposits, and placer deposits of commercial minerals. I, therefore, cannot agree with Ye.V. Shantser's assertion that his proposed "normal classification of alluvium . . . is valid for all three of the dynamic phases proposed by V.V. Lamakin". [12, p. 37].

Particularly noteworthy and important are the genetic and lithologic differences between the instrative, perstrative, and constrictive varieties of bed alluvium, which obviously constitutes the principal facies prevalent in most alluvial strata. The essential nature of these distinctions induces me to advance a suggestion: to consider these three varieties as independent facies of alluvium, since they not only fit perfectly the definitions of facies of unconsolidated deposits as described above, but are also taxonomically analogous to the previously identified alluvial facies.

For instrative alluvium which is present in the thalweg² portion of any river valley, and which rests on bedrock (known in placer geology as "subterranean"), I suggest the term "subterranean facies" [5, 6]. For the constrictive variety of bed alluvium the most appropriate designation, I feel, is "cover facies", which reflects its position in the sections of thick alluvial strata.

The term "bed facies" should be retained to designate perstrative alluvium, the study of which is the basis for Ye.V. Shantser's classification. It is true that this term is not in the same category as those referred to above, but it has become so popular that it is hardly possible to change it at the present time.

The subdivision of alluvium into dynamic phases might be reflected to advantage in the genetic indices applicable in geologic sections and used to designate sequences in this paper.

²The term "Thalweg" has been used for a long time in prospecting practice to designate the line connecting not "the deepest parts of the channel" [4], but the lowest sections along the bottom of a valley. This change in terminology is certainly advantageous, since it gives a definite name to a very important element in the structure of river valleys and eliminates the superfluous synonym for the term "farvater" (German "fahrwasser"; —channel in English).

Alluvial Deposits

The basic subdivisions used in local stratigraphic scales is the formation: "a complex of rocks distinguishable by definite characteristics, formed within the limits of a given region under definite conditions, and occupying a definite stratigraphic position in the given region" [4]. Formations in unconsolidated deposits should also be identified according to this definition. The natural age boundaries in such formations are determinable by the changes occurring during the process of formation of the unconsolidated deposits caused by corresponding modifications in the orientation or rate of development of the relief-forming processes.

Formations of unconsolidated deposits may include formations of different origin. However, alluvium is of such a genetic type that its study usually is of decisive importance in developing stratigraphic maps of unconsolidated deposits. Rivers are known to be relief- and rock-formation agents, and they are very sensitive to all changes in external conditions. Because of this, the structural peculiarities of alluvial deposits reflect all details of the history of geologic development.

The most essential modifications of external conditions lead to changes in the relief-forming processes in rivers. Each such change governs the formation of alluvium belonging to a definite dynamic phase. Thus, the boundaries between the dynamic phases of alluvium must be considered as the natural stratigraphic boundaries of alluvial formations. When alluvial deposits are confined to different terrace levels, these stratigraphic boundaries become particularly clearly manifested.

It should be noted that the reaction of fluvial relief-forming processes to changing external conditions in a number of cases does not facilitate, but rather, complicates considerably, the task of the stratigraphic differentiation of unconsolidated deposits. The rapid change of relief-forming processes, in terms of time and in space, causes the sections of alluvial deposits, located even a short distance apart, to display sometimes a rather dissimilar structure. This, of course, hampers the correlation of sequences, identification of formations, and other stratigraphic units. In most sections of alluvial deposits it is possible to identify several stratigraphic units. But, in order to produce a local stratigraphic scale and distinguish formations, comparison and study of an entire series of such sections relative to a more or less extensive area are required. The stratigraphic units in individual sections may be referred to as formations as tentative identification only, and should be designated by index numbers as was done in the case of the sections included in this paper.

It is well known that formations of bedrock must necessarily be named in accordance with their proper geographic location [11]. Similar geographic names should be assigned to formations of unconsolidated deposits identified on the basis of comparison and correlation of sections, as was recommended by Yu. A. Bilibidze [2].

Of course, the formations of unconsolidated deposits belonging mostly to the Quaternary system are thinner and cover less area than do bedrock formations, just as the Quaternary is less widespread than the other systems. This, nevertheless, should not affect the principles of identification and study of these stratigraphic units.

Formations of unconsolidated deposits, like bedrock formations, can be combined into larger subdivisions, such as series. Their boundaries should coincide with the more substantial changes in the character of sedimentation associated with sharp climatic changes, alterations of the general conditions of tectonic movements, etc.

The creation of well-substantiated, local stratigraphic scales is mandatory before determinations can be made of the position of any of the deposit in the general stratigraphic scheme. In studies of unconsolidated deposits this rule is, regrettably, often disregarded. Considerations relative to the amount and sequence of glaciations, the correspondence of fluvial terrace elevations to specific age boundaries, the coincidence of changes in the character of erosion processes with definite geologic data, etc., dominate in such cases over the factual material. These are the factors which become the main source of errors and differences affecting the stratigraphic maps of unconsolidated deposits.

The lithologic characteristics of alluvial and other unconsolidated deposits belonging to varying formations, their thickness, typical position in sections, and other characteristic features, make it possible sometimes to reconstruct, in detail, the history of sedimentation and relief formation. The brief historical reviews of the formation of unconsolidated deposits in several fluvial valleys described in the following section may serve as an example of such an analysis.

The History of Formation of the Relief and Loose Deposits in a Few River Valleys in one of the Regions of Northwestern USSR

Analysis of the structural characteristics of unconsolidated deposits represented in the preceding sections (Figure 1-4), enables us to compile short historical descriptions of their formation which supplement and clarify these

sections. It should be noted that no comparison of these sections with each other, and with other sections, has been made, and that the "formations" in them are identified conditionally. It may well be that a number of these formations do not appear in these sections, and that some of the identified "formations" may, on the other hand, not be characteristic of the region as a whole.

Ten'ka River. The oldest deposits in the section (Figure 1) are the pebble beds of subterranean facies (ial₁) and the subjacent detrital eluvium (e₁) of "formation" 1, at the base of the terrace on the right.

During the stage of dynamic equilibrium which has replaced the degrading process, formed pebble bed facies (pal₂), pebbly argillaceous deposits belonging to the oxbow facies (pal₂), and the flood-plain facies silty sand (pal₂) were formed — all of them constituting "formation" 2. The over-all thickness of the Ten'ka River flood plain is 10 m, which corresponds to its "normal" thickness of alluvium in the equilibrium section starting a few kilometers above the given section.

At present, the downcutting which replaced the equilibrium stage has deepened the valley of the Ten'ka River by some 10-12 meters, and, judging by the structure of the present-day flood plain, has not yet stopped. Characteristic of the corradng process in the Ten'ka valley is the very important role played by horizontal shifting of the channel. As a result, the instrative flood plain is quite wide and has a flat, step-like surface. "Formation" 3, of which it is composed, is represented not only by detrital eluvium (e₃), boulder deposits and pebble beds of subterranean facies (ial₃), but also by sand, silty sand, pebbly-arenaceous and pebbly-silty deposits of the oxbow and flood-plain facies (ial₃). The deposits of flood-plain and oxbow facies are partially replaced by biogenic-alluvial and biogenic deposits: sandy silt peat (bal₃) and peat (b₃). The deposits comprising the lower, near-channel stage of the flood-plain were, apparently, formed somewhat later than the deposits of the upper stage. However, since the downcutting process continued, and there were no changes in the conditions affecting the formation of the deposits, there is no justification for "formation" 3 to be divided into two independent "formations". "Formation" 3 also includes the loam, argillaceous sand, and pebbly-clay deposits of the alluvial-deluvial facies of the terraced spurs (ald₃), and originated as a result of the shifting of the alluvial deposits from the right terrace produced by the processes of slope denudation. The contact of these alluvial-deluvial deposits with the flood-plain alluvium represents, it seems, a pattern of complex interstratification. In the original records it was recorded as con-

sisting of pebbly, argillaceous and pebbly silt deposits, in the section it is considered as undifferentiated deluvial-alluvial deposits (dal₃). The soil (b₃) is also included in "formation" 3.

Bol'shoi Inyakan Creek. "Formation" 1, the oldest in the section (Figure 2), is represented by pebble beds of the subterranean facies (ial₁) and pebbly-detrital eluvial-alluvial deposits (eal₁) forming the base of the right terrace.

"Formation" 1 was, in all probability, covered by perstrative alluvium which later was totally eroded in the course of subsequent relief development. The first development stage (after the formation of "formation" 1) fixed in this section was the cutting of the Bol'shoi Inyakan creek to a depth of not less than 25 m, which cut a canyon in the left part of the section. The instrative alluvium formed during the cutting of the canyon was not exposed by prospect holes. However, the deluvial material which covers the deposits of "formation" 1 on the right terrace, and which is represented by silt and detrital-silty deposits, must have been, apparently, partially synchronous with this instrative alluvium. The formation of deluvial deposits, evidently, also must have continued later — during the formation of "formations" 3 and 4, as this reflected in its index (d₂₋₄).

The next relief development stage observable in the section was the accumulation of alluvium and partial burial of the canyon. "Formation" 3 which was formed at this time is represented by the clayey pebble deposits of the cover facies (cal₃) and the eluvial-alluvial, pebbly-detrital deposits (eal₃).

The aggradation process terminated when the surface of the Bol'shoi Inyakan floodplain attained the elevation of the present left terrace. The ensuing state of dynamic equilibrium was accompanied by the process of fluvial abrasion — the Bol'shoi Inyakan was scouring its right bank, widening the flood plain, and destroying the right terrace. This period saw the formation of "formation" 4, represented by pebble bed facies (pal₄), silt of the flood-plain facies (pal₄), eluvial-alluvial deposits of pebbly detritus (eal₄), and detrital eluvium (e₄). Since fluvial abrasion continued after the establishment of dynamic equilibrium, the deposits of the bed facies in the central part of the section are separated from the bedrock by only a thin layer of eluvial-alluvial or eluvial deposits.

Dynamic equilibrium was again replaced by another period of erosion during which the Bol'shoi Inyakan valley was deepened for 5-6 m. This still continues at the present time. "Formation" 5, which is presently being formed, is represented by pebble beds of the subterranean facies (ial₅), oxbow silt (ial₅), eluvial-alluvial deposits of pebbly detritus (eal₅), and soil (b₅).

Budennogo Creek. Section (Figure 3) reflects only the most recent formative stages of the Budennogo Creek valley and the unconsolidated deposits filling it. "Formation" 1 is represented by detrital eluvium (e₁), pebbly clay deposits of subterranean facies (ial₂), and the beds of pebbles and detritus belonging to the terraced-spur facies (ald₁). The presence of these spurs betrays the fact that there used to be a terrace on the right slope, which consisted of alluvial deposits. The destruction of this terrace by processes of slope denudation has actually lead to the formation of deposits of the terrace-spur facies. The formation of "formation" 1 deposits continued not only throughout the process of degradation, but also during fluvial abrasion (widening of the valley) which served as the transition stage between downcutting and dynamic equilibrium.

"Formation" 2, represented by pebble beds of the bed facies (pal₂), pebbly silt deposits of the flood-plain (possibly, oxbow) facies (pal₂), and the soil (b₂), — was formed during the dynamic equilibrium stage, continuing to the present time. The deposits belonging to the terraced-spur facies were accumulated as a result of continuing destruction of the right terrace, while the deposits of alluvial facies are the product of protracted reworking of the deposits belonging to the subterranean facies and partly to that of the terraced spurs.

Pavlik Creek. Judging by the fact that the gold-bearing layer of the right terrace (Figure 4) extends along the slope of the buried canyon and, apparently, constitutes an older formation than this canyon, the deposits resting on the terrace at the bottom of the unconsolidated stratum are the oldest formations in the section. "Formation" 1 is represented by eluvial-alluvial deposits of pebbles and detritus (eal₁) and argillaceous-pebbly deposits of the subterranean facies (ial₁).

The great width of the right terrace indicates that a perstrative flood plain must have existed at its level. This justifies the assignment of the pebble beds and the argillaceous-pebbly deposits overlying "formation" 1 to the bed facies (pal₂) of "formation" 2. We can not exclude the possibility, however, that the deposits related to "formations" 1 and 2 were actually formed after the burial of the canyon.

During the process of the corrasion which was responsible for the canyon carving, there the following components of "formation" 3 were laid down: the deposits of boulders and detritus and those of pebbles and detritus belonging to the subterranean facies (ial₃), the pebbly-detrital eluvial-alluvial (eal₃) and the deluvial-alluvial deposits (dal₃).

Degradation gave way not to dynamic equilib-

rium but to accumulation of alluvium which caused the burial of the canyon under the argillaceous-pebbly, pebbly-detrital, and pebbly deposits of the capping facies (cal₄), the pebbly clay deposits of oxbow facies (cal₄), the pebbly-detrital talus-alluvial (dal₄), detrital and detrital-argillaceous talus (d₄) deposits of "formation" 4. The glaciated portion of the argillaceous-pebbly deposits is, in all probability, syngenetic.

Dynamic equilibrium, it seems, must have begun after both the canyon and the right terrace were buried under the unconsolidated deposits. At this time the pebble beds belonging to the channel facies (pal₅) and the pebbly clay and pebbly sand deposits of oxbow facies (pal₅) comprising "formation" 5, were deposited.

Dynamic equilibrium was again replaced by down cutting resulting in the formation of argillaceous-pebbly deposits of the subterranean facies (ial₆) of "formation" 6, distinguishable from the subjacent deposits of the capping facies of "formation" 4 not only by the absence of glaciation, but also by the much smaller amount of cementation. The boundary between "formations" 6 and 1 was traced tentatively, since the original records show no differences between them. It should be stated here that "formation" 1 differs from "formation" 6 by the absence of eluvial-alluvial deposits and the great thickness of the gold-bearing layer "inserted" much deeper into the fissures of the bed rock.

During the subsequent dynamic equilibrium stage, the pebble beds of the bed facies (pal₇), the pebbly-sand deposits of the oxbow facies (pal₇), and the glaciated silty deluvial deposits (d₇) of "formation" 7, were deposited.

The most recent low-amplitude downcutting may be judged mainly on the basis of the relief character, since the pebble beds of the subterranean facies (ial₈) of "formation" 8 are practically undistinguishable from the bed alluvium of "formation" 7. The silty and pebbly clay deposits of the terraced-spur facies (ald₈) and the soil (b₃) were formed simultaneously with the pebble beds of "formation" 8.

CONCLUSIONS

The subdivision of alluvium into dynamic phases, as proposed by V. V. Lamakin [7], is of great importance for a correct understanding of the principal structural characteristics of alluvial deposits.

The destruction of the pre-existent alluvium belonging to different phases occurs only with the formation of instrative alluvium. Perstrative and constrative alluvia in the process of their formation do not replace the previously existing alluvial phases, but cover them up.

The definition of "normal" thickness of alluvium as a difference between the elevations of average freshets and the bottom of average-depth reaches [12] is correct only with respect to perstrative alluvium. In view of the fact that in equilibrium rivers, perstrative alluvium is superposed on instrative deposits, the total "normal" thickness of these deposits, at least for the case of mountain rivers, is always greater than that calculated by Ye. V. Shantser's method [12].

There are no fundamental differences in the facies structures of plains and mountain river alluvia. Mountain-river alluvium is distinguishable only by the greater coarseness of the component material and somewhat less developed oxbow and flood-plain facies.

On the other hand, the distinctions in facies structure and lithologic characteristics of alluvium belonging to different phases are very considerable. They are particularly noticeable in the alluvium called by Ye. V. Shantser [12] bed alluvium. This makes it possible to identify in the bed alluvium characterized by Ye. V. Shantser, three facies: subterranean - in the instrative phase, bed - in the perstrative phase, and capping - in the constrative phase.

The presence in alluvial sections of deposits belonging to deposits belonging to different dynamic phases is determined by the changes in the character of fluvial relief-forming processes and, consequently, reflects the definite historical stages of geologic development. Thus, these deposits of various phases may also be considered as stratigraphic units - formations.

Assimilation and theoretical generalization of the vast material accumulated in the course of geologic prospecting operations in placer deposits will, no doubt, lead to very considerable contributions to the knowledge of the regularities governing the formation of alluvium.

REFERENCES

1. Bilibin, Yu. A., *Osnovy geologii rossyey*. [PRINCIPLES OF THE GEOLOGY OF PLACERS]: GONTI, 1938.
2. Bilibin, Yu. A., *Geomorfologicheskiye issledovaniya kak osnova poiskov i razvedok slozhnykh tipov rossyey*. [GEOMORPHOLOGIC RESEARCH AS A BASIS FOR PROSPECTING AND EXPLORATION OF COMPLEX TYPES OF PLACERS]: Izbr. trudy, t. 2. Izd-vo Akad. Nauk SSSR, 1959.
3. Gavrilov, M. D., *Nekotoryye osobennosti razvitiya gornykh rechenykh dolin*.

[CERTAIN PECULIARITIES OF MOUNTAIN RIVER VALLEY DEVELOPMENT]: Geogr. sborn., t. 1. Izd-vo Akad. Nauk SSSR, 1952.

4. *Geologicheskii slovar'*. [GEOLOGICAL DICTIONARY]: Gosgeoltekhizdat, 1955.
5. Kartashov, I. P., *Flyuvial'nyye rel'yefoobrazuyushchiye protsessy*. [FLUVIAL RELIEF-FORMING PROCESSES]: Tr. Vses. n.-i. in-ta zolota i redkikh metallov. Geologiya, vyp. 29, 1957.
6. Kartashov, I. P., *O plotikovoy fatsii allyuviya*. [THE SUBTERRANEAN FACIES OF ALLUVIUM]: Kolyma, No. 1, 1958.
7. Lamakin, V. V., *O dinamicheskikh osobennostyakh allyuvial'nykh otlozheniy*. [THE DYNAMIC CHARACTERISTICS OF ALLUVIAL DEPOSITS]: Doklady, Akad. Nauk SSSR, t. 57, No. 1, 1947.
8. Lamakin, V. V., *Dinamicheskkiye fazy rechnykh dolin i allyuvial'nykh otlozheniy*. [THE DYNAMIC PHASES OF RIVER VALLEYS AND ALLUVIAL DEPOSITS]: Zemlevedeniye, t. 2(42), 1948.
9. Markevich, V. P., *Ponyatiye "fatsiya"*. [THE CONCEPT OF "FACIES"]]: Izd-vo Akad. Nauk SSSR, 1957.
10. Piotrovskiy, M. V., *K teorii flyuvial'no-denudatsionnogo (erozionnogo) tsikla*. [THE THEORY OF THE FLUVIAL DENUDATION (EROSIVE) CYCLE]: Tr. In-ta geogr. Akad. Nauk SSSR, vyp. 39, 1948.
11. *Stratigraficheskaya klassifikatsiya i terminologiya*. [STRATIGRAPHIC CLASSIFICATION AND TERMINOLOGY]: Gosgeoltekhizdat, 1956.
12. Shantser, Ye. V., *Allyuviiy ravninnykh rek umerennogo poyasa i yego znachenie dlya poznaniya zakonomernostey stroyeniya i formirovaniya allyuvial'nykh svit*. [THE FLATLAND FLUVIAL ALLUVIA OF THE TEMPERATE ZONE AND ITS SIGNIFICANCE FOR THE PERCEPTION OF THE STRUCTURAL AND FORMATIVE REGULARITIES OF ALLUVIAL SERIES]: Tr. In-ta geol. nauk Akad. Nauk SSSR, vyp. 135, geol. ser. (No. 55), 1951.
13. Yakovlev, S. A., et al., *Metodicheskoye rukovodstvo po izucheniyu i geologicheskoy s'yemke chetvertichnykh otlozheniy*, ch. 1. [METHODICAL MANUAL FOR THE INVESTIGATION AND GEOLOGICAL MAPPING OF QUATERNARY DEPOSITS, PART 1]: Gosgeoltekhizdat, 1954.

14. Mackin, H., Concept of a graded river. Bull. Geol. Soc. America, vol. 59, No. 5, 1948.
15. Strahler, A.N., Dynamic basis of geomorphology. Bull. Geol. Soc. America, vol. 63, N. 9, 1952.

The All-Union Scientific-Research Institute
of Gold and Rare Metals,
Magadan

Received May 5, 1960

THE PROBLEM OF THE RELATIONSHIPS BETWEEN MARINE AND CONTINENTAL DEPOSITS IN THE LOWER AND MIDDLE VOLGA REGIONS¹

by

Yu. M. Vasil'yev and P. V. Fedorov

The solution of the problem concerning the relationships between the marine Quaternary deposits of the Caspian Sea and the alluvial formations of the Volga is very important for the general problem of the comparison of the marine and continental deposits in the southeastern part of European USSR.

The Lower and Middle Volga regions, where these relationships must be clarified are well studied. Here the first stratigraphic scale was developed for the Quaternary deposits of the Caspian, and definite ideas were formulated by which the Khvalynsk maximal transgression was compared with the Volga terrace II and with the "Würm" glaciation of the Russian plain, the Khazar transgression - with terrace III: the "Riss" glaciation terrace, and the Baku - with the 4th: "Mindel" glaciation. However, enough factual material was recently accumulated to justify the introduction of certain corrections and amplifications into the existing concepts connected with this matter. The observations conducted by P. V. Fedorov [22] in 1951 along the right bank of the Volga from 'Astrakhan' to Astrakhan' have shown that the highest (48 m) Lower Khvalynsk terrace of the Caspian adjoins not terrace II, but Volga terrace III. M. N. Grishchenko arrived at the same opinion [5] on the basis of a comparison of the Volga and Don terraces.

In 1959 the authors of this paper jointly studied sections of the Quaternary deposits along the lower and middle Volga from Astrakhan' to Kuybyshev. Considered below are mainly the problems of the correlation of the Khvalynian marine terraces with the terraces of the Volga, and also the relationships between certain older alluvial and marine beds. The analysis is based on the data from 1959 observations and earlier materials.

The Baku, Khazar, and Khvalynsk deposits, stratigraphically similar to the corresponding

layers of the entire Caspian basin, were identified on the basis of borehole logs and in natural outcrops in the lower Volga valley. The Novocaspien sediments occur on the southernmost tip of the region along the Caspian coast.

The Baku deposits are represented by thinly-bedded marine clay rocks of dark-grey and brownish colors containing subordinate layers of siltstones and a mollusk fauna characteristic of the entire Caspian basin: *Didacna parvula* Nal., *D. catilus* Eichw., and others. These deposits outcrop in a limited number of places in the Volga valley (Chernyye Yar, Kap Kapustin Yar, and other villages), and were discovered in many borings outside its confines. In the region of Lake Baskunchak the Baku deposits occur in the coastal facies of shingle and coquina beds. At the top of the Baku stage continental formations may be occasionally observed along the Volga.

On rocks of the Baku stage, showing distinct traces of erosion, there are light-yellow, obliquely laminated, alluvial sands grading upward into a layer of grey and dark-grey, thinly-bedded clays with bands of sand which contain an abundance of fresh-water fauna (*Unio Planorbis*, *Sphaerium*, *Corbicula fluminalis* Müll.) and remains of vegetation. In the lower reaches of the Volga, starting at the village of Chernyy Yar, these clays and the subordinate sand layers contain a large variety of Lower Khazar mollusk fauna: *Didacna subpyramidata* Prav., *D. pallasi* Prav., *D. paleotrigonoides* Fed. The alluvial sands at the bottom of this layer, known as the Chernoyarsk sands, contain, according to V. I. Gromov's works [6, 7], the remains of mammals belonging to the Khazar complex: *Elephas trogontherii* Pohl., *Bison priscus* var. *longicornis* V. Grom., *Camelus knoblochi* Nehr., *Elasmotherium sibiricum* Fisch. The facies character of the overlying dark-grey clays permits them to be considered as oxbow-lake deposits which, further south, merge with the freshened Lower Khazar marine sediments. This fact is conclusively evidenced by the afore-mentioned mollusk fauna, similar to that of the Lower

¹ K voprosy o sootnoshenii morskikh i kontinental'nykh otlozheniy nizhnego i srednego povolzh'ya. Izv. AN SSSR, ser. geol. i fiz.-mat. nauch., 1960, no. 1, p. 91 - 99.

Khazar deposits in Dagestan, Azerbaydzhan, and Turkmeniya.

Higher in this stratum one observes a predominance of chocolate-brown clays overlain by a very characteristic horizon of unstratified, lumpy, reddish and reddish-brown loams in which are buried meadow-bog-type soils. These loams are full of crystals and concretions of gypsum. Their tops are covered with frost fissures, the rocks are disturbed by frost heaves. The entire layer represents a single alluvial and alluvial-marine sedimentary complex, which is made up, at the base, of fluvial sands passing upward into oxbow alluvium, which is replaced in the south by the marine and alluvial-marine Lower-Khazar sediments.

Above the sharply scoured top of the Lower Khazar deposits there is a younger alluvial formation at the bottom of which, starting at the village of Nikol'skoye and farther south, are wedged brakish-water, laminated, argillaceous and arenaceous-argillaceous sediments containing a transitional assemblage of mollusk fauna. Included here are shells, still approximating the Khazar stage (*Didacna subovalis* Prav., *D. ex. gr. surachanica* Andrus., *D. ex. gr. crassa* Eichw.), and also the typically Khvalynsk forms, (*Didacna cristata* Bog., *D. ebersini* Fed.), which are clearly prevalent. In addition, one also finds many *Monodacna caspia* Eichw. These deposits belong to the Khazar stage and are identified by P. V. Fedorov [21, 22] as Upper-Khazar. They are underlain by obliquely-laminated sands in the form of lenses and bands.

The top of the Upper Khazar clay deposits is often eroded and are abruptly and unconformably overlain by sands. In a number of places, however, the underlying clays alternate with the covering sands and the former rather gradually grade into the latter. These sediment relationships attest to the estuarine and deltaic conditions of sedimentation. The light, almost white, sands covering, and sometimes underlying, the Upper Khazar brakish-water sediments are characterized by cross bedding and, apparently, constitute the bed facies of the alluvium. Further upward, they gradually become more clayey and grade into sandy and loess-like loams. In tracing this entire formation up-stream along the Volga, one gains the conviction that it represents a single complex, the individual components of which mutually grade into each other or the facies replace one another. Thus, in the region of Stalingrad (Raygorod, Srednyaya Akhtuba, and so on), we deal with a single alluvial formation which begins at the bottom with the bed facies and finishes at the top with sandy-loamy, loess-like formations known in the literature as "Atelian loams".

The Atelian formation was identified by P. A. Pravoslavlev [19] as an independent stratigraphic unit, in which he included not only the loess-like loams and sandy loams, but also the sub-jacent, obliquely laminated alluvial sands associated with the former by gradational contacts. In subscribing to P. A. Pravoslavlev's opinion, we feel that the isolation in the Volga Quaternary section of the Atelian continental formation, presented by alluvial sands and loess-like sandy loams and loams of different origin, but everywhere covered by Lower Khvalynsk marine layers, is wholly justified and expedient. Considering the significant erosion separating this suite (including the mentioned brakish-water sediments) from the top of the Lower Khazar deposits, the stratigraphic unity of the entire Atelian formation, and the absence of a major hiatus between it and the incumbent Lower Khvalynsk marine sediments, it is possible to speak of the greater stratigraphic similarity of the "Upper Khazar" marine layers in the Lower Volga regions to the Khvalynsk stage, than to the Lower Khazar stratum. It would probably be correct to compare these sediments to the "Ghirkanian beds" of G. I. Goretskiy [3] or to the "Lower Khvalynsk beds" of G. I. Popov [18] underlying the Gudilovian ("Bourtassian") sediments and overlying the Karangatian marine beds in the Manych valley.

On the basis of the data from the Lower Volga section it is possible to surmise that the "Upper Khazar" marine beds and their alluvial analogs are associated with the transgression of the Caspian which immediately preceded the maximum Lower Khvalynsk transgression and was separated by a deep regression from the Lower Khazar transgression. Nowhere in the section of this section are there any significant discontinuities. Everywhere the cross-bedded sands grade upward into dusty sands, sandy loams, and loess-like loams.

Thus, in setting up the problem, one may say that the brakish-water sediments in the lower Volga region, usually referred to as Upper Khazar, should no longer be assigned to the Khazar stage, but to a younger formation which should retain its original designation, suggested by P. A. Pravoslavlev, the Atelian formation. The brakish-water sediments are the lowermost section of this formation.

In the lower Volga region, and throughout the entire Caspian lowland, the Atelian formation is overlain by marine Khvalynsk deposits represented by sands and peculiar "chocolate" clays characterized by banded lamination. The Khvalynsk stage deposits in the described territory, as well as in other parts of the Caspian basin, can be subdivided into smaller stratigraphic units. P. V. Fedorov [21, 22] distinguishes here the Lower Khvalynsk deposits which, in the Lower Volga regions, form terraces having absolute elevations of 48, 35, 22-25 m (the first

are composed of loams and sands, the last of (chocolate-colored clays), and the Upper Khvalynsk sandy deposits forming terraces having absolute elevations of minus 2, minus 10-12, and minus 16-17 m. Yu.M. Vasilyev considers the chocolate clays which make up the 22-25 meter Volga terrace as an independent Middle Khvalynsk horizon. The Lower Khvalynsk, marine loamy, sandy, and pebbly deposits comprise the 48-meter terrace located on the right bank of the Volga between Stalingrad and Kamyshin. They also cover the watershed terrace and all of the area in the northern part of Caspian lowland.

Thus, along the lower Volga, two alluvial-marine sedimentary complexes are clearly distinguishable: the lower, or the Lower Chazar, deposited over the plane of erosion on the marine beds of the Baku stage, and the upper, or Atelian, which, at its base, consists of brackish-water sediments. It is separated from the lower complex by a pronounced plane of erosion and distinct traces of freezing processes. Deposited above are the marine Khvalynsk formations which are either separated from the Atelian formation by traces of a hiatus, or have a gradational contact with them.

Up-stream along the Volga, the Lower Khvalynsk marine sediments are replaced by alluvial sands covering the terraces III and IV. Consequently, there are grounds to consider these (Lower Khvalynsk) deposits as the third — the uppermost alluvial stage.

The discussed complexes are also clearly well traceable beyond Stalingrad. For example, near Bukovo village (left bank of the Volga, below the city of Kamyshin), on the southern slope of the ravine, the following sequence can be observed, from the top down:

1. Soil and chocolate-colored clay eluvium; 0.8 m.

kv 2. Chocolate-colored clays, thinly-bedded, alternating with bands of yellow sands, followed by sands. The clays are deposited on a lightly eroded surface of the subjacent rocks, and in some locations, grade into them; 3 m.

at. 3. Sands, yellow, laminated and cross-bedded, in some places the superjacent loess-like sandy loams are preserved; clear traces of erosion are observable on the surface of the underlying deposits; 2-3 m.

hz₁ 4. Brownish-grey, greenish, less frequently reddish-brown loams containing traces of freezing processes; about 1-3 m.

North of the Yeruslan River (outside the limits of the Caspian lowlands) we have observed the following sequences in the scarps of the III and IV.

Near the village of Ilovatka, on the lower reaches of the Kamyshhevakha River, the following sequence is exposed in the scarp of the terrace III.

1. Soil.

kv 2. Chocolate-colored clay eluvium containing an abundance of lime nodules; 0.7 m.

at 3. Yellowish-grey sand containing thin streaks of gypsum; ochre spots are visible. The lamination of the sand is not clear; 1.5 m.

at 4. Yellow-brown porous loam containing streaks of gypsum; replaced below by poorly laminated sand containing clastic stains, mole-hills are observable; 1.5 m.

hz₁ 5. Heavy, brownish-fallow loam, darker in the upper sections, becomes mixed with sand toward the bottom, assumes a lighter coloration, and contains many calcareous streaks. Traces of freezing are observable in the top - pseudomorphs of glacial wedges. Toward the bottom the loam is replaced by sand.

hz₁ 6. Brownish-fallow loam, lighter toward the bottom, at the bottom containing yellowish-grey, occasionally light-grey sand; 1.3 m.

khz 7. Brown loam, 0.3 m.

khz 8. Dark-brown to black loam, gleyed, lumpy, unstratified, with numerous lime spots typical for loess which contains lime nodules, with ochre spots which are particularly abundant at the bottom; below, the rock becomes lighter, ochreous and sandy; 1 m.

hz 9. Greyish-brown loam, finely mottled, containing an abundance of ochreous, greenish, and black spots. At the top — a thin sand layer. The loam becomes mixed with sand at the bottom, assumes a brownish-cinnamon tint, becomes more homogeneous and gives way to bed-10 rocks. Mole-hills are visible. The top of the bed is crumpled into small folds; 1 m.

hz 10. Brownish-cinnamon, fine-grained, compacted sand; the bottom shows bands of brown loam; 1 m.

hz₁ 11. Yellowish-grey sand, light-grey at the bottom, with layers of brown loams and clay. Mole-hills filled with brown loam, 2 m.

Evident here is the distinct boundary between beds 2 and 3. Traces of soil formation are observable in the top of beds 4 and 8.

The following sequence appears in the ravine cutting through terrace III between the villages of Privolzh'ye and Skatovka, near the road from Kamyshin to Engel's.

1. Soil.

hv 2. Yellow-brown loam, columnar, toward the bottom, replaced by non-laminated sand; 3.6 m.

at 3. Brownish-cinnamon loam, humic, lumpy, porous, containing calcareous bands and spots, intensely calcitized; at the base of the bed the loam is dove-colored, greenish, compacted, sandy; 0.8 m.

at 4. Brownish-yellow sandy loam, dark with bluish streaks, containing round limy nodules; mole-hills are visible; 1 m.

at 5. Striated-lamellar sand, fine-grained, clayey, more homogeneous at the bottom; contains sypuchka; 2.5 m.

hz₁ 6. Brownish-cinnamon loam, lumpy, with an abundance of calcite streaks in the lower sections; minor frost-induced deformations are visible at the top; 0.3 m.

hz 7. Brown loam, containing pseudomycelles of lime, 0.15 m.

hz 8. Dark-brown loam, particularly dark at the top, bluish, gleyed, lumpy. The lower sections display an abundance of ochre streaks and spots. At the bottom, loam is replaced by sand; 0.8 m.

hz 9. Compacted, ochre-yellow sand, mottled, laminated, deposited in the form of a lens; 0.3 m.

hz₁ 10. Clayey sand, ash-grey, dark, lighter in the lower sections and less clayey; 1 m.

Here, as near Ilovatka, three horizons of deposits (beds 2, 3-5 and 6-10), separated by buried soils, can be discerned.

A good exposure of the terrace IV section (the elevation of the brow above the Volga River is about 35 m) is observable near Spasskoye village.

1. Soil.

hv 2. Yellow-brown, stratified sand, containing bands of brown loam; 3-4 m.

at 3. Brown loam with greenish bands; the top is eroded, uneven, frost-type deformations are observable, although they are minor; mole-hills are visible in the rocks; 0.5 m.

at 4. Brown loam, occasionally cinnamon, changing near the bottom to yellowish-gray, fine-grained sand, with loamy bands in the top section and homogeneous at the bottom; 6-7 m.

hz₁ 5. Greenish-brown loam abounding with lime spots and pockets of fresh-water mollusk shells; a cinnamon loamy bed is in the top of the section. The top of the bed is slightly eroded; 1.5 m.

hz₁ 6. Brownish-cinnamon loam, sandy near the bottom and gradually replaced by compacted sand, yellow-grey, finely grained; 1.5 m.

hz₁ 7. Yellowish-cinnamon sandy loam containing ochre spots. Small pockets of the overlying sand are observable at the top; 0.8 m.

hz 8. Heavy, lumpy, brown loam; sand filled cracks are observable, manifestations of shrinkage; 2 m.

According to Yu. A. Lavrushin [12] an outcrop of the lower part of the section which was buried under talus deposits — alluvial sands of considerable thickness — is visible near Spasskoye village. Here, as at other stations, one can observe three horizons (layers 2, 3-4, and 5-8).

Considering other available data on the structure of terraces III and IV, it is possible to conclude that the sediment sequence is fairly constant within their range; namely, visible is the lower alluvial formation — sands and shingle — overlain by silty and loamy sediments. Meadow-marshy formations are developed at the top of the formation. The top usually shows signs of frost destruction and scouring. The upper alluvial formation consists of sands overlain by loams. Buried soil is developed on its top, with occasionally observable erosion and traces of freezing. Finally, the topmost member of the terrace section is thin (not more than 5 m thick) sand and loam beds covering the terrace.

All of these horizons and their contacts are clearly visible in the overwhelming majority of the terrace sections, and they can, therefore, be traced successively down-stream along the Volga River. Moreover, the lower alluvial formation, in which the entombed remains of Khazar mammals were discovered — is traceable from the village of Spasskoye to the village of Ilovatka, and can be observed near the city of Nikolayevsk, the village of Bukovo, and further south. Near Stalingrad it is not as thick. In the top, in its roof cultivated soil layer, a Mousterian encampment has been found. In the lower Volga this formation is traceable in all exposures and was described at the beginning of this article as the Lower Khazar formation.

It is important to emphasize that along the entire course of the Volga from the lower reaches to the area of Samarskaya Luka, the Lower Khazar alluvial complex retains not only its specific features and the basic succession

of bedding, but also everywhere contains in its base ("Chernoyarian" sands) remnants of a single faunal complex of mammals — the Khazar complex. This is the lower alluvial formation, according to Ye. V. Shantser's interpretation [23], or the lower stage of alluvium, as identified by A. I. Moskvitin [16]. Almost everywhere in the top of this series one observes fossil soil, traces of erosion and cryoturbation.

The upper alluvium formation is also well traceable in the Lower Volga region. The described structure of this formation was traced by us from the villages of Vladimirovka and Yentayevka (lower Volga) up to the region of Syzran' (Spasskoye village), and, this structure is also known in the areas above Samarskaya Luka. In the opinion of A. P. Mazarovich [14] and Ye. N. Shchukina [24], terrace III, and, consequently, also the deposits of the described formation — follows the aprons of the maximum (Riss) glaciation in the Maryinsky district of the Volga region. Thus, the entire formation as a whole represents a periglacial formation.

In the uppermost horizon of the terrace II-IV section (near the mouth of the Yeruslan River and south of it), marine Lower Khvalynsk fauna was discovered.

Taking all these facts into account, the age of the lower alluvial formation may be determined as Lower Khazar; that of the upper — as Atelian, and that of the uppermost mantle of the terrace — as Lower Khvalynsk. It follows, then that the sediments covering the terrace, and, therefore, the terrace itself (III and IV), date back to Early Khvalynsk time. This three-member structure is determined on the basis of numerous borings drilled in terraces II-IV of the Volga.

The age of the terraces also can be established in tracing them further south.

Terrace I, above the flood plain, is well pronounced in the valley of Volga from Spasskoye village to Stalingrad, and below. On the lower reaches of the Volga, the surface of the terrace levels out and it grades into the Khvalynsk plain of the Caspian, which is covered here by Upper Khvalynsk marine deposits. The same sediments are deposited upon terrace I in the lower-course area of the Malaya Uzen' River (Trans-Volga region). A similar phenomenon was discovered in the lower reaches of the Ural River by A. G. Doskach and I. P. Gerasimov [8].

Terrace II is clearly visible in the Volga valley, north of the mouth of the Yeruslan River and within the confines of the Caspian area. In the vicinity of Stalingrad it merges with the 25-m, Lower-Khvalynsk terrace.

In the opinion of Yu. M. Vasil'yev [1], terrace II is of Middle Khvalynsk age.

Volga terrace III can be traced easily from the village of Privolzh'ye to Ilovatka. Here one observes a uniformity of the component sediments. This excludes the supposed possibility of the Khazar terrace III plunging in the south under the deposits of terrace II, or of the merging of the levels of these terraces, etc. Near the marginal bench of the elevated watershed in the vicinity of Ilovatka village, terrace III widens and merges with the 35-meter Early Khvalynsk Caspian plain. That terrace III and the low North Caspian plain from a single geomorphologic surface is evidenced by their almost identical elevation (the absolute elevation of the terrace and the plain is 30-40 m, the elevation above the Volga — about 40 m).

Terrace IV passes in the south into the correspondingly elevated (48 - 50 m), so-called, waterline marginal terrace at the northern extremity of the Caspian. This terrace consists of marine, faunally characterized, Khvalynsk deposits and represents the abrasion platform of the Khvalynsk sea. P. V. Fedorov [22] considers that it reflects the maximum uplift of the basin level. We agree with Yu. A. Lavrushin [12] that Volga terraces III and IV are, stratigraphically, a single structure, since they consist of identical rocks. But geomorphologically they represent independent sedimentation surfaces associated with two development phases of the Lower Khvalynsk sea.

The Early Khvalynsk age of terraces III and IV is proven also by the fact that their level (below Saratov) does not exceed the highest level of the Early Khvalynsk basin. Its sediments, as well as the alluvium of the Volga which emptied into this sea, overlie the Khazar deposits of the said terraces.

Thus, beyond the limits of the Caspian, in addition to the flood plain there exists in the Volga valley the following terraces: I - the (Late Khvalynsk) terrace above the flood plain, II - the terraces consisting of the chocolate-colored clays (Middle-Khvalynsk), III and IV - the Early Khvalynsk terraces. Analogous conclusions were made earlier by P. V. Fedorov [21, 22], M. G. Kipiani, and L. D. Kolbutov [11].

The problem is could there have been terraces older than Khazar and Baku higher than the Early Khvalynsk terrace in the Volga basin?

Whenever this or that investigator identifies Pre-Khvalynsk terraces, their age is always determined only by the position of these surfaces above the Khvalynsk (or Khazar) terrace. B. A. Mozharovsky [15], in explaining the presence of a Mindel terrace, proposed that

the level of the Baku sea rose to a height of 100 m. In the light of present-day data, this theory appears to be totally unsubstantiated. S.V. Lyutsau [13], A.I. Moskvitin [16], and others identify Baku and Khazar terraces higher than Khvalynian terrace II. But we have already shown above that terraces III and IV ("Khazar" and "Baku") are of Lower Khvalynsk origin. In speaking of the existence of the Khazar and Baku terraces at an elevation of 60, and 90-100 m, one should assign to these terraces a large portion of the southern part of the waterline plain whose surface has absolute elevations below 100 m. However, this plain consists of typically waterline clays. Consequently, in this case, too, identification of Pre-Khvalynsk terraces is senseless.

Analysis of the known facts shows that the Khazar and Baku deposits are buried under the Khvalynsk sediments, and that there can be no Khazar or Baku terraces in the Volga valley. In Baku time a rise of the base level, caused by the transgression of the Baku sea (its maximum level was about 0 m absolute elevation, or slightly higher), led to alluvial and lacustrine-alluvial sedimentation in the Volga valley. In the Khazar age, when the sea level rose to 10-15 m absolute elevation, the Khazar deposits in the Volga valley covered the Baku beds (filling up the deep cuts). It is natural that in Early Khvalynsk time, when the sea level was elevated to 48 m absolute elevation, the Khazar beds were buried under the Khvalynsk deposits. In this case, the Khazar and Baku terraces could not have been higher than the Khvalynsk.

However, the presence of the rising steps of Pre-Khvalynsk terraces may be admitted, if one is to assume that there was an uplift of the watershed area (including the Volga regions beyond the Caspian) and a subsidence in the Caspian area. Then, the Khazar and Baku Volga terraces observable in the relief should exhibit a pitch towards the south and should plunge under the level of the Khvalynsk terrace at the boundary with the Caspian lowland. Incidentally, such a subsidence of the Ural River valley is reported by A.G. Doskach [8]. A similar phenomenon was also noted in the Volga valley. In this case, in tracing the horizons of the Pre-Khvalynsk deposits from the lower reaches of the Volga to the mouth of the Yeruslan River and further north, we should have noticed a rise of the Baku, Lower Khazar and Atelian beds (let it be reminded that they are distinctly observable in the outcrops) and their truncation by the Khvalynsk deposits of terraces III and IV. But on the left bank of the Volga such facts were not noted (and this also applies to the Ural). Here, one observes only a gentle rise of all beds, corresponding to the elevation of the Khvalynsk terrace level. Within the limits of this terrace, both in the Caspian and north of it (near the villages of Ilovatka, Skatovka, and Spasskoye), a series

of successive Khazar beds is observed under the sediments deposited during the maximum transgression of the Khvalynsk sea. G.I. Goretskiy [4] describes precisely such a mode of bedding of the Khazar alluvial formations, which rise up-stream along the Volga and Kama Rivers parallel to the water line.

The Khazar and Baku terraces are buried under the Khvalynsk sediments. The rising steps of terraces can occur only in zones of intense recent upheavals, but the Syrt Trans-Volga region from the very beginning of the Quaternary always was a zone of downwarping.

Some authors, in speaking of the Baku terrace of the Volga, point out that sandy rocks not typical of the Syrt stratum are developed near the river valley. They believe them to be Baku (Mindel) alluvium. However, A.N. Mazarovich [14], in his time, and other investigators have shown that these arenaceous sediments of the "Mendel terrace" are, in the east, replaced along the strike by true Syrt deposits.

REFERENCES

1. Vasil'yev, Yu.M., O sledakh proyavleniya merzlotnykh protsessov v chertvertichnykh otlozheniyakh Severnogo Prikaspiya. [TRACES OF FREEZING PHENOMENA IN THE QUATERNARY DEPOSITS OF NORTHERN CISCASPIAN]: Izv. Akad. Nauk SSSR, ser. geol., No. 12, 1958.
2. Vasil'yev, Yu.M., Chetvertichnyye otlozheniya Severnogo Prikaspiya. [THE QUATERNARY DEPOSITS IN NORTHERN CISCASPIAN]: Izv. Akad. Nauk SSSR, ser. geol., No. 5, 1959.
3. Goretskiy, G.I., O sootnosheniyakh morskikh i kontinental'nykh osadkov Priazov'ya, Primanych'ya i Nizhnego Pridon'ya. [RELATIONSHIPS BETWEEN THE MARINE AND CONTINENTAL SEDIMENTS IN THE AZOV, MANYCH, AND LOWER DON REGIONS]: Tr. Komis. po izuch. chetvertich. perioda, vyp. 13, 1957.
4. Goretskiy, G.I., O vozraste drevnikh alluvial'nykh svit antropogena, pogrebennykh v dolinakh Volgi i Kamy. [THE AGE OF THE EARLY QUATERNARY ALLUVIAL FORMATIONS BURIED IN THE VOLGA AND KAMA RIVER VALLEYS]: Doklady, Akad. Nauk SSSR, t. 110, No. 5, 1956.
5. Grishchenko, M.N., and A.I. Koptev, Materialy k stratigrafii terrasovykh otlozheniy doliny Volgi u s. Privolzh'ye. [MATERIALS ON THE STRATIGRAPHY

- OF THE TERRACE DEPOSITS IN THE VOLGA VALLEY NEAR PRIVOLZH'YE VILLAGE]: Tr. Voronezhsk. un-ta, t. 39, 1955.
6. Gromov, V.I., Stratigraficheskoye znachenie chetvertichnykh mlekopitayushchikh Povolzh'ya. [THE STRATIGRAPHIC SIGNIFICANCE OF THE QUATERNARY MAMMALS IN THE VOLGA REGION]: Tr. Komis. po izuch. chetvertich. perioda, t. 4, vyp. 2, 1935.
 7. Gromov, V.I., Paleontologicheskoye i arkheologicheskoye obosnovaniye stratigrafii kontinental'nykh otlozheniy chetvertichnogo perioda na territorii SSSR (mlekopitayushchiye paleont.). [PALEONTOLOGICAL AND ARCHEOLOGICAL SUBSTANTIATION OF THE STRATIGRAPHY OF THE QUATERNARY CONTINENTAL DEPOSITS ON THE USSR (PALEONTOLOGY OF MAMMALS)]: Tr. In-ta geol. nauk Akad. Nauk SSSR, vyp. 64, geol. ser., No. 17, 1948.
 8. Doskach, A.G., and I.P. Gerasimov, Geomorfologicheskii ocherk nizhnego uchastka doliny r. Ural ot pos. Kalmykovo do g. Gur'yev. [GEOMORPHOLOGICAL OUTLINE OF THE LOWER PART OF THE URAL RIVER VALLEY FROM KALMYKOVA TO THE GUR'YEV]: Tr. In-ta geogr. Akad. Nauk SSSR, vyp. 24, 1937.
 9. Zhukov, M.M., K stratigrafii Kaspiyskikh osadkov Nizhne-Volzhskogo kraia. [STRATIGRAPHY OF THE CASPIAN SEDIMENTS IN THE LOWER VOLGA AREA]: Tr. Komis. po izuch. chetvertich. perioda, t. 4, vyp. 2, 1935.
 10. Zhukov, M.M., Plitsenovaya i chetvertichnaya istoriya severa Prikaspiyskoy vpadiny. [PLIOCENE AND QUATERNARY HISTORY OF THE NORTHERN PART OF THE CASPIAN DEPRESSION]: V kn. "Problemy Zapadnogo Kazakhstana", t. 2. Izd-vo Akad. Nauk SSSR, 1945.
 11. Kipiani, M.G., and A.D. Kolbutov, Novyye dannyye po stratigrafii chetvertichnykh otlozheniy Povolzh'ya i Prikam'ya. [NEW DATA ON THE STRATIGRAPHY OF THE QUATERNARY DEPOSITS IN THE VOLGA AND KAMA REGIONS]: V kn. "Tezisy dokladov Vses. mezhdruvedomstv. soveshch. po izuch. chetvertich. perioda", 1957, Russkaya ravnina. Izd-vo Akad. Nauk SSSR, 1957.
 12. Lavrushin, Yu.A., O sushchestvovanii v Srednem Povolzh'ye "beloyarskoy terrasy". [THE EXISTENCE OF THE "BELOYAR TERRACE" IN THE MIDDLE VOLGA REGION]: Izv. vyssh. shkoly. Geol. i razvedka, No. 5, 1959.
 13. Lyuttsau, S.V., Novyye dannyye po geomorfologii volzhskikh terras rayona Stalingradskogo vodokhranilishcha. [NEW DATA ON THE GEOMORPHOLOGY OF THE VOLGA TERRACES IN THE REGION OF THE STALINGRAD WATER RESERVOIR]: Vestn. Mosk. un-ta, ser. fiz. -mat. i yestestv. nauk, No. 10, 1952.
 14. Mazarovich, A.N., Terrasy Volgi i chetvertichnyye otlozheniya Zavolzhskikh stepey. [THE VOLGA TERRACES AND THE QUATERNARY DEPOSITS OF THE TRANS-VOLGA STEPPES]: Byul. Inform. byuro Assots. po izuch. chetvertich. perioda Yevropy, No. 3-4, 1932.
 15. Mozharovskiy, B.A., Geologicheskaya istoriya Saratovskoy kotloviny. [THE GEOLOGIC HISTORY OF THE SARATOV BASIN]: Izv. Nizhnevolzhsk. Krayevedcheskogo in-ta, vyp. 3, 1929.
 16. Moskvitin, A.I., Chetvertichnyye otlozheniya i istoriya formirovaniya doliny r. Volgi v yeye srednem techenii. [THE QUATERNARY DEPOSITS AND THE HISTORY OF THE FORMATION OF THE MIDDLE COURSE OF THE VOLGA RIVER VALLEY]: Tr. geol. in-ta Akad. Nauk SSSR, vyp. 12, 1958.
 17. Nikolayev, N.I., O vozraste chetvertichnoy volzhskoy fauny mlekopitayushchikh. [THE AGE OF THE QUATERNARY VOLGA MAMMALIAN FAUNA]: Byul. Mosk. o-va ispyt. prirody. Otd. geol., t. 15, vyp. 6, 1937.
 18. Popov, G.I., Istoriya Manychskogo proлива v svyazi so stratigrafiyey chernomorskich i kaspiyskich chetvertichnykh otlozheniy. [HISTORY OF THE MANYCH STRAITS IN RELATION TO THE STRATIGRAPHY OF THE BLACK AND CASPIAN SEA QUATERNARY DEPOSITS]: Byul. Mosk. o-va ispyt. prirody. Otd. geol., t. 30, vyp. 2, 1955.
 19. Pravoslavlev, P.A., O znachenii vertikal'nykh izmeneniy v okraske peschanoglinistykh porod v oblasti nizhnego techeniya rek Bol'shogo i Malogo Uzeney. [THE SIGNIFICANCE OF THE VERTICAL CHANGES IN THE COLORATION OF ARENACEOUS-ARGILLACEOUS ROCKS IN THE LOWER-COURSE REGIONS OF THE BOL'SHOY AND MALYY UZEN' RIVERS]: Iz. Ros. Akad. Nauk, No. 16, 1918.
 20. Pravoslavlev, P.A., Predisloviye k

- rabote V.I. Gromovoy "Novyye materialy po chetvertichnoy faune Povolzh'ya i po istorii mlekopitayushchikh Vostochnoy Yevropy i Severnoy Azii voobshche". [INTRODUCTION TO V.I. GROMOVA'S WORK "NEW MATERIALS ON THE QUATERNARY FAUNA IN THE VOLGA REGION AND ON THE HISTORY OF THE MAMMALS OF EASTERN EUROPE AND NORTHERN ASIA AS A WHOLE"]: Tr. Komis. po izuch. chetvertich. perioda, t. 2, 1932.
21. Stratigrafiya chetvertichnykh otlozheniy i noveyshaya tektonika Prikaspiyskoy nizmennosti. [STRATIGRAPHY OF QUATERNARY DEPOSITS AND THE MOST RECENT TECTONICS OF THE CASPIAN LOWLAND]: Izd-vo Akad. Nauk SSSR, 1953.
22. Fedorov, P.V., Stratigrafiya chetvertichnykh otlozheniy i istoriya razvitiya Kaspiyskogo morya. [STRATIGRAPHY OF QUATERNARY DEPOSITS AND THE HISTORY OF THE CASPIAN SEA]: Tr. Geol. in-ta Akad. Nauk SSSR, vyp. 10, 1957.
23. Shantser, Ye.V., Allyuviiy ravninnykh rek umerennogo poyasa i yego znacheniye dlya poznaniya zakonomernostey stroyeniya i formirovaniya allyuvial'nykh svit. [ALLUVIUM OF THE PLAINS RIVERS OF THE TEMPERATE-ZONE AND ITS SIGNIFICANCE FOR UNDERSTANDING THE STRUCTURAL AND FORMATION REGULARITIES OF ALLUVIAL FORMATIONS]: Tr. In-ta geol. nauk Akad. Nauk SSSR, vyp. 135, ser. geol., No. 55, 1951.
24. Shchukina, Ye.N., Terrasy Verkhney Volgi i ikh sootnosheniye s lednikovymi otlozheniyami Gor'kovsko-Ivanskogo kraya. [THE UPPER VOLGA TERRACES AND THEIR RELATIONSHIP TO THE GLACIAL DEPOSITS IN THE GOR'KOVSKO-IVANOVSKIY AREA]: Byul. Mosk. o-va ispyt. prirody. Otd. geol., t. 11, No. 3, 1933.

Geological Institute of the
U.S.S.R. Academy of Sciences,
Moscow

Received February 24, 1960

METHODS

SEPARATION OF CLAY PARTICLES BY ELECTROPHORESIS¹

by

N. V. Logvinenko, and A. A. Lazarenko

The separation of clay fractions ($<1\mu$) is a very lengthy and cumbersome operation. In order to produce the few grams of this fraction required for a complex mineralogical investigation (by means of thermography, chemical and X-ray analyses), it is usually necessary to precipitate for several days, and then to evaporate clay suspensions. For large-scale operations this method is inadequate.

These difficulties can be overcome generally by utilizing the method of electrophoresis. Essentially it consists of the following. Clay particles dispersed in water from colloidal systems, and the particles carry negative electric charges on their surfaces. In the field of a direct current these particles migrate to the anode where they are discharged and deposited. This process ordinarily is not used for coarse particles of clay minerals and accessory minerals.

It is reasonable to assume that if other negatively charged colloids (humate, silica) are present in the rock, they, too, will settle on the anode.

The method of electrophoresis (often erroneously called electroosmosis) has been utilized for the technical enrichment of clays and kaolins for a long time. A diagram of the equipment used for these purposes is given in Figure 1 (1.3). A dense clay suspension is pumped from a mixer through tube d into a tank, where it is maintained in a suspended state by means of agitators c. When the electric current is switched on, the clay particles, in passing through the cathode grid b, are attracted to the anode surface a, where they settle and are removed by scraper f. The voltage required for this operation amounts to 75-100 v, the current is 100 a, and the density does not exceed 0.01 a/cm². The output of the unit is several tons of enriched product per day (approximately 1 t per 30-70 kw/h). A prerequisite condition for the in-

dustrial application of electrophoresis is the addition of alkaline electrolytes - peptizing agents (liquid glass, alkali) to impart greater stability and to increase the electrokinetic properties of the clay particles in suspension.

Electrophoretic enrichment of clays is practiced both abroad and in domestic industries.

To produce a few grams or tens of grams of fine clay particles in lithological laboratories no cumbersome installations with high current parameters are required. Depending on the available resources it is possible to utilize various operational techniques. However, to assure adequate efficiency and rapid separation of clay particles, a number of conditions must be observed. The most important of these are: 1) sufficient current density at the anode immersed in the suspension: up to 0.010-0.005 a/cm²; 2) stability and sufficient density of the clay suspension (not less than 1.005-1.010); 3) optimum distance between the electrodes, 15 - 25 mm; 4) source voltage of 75-150 v.

Suspensions utilized under laboratory conditions must not contain electrolyte ions, since they are being sorbed by the clay particles and the latter may change their chemical composition or coagulate (in the presence of positively charged electrolytic particles).

In designing electrophoresis equipment the most important condition is the selection of a sufficiently powerful direct-current source, preferably with controllable parameters. VSA-5, continuous control selenium rectifiers, or type VSA-6, which may be satisfactorily controlled by LATR-1 transformer are suitable for this purpose. Use should be made of converters (motor-generators) with current of up to 5-10 a and 65-150 v. The very popular gas-filled tube rectifiers produced by the Elektrodelo Plant, usually lack sufficient power (6a, 24 v), particularly with respect to rectified voltage. This defect may be eliminated by connecting a few of these units in series. This method is also applicable for combinations with different current sources.

¹Vydeleniye glinistyykh chastits pri pomoshchi elektroforeza. pp. 100 - 103.




FIGURE 1. Diagram of a machine used for clay enrichment by the electrophoresis method.

(After Schwerin, 3).

Upon selection of a suitable direct-current source, it is necessary to calculate the anode area to make sure that the current density on the surface immersed in the suspension will, under average conditions, amount to 0.01 a/cm^2 . The density of current, evidently, depends not only on its source, but also on the distance between the electrodes, as well as on the density of the suspension. In order to increase its strength the anode can be plated by pure lead, or, even better, by an alloy of lead containing antimony (up to 15% Sb). Utilization of platinum for electrodes is impracticable in view of their large surface and correspondingly large quantities of metal required. The cathode, in the form of a copper or brass grid with 0.25-0.10 mm openings, should be approximately of the same size. The distance between the electrodes and the degree of anode immersion into the suspension must be regulated.

Depending on the size of the electrodes and the volume of the clay suspension a corresponding bath may be selected. Usually a glass crystallizer may be used for this.

To control the operating conditions it is necessary to include an ammeter and voltmeter in the circuit. It is useful also to provide for a thermometer to take the temperature of the suspension, which may rise very considerably between the electrodes because they are too close or there is a bad contact between them and the leads. The latter must be well insulated. Excessive current strength, as compared with its normal value, may be reduced by inserting a rheostat into the circuit.

The circuit diagram of a laboratory unit is shown in Figure 2.

The quantity of clay produced by electrophoresis is, as was previously stated, a function of the current source parameters, suspension density, distance between the




FIGURE 2. Basic diagram of the laboratory unit used for electrophoresis.

electrodes, current density on the anode, time, and other factors. The effects of these factors can be judged from the data contained in Table 1.

The initial material for the investigation was a suspension (size of particles $<5\mu$) of the same rock - Jurassic hydromicaceous clay taken from the northwestern part of the Donetz basin. In all cases, an anode in the form of a round lead plate having an over-all area of 450 cm^2 was inserted in a horizontal position into the suspension whose volume was about one liter.

Table 1 shows that the density of current during the operation was, on the whole, insignificant: $0.0025-0.0050 \text{ a/cm}^2$, mainly due to the relatively large surface of the anode. This indicates that the method of clay electrophoresis is operative even under such unfavorable conditions.

Generally, however, because of low suspension density and the absence of electrolytic additions, in operations designed for special purposes and conducted under laboratory conditions, the deposition of clay particles per identical unit time and unit anode area is smaller than in the industrial enrichment processes.

Greater efficiency can be achieved by increasing the current-source parameters and augmenting the density of the initially prepared suspension. For example, in the Petrography Department Laboratory of Khar'kov University, a 180-v and 13-a converter serves as the source of current. The strength of the current in the circuit usually amounts to $2.5 - 5 \text{ a}$. The area of the anode lead cover is 37 cm^2 . With a density of suspension of $1.008-1.011$, the output of clay

Table 1

Effect of different conditions in the electrophoresis process.

Variables	Voltage	Current strength in amperes	Distance between the electrodes, mm	Density of suspension	Duration of operation, min.	Current density at anode, a/cm ²	Quantity of separated clay (dry), g	Remarks
Voltage (and strength) of current	55	0,25—0,35	20	1,010	6	0,0006—0,0008	0,80	Coagulation of suspension during operation
	75	0,60—0,80	20	1,010	6	0,0013—0,0018	1,08	
	100	1,00—1,25	20	1,010	6	0,0022—0,0028	1,49	
	125	1,30—1,80	20	1,010	6	0,0029—0,0040	1,98	
	150	1,50—2,50	20	1,010	6	0,0033—0,0056	5,67	
Density of suspension	100	0,60—0,80	20	1,005	10	0,0013—0,0018	0,97	Noticeable coagulation of suspension
	100	0,85—1,10	20	1,010	10	0,0019—0,0024	2,15	
	100	0,90—1,20	20	1,015	10	0,0020—0,0027	2,35	
	100	1,00—2,85	20	1,020	10	0,0022—0,0063	2,87	
	100	1,00—2,90	20	1,030	10	0,0022—0,0065	4,41	
	100	1,00—5,40	20	1,050	10	0,0022—0,0120	5,11	
Distance between the electrodes	100	3,00—10,00	5	1,005	4	0,0067—0,0222	1,29	Very considerable heating of suspension - termination of operations Considerable heating of suspension
	100	1,90—7,00	10	1,005	6	0,0042—0,0156	1,66	
	100	1,25—1,50	15	1,005	7	0,0028—0,0033	2,10	
	100	1,30—1,15	20	1,005	7	0,0029—0,0026	2,13	
	100	0,80—0,50	25	1,005	7	0,0018—0,0011	1,99	
	100	0,70—0,50	30	1,005	7	0,0016—0,0011	1,86	
	100	1,00—1,25	20	1,010	5	0,0022—0,0028	1,25	
Times	100	0,85—1,10	20	1,010	10	0,0019—0,0024	2,15	Noticeable coagulation of suspension
	100	1,00—2,00	20	1,010	15	0,0022—0,0044	2,50	
	100	1,00—2,10	20	1,010	20	0,0022—0,0047	2,85	
	100	1,00—2,10	20	1,010	20	0,0022—0,0047	2,85	

NOTE: Commâ represents decimal point.

at the anode amounts to 20-30 g per 6-8 min of work.

Efforts should be made to assure the greatest possible degree of dispersion and purity of the clay particles deposited at the anode by systematic microscopic checks. The data obtained indicate that sufficiently pure clay is precipitated at the anode from the suspension with particles less than 5μ in diameter and there are almost no fragmental grains of quartz and other minerals.

The moisture of the separated clay usually

exceeds 50% (amounting to 65-80%). It is in inverse relation to the magnitude of voltage and current strength, as well as to suspension density.

General Sequence of Operation. A stable suspension in a volume of 1-2 liters, (depending on the size of the apparatus), is prepared from a corresponding batch of clay rock and distilled water. One hour and 20 minutes later the top 10 cm of suspension containing particles measuring less than 5μ are decanted and used for electrophoresis. The density of the suspension measured by an areometer

must not be less than 0.005-0.010. The suspension is poured into a bath in which the electrodes have been previously installed. The optimum distance between them should be 15-25 mm. After a few minutes of work the anode is withdrawn from the suspension and clay is removed from its surface. A few smears are made on the object glass for microscopic verification of the purity of the deposits. If the amount of clay produced is insufficient the process may be repeated, with the current increased by putting the electrodes closer together. After the work is finished, the electrodes should be thoroughly washed and rinsed in distilled water.

A set of different sizes of electrodes should be available for each electrophoresis apparatus. In varying the different procedural parameters, it is possible in each particular case to create the most favorable conditions for deposition of clay fractions.

REFERENCES

1. Lysin, B., Ochistka gliny pri pomoshchi elektricheskogo toka. [CLAY REFIN-

ING WITH THE AID OF AN ELECTRICAL CURRENT]: *Keramika i steklo*, No. 5, 1925.

2. Kukolev, G.V., *Tekhnicheskiye svoystva glin v svete kolloidno-khimicheskikh vozzreniy*. [TECHNICAL PROPERTIES OF CLAYS FROM THE COLLOIDAL-CHEMICAL POINT OF VIEW]: *Ogneupory*, No. 12, 1948.
3. Eyken, A., *Elektricheskiye i magnitnyye sposoby razdeleniya materialov*. [ELECTRIC AND MAGNETIC METHODS FOR SEPARATING MATERIALS]: ONTI, Khar'kov, 1938.
4. Hind, S.R., *Transactions of the British Ceramic Society*. London, vol. 24, No. 73, 1924.

Khar'kov State University

Received April 2, 1960

THE 1961 LAUREATES OF THE LENIN PRIZES¹

by

N. M. Strakhov — Originator of the Theory of Lithogenesis

The appearance of N. M. Strakhov's monograph, The Principles of the Theory of Lithogenesis, for which the author was awarded the well-deserved, highest prize — the Lenin Prize —, came as no surprise to geologists. This event was well argued by Nikolai Mikhaylovich's numerous previous works and his entire scientific activity. Beginning in 1937, N. M. Strakhov, with the natural talent and concentration so characteristic of him, focused all of his efforts on the elaboration of the theory of the sedimentary process, believing that the method of comparative lithologic analysis would lead him more directly to his goal. N. M. Strakhov's monograph is proof that he selected the correct path. In terms of its value to the science of lithology, there is no work equal to his Principles of the Theory of Lithogenesis, either in this country, or abroad.

The first volume of the monograph contains the doctrine on the types of lithogenesis: glacial, humid, arid, and effusive-sedimentary. A detailed scheme is given for the humid-type of lithogenesis, a thorough analysis is made of the processes occurring in catchment areas, of the mobility of various elements in the weathered mantle, the laws of mechanical and chemical denudation, the types of fluvial transportation of materials, the origin of sediments in basins, as well as of the diagenetic and epigenetic changes in the process of transformation of sediments into rocks.

Having discussed the stages of humid-type lithogenesis, the author proceeds to the description of humid formations as related to the paragenesis of sedimentary rocks, and in doing so, bases his discussion on a clearly-defined genetic concept of the process. He establishes the decisive role played by topography and structural environment in the formation of humid rocks and their associations, and defines the peculiarities of formations

developed on the platforms and in geosynclines at different stages of their development. Against a background of minutely described humid lithogenesis, the main features of the remaining three types of rock formation — glacial, arid, and effusive-sedimentary — become more graphically apparent. In discussing the distribution of the climatic types of lithogenesis on the Earth's surface in Post-Proterozoic time, the author develops a pattern of past climatic zonality.

N. M. Strakhov conclusively demonstrates the advantages of lithologic indicators over the paleontological for the reconstruction of climate. These chapters are illustrated with very interesting paleoclimatic maps. Since the maps are based on the distribution of such indicator-rocks as bauxites, iron, and manganese ores, the crust of weathering, coals, rock and potash salts, they also simultaneously serve as patterns of zonal distribution for commercial minerals throughout the entire period of the Earth's geologic history. The analysis of paleoclimatic maps leads the author to a conclusion concerning the repeated migrations of all climatic zones throughout the Earth's history, associated with the changing positions of its axis of rotation relative to the mantle.

The second volume of the monograph is devoted to an analysis of the regularities governing the composition and distribution of humid deposits. Considered attention is paid thereby to the conditions and the mechanism of formation of high aluminum, iron, manganese, phosphorus, CaCO_3 , MgCO_3 , and SiO_2 concentrations, as well as to polarity zones in the distribution of rare and dispersed elements. A step-by-step examination is made in this volume of the granulometric composition of silty-argillaceous rocks of various types of facies formed under varying climatic and structural environments; delineated are the distribution zones of components the complex forms of migration: iron, manganese, phosphorus, and minor elements. Sedimentary ore deposits are subdivided, for the first time in the literature, into three facies-genetic groups.

¹ Laureaty Leninskikh Premiy 1961. N. M. Strakhov — Osnovopolozhnik Teorii Litogeneza. pp. 104 — 105.

Identified for each ore component, is its facies profile, and the formative mechanism of ore concentrations. The emplacement of ores in synchronous tectonic structures and metalliferous formations is analyzed, and the evolution of ore deposition in the history of the Earth discussed.

The last chapters of the second volume are restricted to a detailed examination of the diagenesis of subaqueous deposits, the basic premises for which were formulated in 1953. In the new monograph, the author delved deeper into the substance of minerogenetic processes and the redistribution of matter in diagenesis. He also outlined the development peculiarities of diagenetic processes in basins of varying physical and geographic characteristics.

This year will see the publication of the third volume of his Principles of the Theory of Lithogenesis devoted to arid lithogenesis. Discussed here exhaustively are the particu-

lars of sedimentation and ore-formation in an arid environment, the basic features of modern halogenesis, as well as halogenetic evolution in Post Algonquian time.

Manifest in the third volume, just as in the preceding two, is N. M. Strakhov's persistent attempt to disclose the physiochemical essence of the processes involving the formation of sedimentary rocks.

At the present time N. M. Strakhov directs his own efforts and those of his associates toward the study of the geochemistry of sedimentary rocks. He has advanced the problems of the facies-genetic relations between the scattered concentrations of elements and their ore accumulations in humid zones. Solution of this problem will make it possible to understand the distribution and accumulation mechanism of elements, i. e., brings us closer to the development of the theory of geochemical processes, and help us to provide better substantiation in diagnosing ore occurrences.

REVIEWS AND DISCUSSIONS

THE PROBLEM OF THE NOMENCLATURE OF EFFUSIVE ROCKS¹

by
T. Ya. Goncharova

In Ye. K. Ustiyev's article The Problem of the Nomenclature of Effusive Rocks, published in *Izvestiya Akademii nauk SSSR, seriya geologicheskaya*, No. 11, 1959, an important problem was raised concerning the unfortunate situation concerning the classification of effusive rocks which is causing a great deal of concern among geologists engaged in the study of sedimentary-volcanic complexes and the related ore deposits. The poorly developed nomenclature of effusive rocks and their pyroclastic varieties considerably hampers the detailed mapping of mining districts and complicates the study of deposits enclosed in extrusive rocks.

Ye. P. Ustinov's article and the critical reviews published by S. N. Ivanov and V. I. Lebedinskiy in the *Izvestiya AN SSSR, seriya geologicheskaya*, No. 7, 1960, discuss the advisability of two parallel nomenclature systems for fresh cenotypal and altered paleotypal effusive and subvolcanic rocks.

Ye. K. Ustiyev proposed that the two parallel nomenclatures for effusive rocks be abandoned and a single nomenclature scale corresponding to the cenotypal series be adopted. Rock variations in his opinion, could be indicated by adding the word "paleotypal" or, by the words "altered" or "decomposed" which are preferred in the Russian language and which more accurately reflect their meaning.

S. N. Ivanov and V. I. Lebedinsky strongly object to Ye. K. Ustinov's proposals and defend the necessity of maintaining a double nomenclature for effusive rocks.

The above papers, it seems to us, outline two different trends in the specification of classification principles: principal emphasis is on the primary properties of rocks, or,

conversely, the secondary properties of effusives acquired as a result of subsequent metamorphism.

We believe that both property of extrusive rocks are equally important.

1. The primary character of effusive rocks, which can be discerned in the altered varieties only after thorough analysis of their optical properties and chemical composition, makes it possible to reconstruct the conditions under which the rock was formed and to relate it to a definite magma. Only the primary properties of a rock can furnish the correct representation of the geologic environment in which it was formed, information of crucial importance for further geologic and metallogenic generalizations

2. Of no less importance are the secondary properties of effusive rocks acquired in the process of aging, and by structural and metamorphic transformations. The present state of a rock certainly reflects its geologic history, since the rocks, as was aptly stated by V. I. Lebedinskiy, are the monuments of geological processes.

Thus, both of these elements must be reflected in the names of effusive and subvolcanic rocks.

In our opinion, composition-wise more correct and more convenient to use is that classification of extrusive rocks in which the names of both, the fresh and the altered, lavas of one genetic series have the same roots. We, therefore, propose to preserve the roots of the nomenclature series embracing the cenotypal effusives for all of their altered analogs. To express the subsequent transformations experienced by the rocks, it is recommended to append characteristic prefixes (or suffixes).

For example, in taking the cenotypal family of effusives given in Ye. K. Ustiyev's paper as a basis, it is possible, in accordance with M. A. Usov's views concerning the existence of three constitutional phases of effusive rocks, to isolate three petrographic series.

¹K voprosy o nomenklature effuzivnykh gornykh porod. pp. 106 - 107.

Cenotypal	Paleotypal	Greenstone (metamorphic)
1. Liparites	1. Paleoliparites	1. Metaliparites
2. Sodium liparites	2. Sodium paleoliparites	2. Sodium metaliparites
3. Liparite-dacites	3. Paleoliparite-dacites	3. Metaliparite-dacites
4. Dacites, etc.	4. Paleodacites	4. Metadacites

We do not aspire to have precisely this classification scheme introduced for effusive rocks — this is merely an example of one of the possible variants. It is quite possible that other more appropriate prefixes and suffixes ("apo" instead of "paleo", suffix "oil" instead of prefix "meta") may be found. This matter should become the subject of further discussions. However, we believe that the principle used as the basis of the proposed scheme, is the most suitable of all.

We are firmly convinced that the classification of paleotypal and metamorphized effusives should be revised. The long, and often unsuitable, designations of effusive rocks must be replaced by new terms, which should, of necessity, retain the roots of the initial cenotypal analogs. The presence of roots pertaining to cenotypal effusives in the names of the altered varieties emphasizes their consanguinity, pertinence to a single genetic series, and considerably facilitates the development of a nomenclature scheme. Along with further elaboration and detailed development of the classification for cenotypal effusive series, the nomenclature of altered rocks will be correspondingly improved. All the identified varieties of cenotypal effusives, in-

cluding those of intermediate composition, which as yet have no analogs in the petrographic nomenclature for altered series, will find their place in the proposed classification scheme. The new terminology can be adapted easily to pyroclastic rocks.

Despite the many years of study on the old, predominantly Paleozoic, strongly altered volcanic formations surrounding the chalcopiritic deposits in the Caucasus, Altay, and partially in the Urals, we cannot share S. N. Ivanov's apprehensions concerning the great confusion which will arise if, in following Ye. K. Ustiyev's suggestion, the paleotypal designations of rocks are replaced by cenotypal terms on the geologic maps of the regions where altered effusives predominate. On the contrary, we feel that the maps can only gain thereby and will be of greater value. This, of course, provided this replacement is made not on the basis of "the discretion of individual map authors" (S. N. Ivanov), but after a detailed and considered petrographic study of the effusives. A reflection of the characteristic alteration peculiarities in the description of rocks is also a prerequisite condition.

The identification of the primary aspect of altered effusives on the basis of relict mineralogical composition and texture, and the reconstruction of their geologic life, represent one of the main tasks in the petrography of volcanic complexes.

Thus, the suggestions briefly outlined in this article should be considered as a further expansion and development of Ye. K. Ustiyev's ideas.

Received November 1, 1960

THE NOMENCLATURE OF EFFUSIVE ROCKS¹

by
I. M. Speranskaya

The symposium on the classification, nomenclature, and terminology of volcanic rocks constituted an important part of the First All-Union Volcanological Conference. However, the overwhelming majority of papers dealing with this subject was devoted to pyroclastic rocks, whose classification is still very poorly developed. Only one author — Ye. K. Ustiyev — paid serious attention to the sub-

stantial defects in the nomenclature and the classification principles of paleovolcanic extrusive rocks. Later on, he published a special article restricted to the classification of effusives (Izvestiya AN SSSR, seriya geologicheskaya, No. 11, 1959), in which he again stressed the need to regulate the nomenclature and suggested that the nomenclature referring to "paleotypal" lavas be discarded.

The development of more precise specifications for the classification principles and the existing nomenclature of effusive rocks is long

¹ O nomenklature effuzivnykh gorn'nykh porod.
pp. 107 - 108.

since overdue. At the Volcanological Conference, attended by the leading petrographers-volcanologists of the Union, the speakers often used a very diversified nomenclature for effusives which was not always reconcilable with any definite principle of classification. This situation has a bad effect on the geological surveys carried out by field geologists in different volcanic regions of our country.

For purposes of relatively small-scale geologic mapping, an important segment of field geologic operations, the problem of a coordinated nomenclature for effusive rocks is, at present, of particular importance. This is due to the fact that, in various old and recent volcanic regions throughout the USSR, a great deal of factual material on the geology and petrography of volcanic formations has been assembled and still continues to accumulate. One of the methods by means of which all this material could be generalized is the preparation of the State geologic maps (on various scales) for the entire territory of the Union. To facilitate the future comparisons of volcanism in various regions, it is advisable that a single nomenclature relative to the effusive rocks should be used in the legends of these maps. Considering the very brief explanatory notes accompanying the maps, the terminology problems are particularly important.

During the many years devoted to the study of Mesozoic effusives in the Okhotsk-Chaun volcanic zone in Northeast Asia, and to the editing of geologic maps covering this territory, I was repeatedly confronted with the great confusion prevailing in the terminology of effusive rocks. This creates considerable problems with respect to mapping, and sometimes it is impossible to make conclusive comparison of the available materials with the published data.

At the same time, it should be emphasized that a considerable proportion of geologists working on Northeast Asia utilize a single "cenotypal" classification of effusives. At the Interdepartmental Stratigraphic Conference (Magadan, 1957), where problems of the stratigraphic differentiation of eruptive formations were also discussed, the authors of many reports used "neo" volcanic nomenclature. The widespread use of "cenotypal" nomenclature for effusive rocks by the geologists concerned with the Northeast is not accidental. This is primarily attributable to the peculiarities of the Mesozoic stages in the geologic-tectonic history of Northeast Asia, where the most powerful manifestations of volcanism are related to Upper Mesozoic and Cenozoic tectonics. In the Okhotsk-Chaun volcanic zone, which used to serve as the main arena for powerful Cretaceous volcanism, a major tectonic reconstruc-

tion must have, apparently, occurred in the Pre-Devonian time. Considerable tectonic movements are found also on the boundary between the Cretaceous and Paleogene.

Thus, Cretaceous volcanism developed under conditions of considerable tectonic mobility, which was particularly typical during its early stages. The volcanic rocks are, therefore, altered in varying degrees. Although a general strengthening of the "paleotypal" characteristics was established as we progress from younger to older rocks, the degree of secondary alteration, even in the synchronous effusives, was found to be extremely uneven. Among the relatively cenotypal lavas there sometimes appear considerable zones of paleotypal rocks. Moreover, in the neighboring outcrops, or even in the very same ones, it is possible to observe extrusive rocks forming one and the same cover, but sharply differing in the degree of secondary alterations. Here one may observe the entire range of transitions from the fresh to strongly altered rocks with identical composition. Under these circumstances, the determination of the degree of "paleotypal" development (and, consequently, also the use of one term or another) becomes, to a considerable extent, a matter of subjective judgement. In connection with the unavoidable subjectivity of determinations, different nomenclatures are frequently used to designate effusive rocks of the same formation in different regions. This makes the correlation of stratigraphic sections exceedingly difficult.

Only the utilization of a single "neo"-volcanic nomenclature would permit an objective characteristic to be given to rocks typical of volcanic complexes, and to single out areas having special types and different degree of secondary alterations, without isolating these rocks from the single complex through the application of varying designations. Exclusive utilization of "cenotypal" classification also may have a favorable effect on the methods used in the investigations of effusives.

In studying, for example, a stratum of relatively cenotypal andesite lavas, the investigator, upon finding a zone of altered rocks, will no longer be able to simply call them "porphyrites". Experience shows that the following cases are not atypical in geologic mapping: when "formations of porphyrites" are identified without sufficient analysis of the altered rocks, while, in reality, they have no independent stratigraphic importance. In making use of the "cenotypal" classification, the geologist necessarily will have to compare the altered rock with the same type of effusive rocks of cenotypal aspect, to identify the nature of secondary alterations, and only then draw a conclusion as to the relationship between the cenotypal and paleotypal representatives.

Thus, in describing the Mesozoic volcanic formations in Northeast Asia, one should refrain from using a dual nomenclature for the effusive rocks, especially, because this territory is a component of the enormous Pacific volcanic belt, for a large section of

which (The American littoral, the Japanese Islands, and China) geologists use a uniform "cenotypal" nomenclature for effusive rocks.

Received September 6, 1960

MAIN RESULTS OF THE DISCUSSION ON THE NOMENCLATURE OF EFFUSIVE ROCKS

by
Ye. K. Ustiyev

Sometime ago I made a report [15] and published a paper [14] proposing a change in the nomenclature of effusive rocks. The main objective of this long overdue reform is the elimination of a dual set of designations for effusive rocks differing only in geologic age or degree of post-volcanic alteration. The following quotation sums up the basic conclusions contained in these papers: "The 'nomenclatural' subdivision of extrusive rocks into 'young' and 'old', or 'paleotypal' and 'cenotypal', is historically inherited from European, (including the Soviet) petrographers from the early development stages of geology and petrography in Central Europe. This hangover of a long past stage, which even in the last century had an inhibitive effect on the development of petrographic nomenclature, has, in our time, become a crying anachronism."

A general liquidation of the entire nomenclature table relating to the "paleotypal" lavas will contribute to the removal from petrography of a number of superfluous, logically unsubstantiated, designations, and will lead to greater uniformity and harmony in the world's petrographic terminology.

As one might have expected, the suggested regulation of the nomenclature for effusive rocks attracted the attention of many petrographers and gave rise to lively discussion. At present, it has become definitely possible to sum up the published remarks.

T. V. Dianova [2] writes that "proposal made by Ye. K. Ustiyev is timely and fundamentally correct". N. D. Zlenko and M. A. Tarkhova [4] believe that the "article by Ye. K. Ustiyev is very timely, and the positions stated in it are definitely progressive". T. Ya. Goncharova [1] and I. M. Speranskaya

[10] also insist on the necessity of correcting the "essential defects in the nomenclature and classification principles" of paleovolcanic effusives and their pyroclastic varieties. Both authors feel that these shortcomings are reflected not only in the present-day state of our theoretical petrography, but also "strongly hamper the mapping of mining districts and complicate the study of the deposits in the volcanic rocks" (T. Ya. Goncharova).

Almost all of these authors not only declare themselves in favor of the proposed uniform nomenclature for effusive rocks, but further develop it in specifying certain details of the set of designations for altered lavas and the comagmatic extrusive rocks of veined nature. Such initiative should be welcomed. It attests not only to the vitality of the idea, but also to the inherent promise of progress, both in the theoretical and applied (geologic mapping) aspects of petrography.

The essence of the conclusions arrived at by all five of the authors referred to above may be summarized in the following four [sic] points: 1) a fundamentally single nomenclature should be adopted for the effusives of any age or stage of alteration; 2) as a basis for further improvement of the nomenclature for altered effusives, it is necessary to make use of the nomenclature for the cenotypal family of lavas distinguishable by particular richness and flexibility due to the historical conditions of petrographic development; 3) the designations of effusive rocks must be derived by means of adjectives, suffixes, and prefixes, retaining the root which indicates a corresponding cenotypal lava, for example: basalt — altered basalt (paleobasalt) — metamorphosed basalt (metabasalt) — deeply metamorphosed basic effusive rock (metabasite); 4) the nomenclature of effusive rocks must be made to correspond

¹ Osnovnyye itogi o nomenklature effuzivov. pp. 109 — 114.

with the parallel series of intrusive and vein rocks.

It is easy to perceive the harmony in spirit and form of these conclusions with the proposals which were used as their basis. Quite reasonable is the suggestion that the details of this problem be considered by a special nomenclature commission. It is obvious that the recommendations of such a commission must be decisive for the development of a corresponding division in our petrography.

At the same time it would have been wrong to believe that the discussed reform in the nomenclature of effusive rocks would meet with opposition. It is a well-known fact that even the most modest progress in science is always confronted with obstacles and cannot be achieved without overcoming the "traditions" it contradicts.

Two of the published reviews — one by V. I. Lebedinskiy [6], the other by S. N. Ivanov [5] — reject the idea of a uniform nomenclature for effusives. Many of the arguments advanced have already been refuted by T. V. Dianova, N. D. Zlenko, M. A. Trakhova, T. Ya. Goncharova, and I. M. Speranskaya. Nevertheless, an additional analysis of the objection against the reform may prove useful.

I. V. Lebedinskiy's criticism is based on a perfectly correct idea that it is important to maintain "a geologic approach to effusive rocks" and that "anti-historical concepts" are inadmissible in petrography. However, lack of consistency in the discussion of various aspects of the problem has brought him to erroneous deductions drawn from a reasonable premise.

In dealing with the first question he writes: "Ye. K. Ustiyev for some reason forgets about M. A. Usov who has made a special study of the existence of conditions of extrusive rocks after their formation depending on the peculiarities of the geologic environment, and has developed, on this basis, their classification and nomenclature. . . M. A. Usov has identified three families of rocks — the primary phase (cenotypal), the diagenetized (paleotypal) and the greenstones. . .".

This approach is based on two errors. First of all, it is impossible to consider seriously that the "geologic approach" to the nomenclature of effusive rocks will be destroyed if "porphyrite" is to be called "altered (paleotypal) andesite", and "melaphyre" be referred to as "altered (paleotypal) basalt". At the same time, it is clear that the very opportunity of identifying in some ancient volcanic formation, instead of the rather vague "porphyrites", at least a more specific formation of paleotypal andesite-basalt, or

trachy-andesite (for which there is no designation in the "orthodox" nomenclature defended by V. I. Lebedinskiy), is already an advantage, since this saves the petrographer from the danger of distorting its geologic and magmatic history.

In addition, the assumption that "Ye. K. Ustiyev has forgotten about M. A. Usov" is wrong. In a very similar way, V. I. Lebedinskiy seems to have "forgotten" the proposal to divide into diagenetized "phases" not only effusive, but also intrusive, rocks.

As is well known, the classification of igneous rocks developed by M. A. Usov [12, 13] on the basis of this principle can be represented in the form of the table.

I am not aware, however, of any published work in which V. I. Lebedinskiy would have designated fresh granites as charnockites, granodiorites as kavkazites, and diorites as andendiorites; just as he fails to see the necessity of special designations even for the totally altered (uralitized, scapolitized, chloritized, etc.) gabbro, diorite and other rocks of intrusive origin. Thus, he obviously refuses to recognize a dual nomenclature for intrusive rocks, but cannot do without a dual nomenclature where effusive rocks are concerned! It may be clearly deduced therefrom that the matter in question is not the "forgetfulness" of this or that petrographer, but rather the consistency with which he develops his views on the nature of matter. It should be said that any strict classification of natural phenomena, first of all, requires uniformity of measurement units, or, at least, of the principles on which this classification system is based. However, it is precisely this uniformity in approach to the basic principles of classification and systematization of igneous rocks that V. I. Lebedinskiy, regretfully, lacks.

The opinion regarding the "anti-historic character" of a single nomenclature for effusive rocks easily can be refuted with the aid of V. I. Lebedinskiy's own materials. In the very recently examined [7] example of "spilitic-keratophyre" effusives of the Crimea, studied long ago by F. Yu. Levinson-Lessing [8], he described a lava stream, whose top is composed of andesites, the central part — of hydrothermally altered andesites (which, of course, are called by the author "porphyrites"), and the bottom — of almost totally reworked andesites (referred to as "keratophyres"). Thus, one and the same lava, from one and the same sheet, but considerably altered by post-volcanic processes, was given three different designations usually associated by all supporters of dual nomenclature with different historical diagenetic phases and remote families of rocks.

If such ideas of a natural process are to

Primary phases		Diagenetized phase		Greenstone phase	
Effusives	Intrusives	Effusives	Intrusives	Effusives	Intrusives
Liparite	Charnockite	Quartz-porphry	Granite (anoterite)	Quartz-keratophyre	—
Dacite	Kavkazite	Quartz-porphry	Granodiorite (opdalite)	Quartz-keratophyre	—
Andesite, etc.	Andendiorite, etc.	Porphyrite, etc.	Diorite, etc.	Diabase, etc.	—

be considered "historical", what then should one regard as "anti-historical"? It is perfectly clear, that in calling the cap lava - andesite, that of the middle section - chloritized andesite, and the bottom lava-silicified and albitized andesite, we would have related the description to the geologic environment, and would have, thereby, made it historical in the precise meaning of the word.

It is clear, that in systematizing effusives, one should take into account the characteristics of their composition as related to the post-volcanic alterations, but it is no less clear that in relating the system only to these features, we are distorting the history of a natural phenomenon. A heart, even if affected by sclerosis, is still called a heart so long as it does not turn to ashes. Consequently, only the total disappearance of the initial characteristics justifies a change in designation. In the given case, it is obvious that only a transformation of the effusive into a metamorphic rock may constitute such a boundary. The problem thus is reduced to determining the place and character of this boundary, which, evidently, is located outside of the area to which the terms "porphyry" and "porphyrite" apply.

Now, may we make a last remark with respect to V. I. Lebedinskiy's criticism. In relating the necessity of preserving a dual nomenclature to M. A. Usov's views relative to the "diagenetized" and "greenstone phases" of effusives, he writes that "this achievement of Soviet geologic thought cannot be simply written off". A reference has already been made above to the necessity of further study and consideration of the post-volcanic transformations of effusives of which M. A. Usov has written in his work (even though his views require substantial revision, as was reasonably noted by V. I. Lebedinskiy himself). As to the dual nomenclature, which, incidentally, was created long ago in Germany, one could attribute it to Soviet geologists only in consequence of an obvious error.

The next article — by S. N. Ivanov — is noteworthy for its even greater "orthodoxy". The author stands pat on the positions of descriptive petrography. He considers as totally unacceptable not only the proposed introduction

of a single nomenclature, but he also rejects the requirement advocated by V. I. Lebedinskiy for a "geologic approach to effusive rocks". S. N. Ivanov's polemic article first of all decries the proposed introduction of a single set of designations for effusive rocks as an anachronism and a step backward. In order to appreciate the paradoxical nature of such a view, one should recall that it is based on the principles formulated by F. Zirkel [16] exactly 95 years ago! It is true that S. N. Ivanov makes no mention of F. Zirkel, G. Rosenbusch, and other petrographers who laid the foundations for a nomenclatural differentiation of fresh and altered effusives. He refers only to A. N. Zavaritskiy's proposal for a modernized system of designations for effusive rocks consisting in appending the name of a corresponding paleo-typal variety to its cenotypal equivalent ("liparitic porphyry", "andesitic porphyrite", etc.). No doubt, however, this suggestion is only a compromise solution, which stressed a search for a way out of the nomenclature crisis (for there would have otherwise been no need to advance this suggestion). I have already written that this "effort to achieve the greatest possible etymological approximation of the two nomenclature systems points out the imminent necessity for their total merger (or, more precisely, for the elimination of one of them)".

It should be stressed here that A. N. Zavaritskiy himself, regardless of the compromise solution he advocated, has clearly expressed his own attitude toward the dual nomenclature for effusives. In discussing the historical reasons for its emergence, he writes [3]: "Initially, in the development of the science of geology... it was assumed that at different periods of the Earth's history... the conditions of rock formation were sharply different... Thus, at the initial development stage of petrography the geologic age was the basis for classification. But later, the Paleozoic and more recent (Tertiary and Post-Tertiary) effusive rocks were divided into two different genetic groups, regardless of their obvious affinity, and this differentiation came to an expression also in the petrographic terminology... The duality of designations... is to a certain extent a left-over from the past and essentially abandoned concepts (underscored by the author). This is the reason that we do not find this duality in the groups of rocks

identified recently, when these ideas have already become obsolete... The chief objective difference between the Paleozoic effusive rocks and the corresponding younger rocks is the degree of alteration resulting from the more recent secondary processes. If this characteristic is to be given a classification significance, then the dual nomenclature of rocks may be justified, but it should be applied depending on the general aspect of the rock, on the degree of deuteric alterations, and not on age".

Hence, the attitude of A. N. Zaritskiy, to whom S. N. Ivanov refers in support of his own views, is perfectly clear. A dual nomenclature is a left-over from an earlier stage of the development of petrography and may be partly justified only if totally dissociated from the age principle, provided the modernized designations recommended by him are duly introduced. From here there remains only one logical step to total rejection of dual nomenclature. Only a biased attitude could induce one to deny the fact that the term "altered (or paleotpal) liparite" better conveys an idea of the basic properties of this effusive than does the less definite and ambiguous² designation: "liparitic porphyry".

Nonetheless, S. N. Ivanov continues to insist that a rejection of dual nomenclature is a dangerous simplification, which will unavoidably lead petrography to impoverishment and regression. The existence of the science without the terms of "porphyry" and "porphyrite" appears impossible to him. He even considers enriching it by new designations: quartzephyre, ap quartzephyre, etc.

One should, in the first place, re-emphasize the fact that a single nomenclature is not poorer, but richer in terminology than a nomenclature system corresponding to special designations for altered effusive rocks. Furthermore, rejection of certain obsolete concepts is a prerequisite for any progress. The Russian orthography in discarding the letters ъ, ѓ, і, ѣ, and ѱ will formally become "poorer" through loss of five letters, but this "simplification" turned out to be an indispensable element in the development of our language. Of course, far from all were satisfied with the reform. In this case too, there appeared to be some defenders of the old traditions, who believed that without the letter ъ, the Russian people

would never survive. Prof. A. I. Tomson [11] wrote in this connection: "I have already shown that the elimination of the letters ъ, ѓ, і, ѣ, and ѱ will directly complicate and disorganize the harmonious system of Russian declensions and conjugations. With the abolishment of these letters the difference would disappear between such words as "сведение" and "свѣдѣніе", "ѣсть" and "есть", "міръ" and "миръ", etc. These changes in orthography threaten to dislocate the established mechanism of thought and weaken mental activity among the literate". A similar situation engendered in both professors a similar reaction!

S. N. Ivanov's article shows, however, that this opposition is directed not only against the single nomenclature, but also against the very principle serving as the basis for systematization and classification of effusive rocks. S. N. Ivanov in this case holds fast to the rule whose origin may be traced back to the epoch when microscopic physiography was the leading science. He writes: "It seems to me that preference should be given to those designations of rocks which are determined not on the basis of some abstract concept but by actually existing and unquestioned petrographic characteristics observable in every fragment or, at least, in every natural bedrock outcrop".

This quotation explains the true reason for the difference in opinions. It is important to emphasize right away and most decisively that this approach to rocks, which in its time was known as "fragment petrography", has, in our day become largely obsolescent. The rapid development of geology and petrography has upset many traditional ideas of the past era. It has, first of all, brought to the foreground genetic problems of effusive rocks which turned out to be far more complicated than they appeared to petrographer-microscopists. The ever emerging new proofs of the major role played by the phenomena of heterogenesis and convergence of features seriously complicate the task of rock determination "in fragment" and even in "bedrock outcrop". Not so isolated are the cases when a good knowledge of the geologic circumstances in the entire investigated area is required for a correct determination of the origin, history, and the name of a rock "fragment". Without this knowledge it is difficult, and impossible at times, to distinguish granite from granitized sedimentary rock, altered basalt from altered liparite, ignimbrite from lava, and so on. Even so distinguished a scientist as Ye. S. Fedorov has made a major error when, guided by principles of "fragment petrography", he described the metamorphic rocks, "kelabekite" and "drusite", as extrusive.

Thus, a geologic-genetic approach to rocks

²Let it be recalled that the word *porphyros* originally means "red", and *porphyros lithos* means red stone. Hence, such word formations as, for example, "greenstone porphyrite", "grey porphyrite", "black porphyrite" and so on, appear to be obviously unreasonable. This fact escapes our attention only because of our continuous use of foreign words, the true meaning of which we tend to forget.

constitutes that important element without which correct understanding of the nature of rock-forming processes and a correspondingly correct determination of the rock sometimes prove impossible. Moreover, it is obvious that the petrographer must designate his rocks, certainly not on the basis of "abstract concepts" as implied by S. N. Ivanov, but on the basis of a study of the specific geologic environment which serves as no less a "realistic characteristic", than the mineralogical composition determined under the microscope.

As F. Yu. Levinson-Lessing once stated very appropriately: "the decisive word in petrogenetic problems always belongs to geology" [9].

From the aforesaid follows then the answer to the "baffling" question posed by S. N. Ivanov: what should one call a strongly altered basic effusive rock, the primary nature of which cannot be precisely determined? Of course, the inquirer believes that "porphyrite" is the sole acceptable name. But is it not clear that by giving an unidentified phenomenon an obscure name one can hardly claim to have defined it! Would it not be more scientifically correct to designate such a rock as "altered basic effusive", or even simply "altered effusive"? One should add here that if the petrographer does not entirely rely on the omnipotence of the microscope, and studies the rocks in the field, approaching them not as "fragments of stones", but as geologic objects, such "difficult" cases will actually prove to be not too numerous.

In order to exhaust the list of questions posed by S. N. Ivanov, it remains to refer to the problem of the nomenclature of effusive rocks in geologic mapping. S. N. Ivanov writes: "I would like to hear Ye. K. Ustinov's explanation as to the precise nature of the confusion arising from the classification of rocks of volcanic origin into cenotypal and paleotypal. Why, for example, should a geologic map showing andesites and andesitic porphyrites become more legible and valuable if, on the basis of the preserved remnants of primary features, we referred to andesitic porphyrites simply as andesites?"

First of all, it is obviously wrong to call altered andesites "simply andesites" and no one actually suggests this. It is better to call them "altered andesites" (chloritized, carbonatized, sulfidized, etc.). This corresponds more closely to the nature of the rock and conveys more accurately the meaning of the term. Moreover, one should not speak only of andesites, but also of many other rocks, for which no designations are available in the scale of paleotypal varieties and which, therefore, are never identified in geologic sur-

veys in the areas where ancient volcanic formations occur. Is it possible to conclude, therefore, that in the Mesozoic or Paleozoic there was no extrusion of theolites, phonolites, trachyandesites, and so on, within the confines of the folded zones and platforms of the Soviet Union? Certainly not, but all of these effusives are comfortably camouflaged today under the all-leveling designation of "porphyrites", so convenient in its indefiniteness. If one is to add here that concealed under the mask of albitophyres and keratophyres one often finds strongly altered effusive rocks by no means of acidic, but rather of average, or even basic composition, then the picture of the serious distortion of the historical development of magmatic phenomena in the process of geologic mapping becomes evident. It seems hardly necessary to argue that the problem of the accurate reconstruction of the evolution of volcanism and the volcanic rock compositions also has practical importance in addition to its scientific significance.

In protesting the proposal for a single nomenclature of effusive rocks, S. N. Ivanov refers to his experience gained during work in the Ural areas where there are widespread occurrences of paleotypal effusives. It is important to note that support for this suggestion comes from petrographers who were also engaged in mapping the older volcanic complexes in the Urals, the Caucasus, the Altay, and the Far East. Thus, S. N. Ivanov's experience is in direct opposition with the experiences of T. V. Dianova, N. D. Zlenko, M. A. Tarkhova, T. Ya. Goncharova, and I. M. Speranskaya.

T. Ya. Goncharova writes, in particular: "Despite a prolonged study of the old, primarily Paleozoic, strongly altered volcanic formations containing chalcoppyritic mineralization in the Caucasus, the Altay, and partly in the Urals, we failed to come to a conclusion, like S. N. Ivanov, concerning the great confusion which would ensue should the paleotypal rock designations in the geological maps of regions containing predominantly effusive formations be replaced by cenotypal designations as suggested by Ye. K. Ustiyev. On the contrary, we believe that the maps can only gain thereby and be of greater value, provided of course, that definite conditions are observed, that these changes are not entirely left to the "discretion of the authors of individual maps" (S. N. Ivanov) and that they are made after a detailed and considered petrographic study of the effusive rocks".

Finally, still one more and last detail. S. N. Ivanov has several times underlined the negative features of a single nomenclature of effusives, calling it "Anglo-American", and opposed to it, the advantages of a dual nomenclature, to which he refers as the "domestic" system. I have written quite plainly that a single nomenclature for effusives is used by many geologists and

petrographers of Europe and the "countries of the Pacific belt", which, as is well known also includes China and Japan, Indonesia and Vietnam, and scores of other nations which, by no stretch of imagination, could be called "Anglo-American". As to the epithet "domestic" with respect to the dual nomenclature, it might have been considered appropriate had S. N. Ivanov been a native of Leipzig or Heidelberg, where F. Zirkel and G. Rosenbusch, the main architects of dual nomenclature principles, actually lived and worked.

In conclusion, it is well to mention the definite good that may have resulted from this discussion. First of all, it has permitted the supporters of a single nomenclature for effusives to delineate clearly their positions and to reveal the weak points of its opponents. Furthermore, it has clearly evidenced the growing dissatisfaction among petrographers with the dual nomenclature system, which has begun to threaten the progress of field and theoretical petrography. One would think that the argumentation of the defenders of the dual nomenclature contributes more than anything else to its gradual disappearance. Its hundredth anniversary which is due within a short time will, apparently, also be its last.

REFERENCES

1. Goncharova, T. Ya., K voprosu o nomenklature effuzivnykh gornyx porod. [THE PROBLEM OF THE NOMENCLATURE OF EFFUSIVE ROCKS]: Izv. Akad. Nauk SSSR, ser. geol., No. 9, 1961.
2. Dianova, T. V., Nekotoryye zamechaniya k voprosu o nomenklature effuzivnykh gornyx porod. [A FEW REMARKS ON THE PROBLEM OF THE NOMENCLATURE OF EFFUSIVE ROCKS]: Izv. Akad. Nauk SSSR, ser. geol., No. 1, 1961.
3. Zavaritskiy, A. N., Izverzhenyye gornyye porody. [IGNEOUS ROCKS]: Izd-vo Akad. Nauk SSSR, 1955.
4. Zlenko, N. D., and M. A. Tarkhova, K voprosu o yedinoi nomenklature effuzivnykh i zhil'nykh gornyx porod. [THE PROBLEM OF A SINGLE NOMENCLATURE FOR EFFUSIVE AND VEIN ROCKS]: Izv. Akad. Nauk SSSR, ser. geol., No. 1, 1961.
5. Ivanov, S. N., K voprosu o nomenklature effuzivnykh gornyx porod. [THE PROBLEM OF THE NOMENCLATURE OF EFFUSIVE ROCKS]: Izv. Akad. Nauk SSSR, ser. geol., No. 7, 1960.
6. Lebedinskiy, V. I., Zamechaniya k stat'ye Ye. K. Ustiyeva "K voprosu o nomenklature effuzivnykh gornyx porod". [COMMENTS ON YE. K. USTIYEV'S ARTICLE "THE PROBLEM OF THE NOMENCLATURE OF EFFUSIVE ROCKS"]: Izv. Akad. Nauk SSSR, ser. geol., No. 7, 1960.
7. Lebedinskiy, V. I., Novyye dannyye o genezise spilito-keratofirovoy formatsii gornogo Kryma. [NEW DATA ON THE GENESIS OF SPILITIC-KERATOPHYRE FORMATIONS OF THE CRIMEAN UPLAND. From the book, "PROBLEMS OF MAGMA AND THE ORIGIN OF IGNEOUS ROCKS"]: Tez. dokl. k simpoziumu, posv. 100-letiyu so dnya rozhhd. F. Yu. Levinson-Lessinga, 1961.
8. Levinson-Lessing, F. Yu., and Ye. N. D'yankonova-Salvel'yeva, Vulkanicheskaya gruppa Karadaga v Krymu. [THE KARADAG VOLCANIC GROUP IN THE CRIMEA]: Izd-vo Akad. Nauk SSSR, 1933.
9. Levinson-Lessing, F. Yu., Problemy genezisa magmatischeskikh porod i puti k yeye razresheniyu. [THE GENETIC PROBLEM OF MAGMATIC ROCKS AND WAYS FOR ITS SOLUTION]: Izd-vo Akad. Nauk SSSR, 1934.
10. Speranskaya, I. M., K voprosu o nomenklature effuzivnykh gornyx porod. [THE PROBLEM OF THE NOMENCLATURE OF EFFUSIVE ROCKS]: Izv. Akad. Nauk SSSR, ser. geol., No. 9, 1961.
11. Tomson, A. I., Neobkhodima reforma ne pravopisaniya, a prepodavaniya pravopisaniya. [A REFORM IS REQUIRED OF ORTHOGRAPHY TEACHING METHODS, NOT OF ORTHOGRAPHY ITSELF]: Odessa, 1905.
12. Usov, M. A., Fazy effuzivov. [EFFUSIVE PHASES]: Tomsk, 1929.
13. Usov, M. A., Fatsii i fazy intruzivov. [THE FACIES AND PHASES OF INTRUSIVE ROCKS]: 2-ye izd., Tomsk, 1932.
14. Ustiyev, Ye. K., K voprosu o nomenklature effuzivnykh gornyx porod. [THE PROBLEM OF THE NOMENCLATURE OF EFFUSIVE ROCKS]: Izv. Akad. Nauk SSSR, ser. geol., No. 11, 1959.
15. Ustiyev, Ye. K., K voprosu o nomenklature effuzivnykh gornyx porod.

[THE PROBLEM OF THE NOMENCLATURE OF EFFUSIVE ROCKS]: B sb.
 "Problemy vulkanizma", Yerevan, 1959.

16. Zirkel, F., Lehrbuch der Petrographie, Bonn, 1866.

Received March 7, 1961

ON A BOOK BY G.M. ZARIDZE AND N.F. TATRISHVILI "MAGMATISM IN GEORGIA AND ASSOCIATED ORE FORMATIONS"

by
 A. P. Lebedev

The book being reviewed may, to a certain extent, be considered an expansion of the composite, generalized studies on magmatism in Georgia published earlier by the same authors, or by one of them (G. M. Zaridze) [1, 2]. A characteristic mainly of the young effusive and intrusive magmatism in Georgia was given in the first study [1]. The second sheds some light on the magmatic formations of predominantly Paleozoic age [2]. The new summary covers both the first and the second groups of rocks.

There is a dual purpose in compiling extensive review studies on the petrography of individual major provinces such as the Georgian SSR. On the one hand, the object is to provide reference material where the reader can find the required information on all formations and rocks occurring in this region, along with the newest data on its petrography and geology. On the other hand, a general survey of materials on different forms of magmatic manifestations in a definite major province enables the authors of such a survey to make a certain critical analysis of the material from the standpoint of this or that petrological concept. It is deemed useful, therefore, to review this work with due consideration of these positions.

In the descriptive part the material in this study may be subdivided into two sections: 1) the old (Pre-Mesozoic) formations, and 2) the recent formations. Far more attention is devoted to the former, about three-quarters of the descriptive part in the book being devoted to them. Here one finds a description of rocks occurring in the Dzirul'skiy, Khram'skiy, and other mountain massifs. These descriptions provide a detailed description of

separate bodies and their sections, of individual types of rocks, their contacts and hydrothermal products, etc. The description is very detailed, sometimes excessively burdened with unnecessary particulars. One of the shortcomings of this section (as well as of the entire descriptive part) is the isolation of the recorded chemical analyses and quantitative estimates from the specific descriptions of the rocks. All of these analyses are given without "an address", without precise reference to locations, often in a generalized form. This, of course, considerably reduces their value to the reader for purposes of comparison, or as reference material. The absence of schematic charts showing distribution, sampling locations, etc., also must be considered as a defect. All this results occasionally in too much generalization and isolation of the descriptions from the actual environment.

The second group covers Meso-Cenozoic formations. It includes the "neointrusions", well known in the petrographic literature on the Caucasus, of acidic and average rocks, the Jurassic volcanic rocks of Racha, Southern Osetiya, Khevsuretiya, Kakhetiya, and Southern Georgia, the complex of young effusive formations of the Kazbek, and other volcanic centers. This group of formations was described in G. M. Zaridze's summary of 1947 [1] referred to above. In contrast to this summary, the descriptions given here are in an extremely concise, often almost synoptical, form. A characteristic feature of this part of the book, which, incidentally, distinguishes it from the first ("pre-Mesozoic") part, is the lack of uniformity in the depiction of individual areas, imparting to the entire section and eclectic character and leaving an impression of incompleteness. The selection of individual sections, bodies, rocks, and entire formations for more detailed treatment appears to be made at random. Some of the descriptions are overly detailed, others are presented synoptically, and a few features are overlooked altogether. In arranging and compiling this

¹ O knige G. M. Zaridze i N. F. Tatrishvili "Magmatizm gruzii i svyazannyye s nim rudooobrazovaniya", pp. 114 - 116.

section the authors, we feel, should have followed the course of a more profound and generalized characterization of individual groups or formations, bearing in mind that more detailed factual material has already been presented with adequate completeness in the well-known surveys of Georgia prepared by G. S. Dzotsenidze, N. P. Skhirtladze, D. S. Belyankin, and V. P. Petrov. In this manner, generalized characteristics would have made it possible to throw, in a more systematic way, a more even light on this rich and diversified material.

The positive features, which favorably distinguish these descriptions from G. M. Zaridze's work published in 1947, is the method of region-by-region treatment of materials adopted in this study which facilitates its general use.

In summing up the characteristics of the book's descriptive part (Chapters 3 to 5), one should say that on the whole, in spite of the mentioned shortcomings, it still gives a fairly complete idea of the principal magmatic and metamorphic complexes in Georgia. In a new edition of this book is prepared, the authors would do well to eliminate the mentioned shortcomings (absence of schematic charts and adequate records on chemical analyses, etc.).

In the conclusive chapters of the book, which present an independent interest, the authors attempt to analyse the genetic problems of certain magmatic formations in Georgia and their metallogeny, from a definite viewpoint.

Basically, the petrogenetic deductions of the authors are related to the granitoids in the old and recent formations and to acidic effusives. They relate the origin of all intrusive acidic formations to the processes of silica, potassium, and sodium replacement. In their opinion, the granitoids appeared as a result of the transformation of the old country rocks (metamorphic and sedimentary) and the old basic rocks: gabbro and amphibolites (in the Dzirul'skiy massif). The acidic effusives, on the other hand, originated from basic magma contaminated by country-rock materials. The authors, apparently, totally discount the existence of intrusive or plutonic acidic magma. This point of view stands, thus, in contradiction to prevalent ideas on the genesis of the Caucasian granitoids.

What factual material then leads the authors to such conclusions? On the whole, this is material accumulated by observations carried out in the zones of mineralization and microclinization in the deeply eroded massifs constituting the remnants of the old crystalline basement (Dzirul'skiy mountain massif, the zone of crystalline schists in the

Great Caucasus). It is true that in these regions one can observe pretty clearly the workings of the processes of metasomatic replacement of gneisses and schists by quartz and feldspathic injections accompanied by the formation of porphyroblastic microcline, petritization, albitization, silification, and other phenomena. Although some of these manifestations may, possibly, be associated with other processes (in particular, with metamorphic differentiation), nevertheless, it is really difficult to deny the great role played here by metasomatic replacement, and the authors are perfectly correct in calling the attention of petrographers to these phenomena. One can also agree with a few interesting considerations formulated in the same chapter with respect to the genesis of the accessory minerals of certain types of granitoids. The role of metasomatic processes in their formation is reflected very clearly in some places (apatite, sphene, and less distinctly with regard to zircon and orthite). Indications of metasomatic growth of the accessory minerals can, indeed, be traced also in some basic rocks, for example, in the Siberian traps.

Thus, the authors are perfectly right when they put strong emphasis on the great petrogenetic significance of alkaline and siliceous metasomatic processes in the described cases. In drawing attention to this aspect of granitoid petrogenesis in Georgia, and in providing a number of conclusive examples to this effect, the authors, no doubt, have rendered considerable service.

While recognizing along with them the great importance of the processes of metasomatic granitization in the formation of certain types of old granitoids in Georgia, one still cannot refrain from saying that the theory advanced by them that all granitoid intrusions in the Caucasus were formed exclusively in this manner, seems to be hardly acceptable, particularly when this concerns Post Jurassic intrusions. This theory fails to provide a satisfactory explanation for a considerable number of facts related to specific intrusions. Here are a few of these facts, 1) the obvious signs of high-temperature contact action produced by acid intrusions on the country rocks and followed by the formation of high-temperature hornfels (for example, in Teplinsk, Kazbek, and other neointrusions); 2) the shape of the intrusive bodies and the peculiarities of their internal structure point to the autonomous character of their development and to the phenomena of magmatic flow; 3) the sharply discordant, cross-cutting, positioning of the individual intrusive bodies with respect to the enclosing rocks; 4) a tendency toward a more uniform composition observable in major intrusions, and assimilation of xenoliths, 5) the petrochemical and (in a broader sense) the general geochemical characteristics of

individual intrusions and intrusive complexes (as described, for instance, in G.D. Afanasyev's works on the Greater Caucasus); 6) the multi-mineral composition of granitoids, which, according to D.S. Korzhinskiy, is at variance with their metasomatic genesis (the explanations furnished by the authors in this respect do not sufficiently clarify the question).

To accept the theory of metasomatic granitization on so large a scale as suggested by the authors, i. e. for the explanation of the formation of all granitoid intrusions (and not only for the rocks occurring in the old magmatic and injection zones), one must find answers to many new questions. First of all, that concerning the reasons for the selective character of metasomatism, which, in some cases, is of potassic, and in other cases of sodic, or combined type. Are the relations between them determined by tectonic factors, by the depth of intrusive body formation, by temperature, the composition of wall rocks, or by some factors — all these problems remain essentially unsolved by the authors. Yet, without solving them, it is impossible to answer the basic problems of petrology and metallogeny in the area under review. One should also note the lack of clarity in the authors' views on the nature of the granitizing emanations themselves. In certain instances they seem to approximate closely to the views held by V.S. Koptev-Dvornikov relative to the existence of "alaskitic" or "leucocratic" magma (p. 201), but in other cases they speak of "solutions consisting of silicium, potassium, and sodium" (p. 214).

In analyzing the problems concerning the relationship between endogenic mineralization and magmatism (chapter 7) the authors, along with the theory of the plutonic origin of ore elements, also recognize the role of processes causing the extraction of ore elements from the country rocks, and their subsequent concentration. It is possible to agree that this method of ore-substance accumulation (still insufficiently studied on specific examples) actually may have had considerable significance in the metallogeny of the Caucasus.

In considering the petrogenetic part of the book as a whole, we are inclined to acknowledge that the principal merit of the authors is that they were, possibly, the first authors dealing with the Caucasus to underline the great importance of the metasomatic processes in the genesis of the ancient-complex granitoids. However, the role of these processes in the genesis of acid intrusions in the upper structural stage appears to us to be exaggerated because of obvious under-estimation and insufficient appreciation of the processes unquestionably magmatic in character.

On the whole the reviewed book, no doubt, will contribute to a further expansion of our ideas of magmatism in Georgia. As to the original theoretical considerations presented in the book, they, in spite of their debatable nature, will stimulate a more profound study of the problem relative to the formation of granites not only in the Caucasus, but also in other regions.

REFERENCES

1. Zaridze, G.M., *Zakonomernosti razvitiya vulkanizma v Gruzii i svyazannyye s nim rudoobrazovaniya*. [THE DEVELOPMENT REGULARITIES OF VOLCANISM IN GEORGIA AND THE ASSOCIATED MINERALIZATIONS]: Gostekhizdat, GruzSSR, 1947.
2. Zaridze, G.M., and N.F. Tatrishvili, *Vvedeniye v magmaticheskuyu geologiyu Gruzii*. [INTRODUCTION TO THE MAGMATIC GEOLOGY OF GEORGIA]: Gostekhizdat, GruzSSR, 1947.

Received September 30, 1960

BIBLIOGRAPHY

GEOLOGIC LITERATURE RECEIVED BY THE
LIBRARY OF THE SECTION OF GEOLOGIC
AND GEOGRAPHIC SCIENCES, THE U.S.S.R.
ACADEMY OF SCIENCES, IN JUNE, 1961

A. Articles in Periodicals

GEOLOGY

1. Alekseyeva, K. A., and V. V. Kravets, Sravnitel'nyye dannyye opredeleniya uprugikh svoystv kamennykh meteoritov i nekotorykh gornyykh porod. [COMPARATIVE DATA ON THE DETERMINATION OF THE ELASTIC PROPERTIES OF STONY METEORITES AND CERTAIN ROCKS]: Geol. zh. Akad. Nauk UkrSSR, t. 21, vip. 2.
2. Aliyev, A. G., and V. P. Akayeva, O monografii A. G. Babayeva i A. M. Akramkhodzhayeva "Paleogeografiya neftegazonosnykh melovykh otlozheniy Uzbekistana". [ON THE MONOGRAPH BY A. G. BABAYEV AND A. M. AKRAMKHODZHAYEVA PALEOGEOGRAPHY OF THE OIL- AND GAS-BEARING CRETACEOUS DEPOSITS IN UZBEKISTAN]: Uzb. geol. zh., No. 1.
3. Alimov, K. A., Sporovo-pyl'tsevoy analiz pri raschlenenii i korrelyatsii yurskikh otlozheniy Ferganskogo khrebta. [SPORE AND POLLEN ANALYSIS IN THE DIFFERENTIATION AND CORRELATION OF THE JURASSIC DEPOSITS IN THE FERGANA RANGE]: Uzb. geol. zh., No. 1.
4. Al'ter, S. P., K stratigrafii chetvertichnykh otlozheniy Priyeniseyskoy zony. [STRATIGRAPHY OF THE QUATERNARY DEPOSITS IN THE YENISEY RIVER AREA]: Inform. sb. VSEGEI, No. 40, 1960.
5. Appolonov, M. K., and G. Kh. Yergaliyev, O stratigraficheskom polozenii yerkebidaikskoy svity Vostochno-Kokchetavskogo progiba. [STRATIGRAPHIC POSITION OF THE YERKEBIDAIIAN SUITE IN THE EAST-KOKCHETAVSKIY DOWNWARP]: Izv. Akad. Nauk KazakhSSR, ser. geol., vyp. 2.
6. Babayan, G. D., and A. K. Bobrov, Struktura Vilyuyskoy sineklizy i Predverkhoyanskogo progiba po krovlе yurskikh otlozheniy. [STRUCTURE OF THE VILYUY SYNECLISE AND THE CISVER-KHOYANSK TROUGH ALONG THE DOME OF THE JURASSIC DEPOSITS]: Geol. i geofiz., Sib. otd. Akad. Nauk SSSR, No. 3.
7. Belyayevskiy, N. A., Vazhnyye problemy geologii. (Na mezhdunarodnom kongresse v Skandinavii). [IMPORTANT PROBLEMS OF GEOLOGY. (AT THE INTERNATIONAL CONGRESS IN SCANDINAVIA)]: Priroda, No. 4.
8. Birina, L. M., O proiskhozhdenii krasnotsvetnoy okraski terrigennykh porod devona i kembriya v severnoy chasti Moskovskoy sineklizy. [ORIGIN OF THE REDDISH COLORATION OF TERRIGENOUS DEVONIAN AND CAMBRIAN ROCKS IN THE NORTHERN PART OF THE MOSCOW SYNECLISE]: Byul. MOIP, otd. geol., t. 36, vyp. 2.
9. Bobkova, N. N., O mestnoy zone Caprinula soluni v verkhnem senomane Yugo-Zapadnogo Darvaza. [LOCAL ZONE OF CAPRINULA SOLUNI IN THE UPPER SENOMAN IN SOUTHWESTERN DARVAZ]: Inform. sb. VSEGEI, No. 42.
10. Bordunov, I. N., Tektogenez i vulkanizm kak glavnyye faktory osadkoobrazovaniya proterozoyskikh zhelezisto-kremnistyykh formatsiy Kremenchugskogo zhelezorudnogo rayona. [TECTONOGENESIS AND VOLCANISM AS THE MAIN FACTORS OF SEDIMENTATION OF THE PROTEROZOIC FERRO-SILICEOUS FORMATIONS IN THE KREMENCHUG IRON-ORE DISTRICT]: Geol. zh. Akad. Nauk UkrSSR, t. 21, vip. 2.

11. Borisyak, M.A., O.P. Kovalevskiy, and T.V. Nikolayeva, K stratigrafii silura khrebta Chingiz. [SILURIAN STRATIGRAPHY IN THE CHINGHIZ RANGE]: Inform. sb. VSEGEI, No. 42.
12. Boyakova, V.D., and V.P. Vladimirovich, Stratigrafiya severnoy chasti Chelyabinskogo burougol'nogo basseyna. [STRATIGRAPHY OF THE NORTHERN PART OF THE CHELYABINSK BROWN-COAL BASIN]: Inform. sb. VSEGEI, No. 42.
13. Bukatchuk, P.D., Spornyye voprosy stratigrafii rifeyskikh otlozheniy Moldavii. [MISCELLANEOUS PROBLEMS ON THE STRATIGRAPHY OF THE RIPHEAN DEPOSITS IN MOLDAVIA]: Geol. zh. Akad. Nauk UkrSSR, t. 21, vip. 2.
14. Vasil'yev, Yu.M., Antropogennyye otlozheniya Yuzhnogo Zavolzh'ya. [THE POST-QUATERNARY DEPOSITS OF THE SOUTHERN VOLGA AREA]: Byul. MOIP, otd. geol., t. 36, vyp. 2.
15. Vasil'yev, Yu.M., and V.S. Mil'nichuk, Strukturnyy plan neogenovykh otlozheniy Ustyurta. [STRUCTURAL PLAN OF NEOGENE DEPOSITS IN USTYURT]: Novosti neft. i gazovoy tekhniki, geologiya, No. 3.
16. Vvedenskaya, A.V., Izucheniye napryazheniy i razryvov v ochagakh zemletryaseniya. [STUDY OF THE STRESSES AND FISSIONS IN EARTHQUAKE FOCI]: Byul. MOIP, otd. geol., t. 36, vyp. 2.
17. Vinukurova, Ye.G., and Ye.A. Zhukova, K stratigrafii melovykh otlozheniy nizov'yev Amu-Dar'ya po skvazhinam v rayonakh Khodzheyli i Chimbaya. [STRATIGRAPHY OF THE CRETACEOUS DEPOSITS IN THE LOWER-REACHES OF THE AMU-DAR'YA BASED ON BORE-HOLE LOGS FROM THE VICINITY OF KHODZHEYLI AND CHIMBAY]: Uzb. geol. zh., No. 2.
18. Vozhzhova, N.N., O metodike raschleneniya geologicheskogo razreza tretichnoy tolshchi Yugo-Zapadnoy Turkmenii po parametru β . [METHOD OF GEOLOGIC SECTION DIFFERENTIATION IN THE TERTIARY FORMATION OF SOUTH-WEST TURKMENIA BASED ON PARAMETER β]: Izv. Akad. Nauk TurkmSSR, ser. fiz.-tekh., khim. i geol. nauk, No. 1.
19. Vorob'yeva, O.A., O rabote petrograficheskoy sekti XXI sessii Mezhdunarodnogo geologicheskogo kongressa i vpechatleniyakh ob ekskursii po Shvetsii. [WORK OF THE PETROGRAPHIC SECTION AT THE 21ST SESSION OF THE INTERNATIONAL GEOLOGICAL CONGRESS AND IMPRESSIONS ON THE EXCURSION IN SWEDEN]: Byul. MOIP, otd. geol., t. 36, vyp. 2.
20. Voronov, P.S., Opyt restavratsii lednikovogo shchita Antarktidy epokhi maksimal'nogo oledeneniya Zemli. [EXPERIMENT IN RESTORING THE ANTARCTIC GLACIER SHIELD DURING THE EPOCH OF MAXIMUM GLACIATION OF THE EARTH]: Inform. byul. Sov. Antarkt. eksp., No. 23, 1960.
21. Voronov, P.S., Skhema noveyshey tektoniki Antarktidy. [SKETCH OF THE MOST RECENT TECTONICS OF THE ANTARCTIC]: Inform. byul. Sov. Antarkt. eksp., No. 24, 1960.
22. Gerasimov, I.P., Sovremennyye laterity i lateritnyye pochvy. [MODERN LATERITES AND LATERITIC SOILS]: Izv. Akad. Nauk SSSR, ser. geogr., No. 2.
23. Glazunov, O.M., Lysanskiy gabbropirokseno-serpentinitovyy kompleks v zapadnoy chasti Vostochnogo Sayana. [THE LYSANSK GABBRO-PYROXENE-SERPENTINITE COMPLEX IN THE WESTERN PART OF EASTERN SAYAN]: Geol. i geofiz., Sib. otd. Akad. Nauk SSSR, No. 3.
24. Golenetskiy, S.I., Opredeleyeniye moshchnosti zemnoy kory po nablyudeniym voln, otrazhennykh ot yeye podoshvy i glubiny ochagov aftershokov Srednebaykal'skogo zemletryaseniya 29 avgusta 1959 g. [DETERMINATION OF THE THICKNESS OF THE EARTH'S CRUST BASED ON THE OBSERVATION OF WAVES REFLECTED FROM ITS SURFACE AND ON THE DEPTH OF THE SEISMIC AFTER-SHOCK FOCI OF THE CENTRAL BAYKAL EARTHQUAKE OF AUGUST 29, 1959]: Geol. i geofiz., Sib. otd. Akad. Nauk SSSR, No. 2.
25. Gol'man, Ye.I., O nakhodke kamennougol'nykh otlozheniy v tsentral'noy chasti Vostochnogo Sayana. [DISCOVERY OF COAL MEASURES IN THE CENTRAL PART OF EASTERN SAYAN]: Inform. sb. VSEGEI, No. 40, 1960.
26. Gol'ts, S.I., Chetvertichnaya istoriya doliny r. Klyaz'my (v predelakh Meshchery). [THE QUATERNARY HISTORY OF THE KLYAZ'MA RIVER VALLEY (IN THE MESHCHERA REGION)]: Byul. MOIP, otd. geol., t. 36, vyp. 2.

7. Goretskaya, Ye.N., Magmaticheskiiye formatsii Tyan'-Shanya. [THE MAGMATIC FORMATIONS OF THE TIEN'-SHAN']: Zap. Vses. min. o-va, ch. 90, vyp. 2.
8. Greshner, S.G., O rasprostraneni effuzivov kislogo sostava v stratigraficheskoy razreze silura Zapadnoy zelenokamennoy zony Mugodzhara. [DISTRIBUTION OF ACID EFFUSIVES IN THE SILURIAN STRATIGRAPHIC COLUMN IN THE WESTERN GREENSTONE ZONE OF MUGODZHAR]: Vestn. Akad. Nauk KazakhSSR, No. 4.
9. Grigor'yev, K.A., O bazal'nykh konglomeratakh nizhnokamennougol'noy krasnotsvetnoy svity severnogo sklona khr. Moldotau (Severnaya Kirgiziya). [BASALTIC CONGLOMERATES OF THE LOWER CARBONIFEROUS RED BEDS ON THE NORTHERN SLOPE OF THE MOLDOTAU RANGE (NORTHERN KIRGHIZIYA)]: Inform. sb. VSEGEI, No. 41, 1960.
10. Gridnev, N.I., and K.A. Nabyev, O litologii chetvertichnykh nadvodno-del'tovykh otlozheniy Amu-Dar'ya. [LITHOLOGY OF THE QUATERNARY SUPRA-AQUEOUS DELTAIC DEPOSITS OF THE AMU-DAR'YA RIVER]: Uzb. geol. zh., No. 1.
11. Grosval'd, M.G., O proiskhozhdenii tak nazyvayemykh "ziyayushchikh tektonicheskikh treshchin" v Sayanakh i Pribaykal'ye. [ORIGIN OF THE SO-CALLED "GAPING TECTONIC FRACTURES" IN THE SAYAN MOUNTAINS AND BAYKAL REGION]: Byul. MOIP, otd. geol., t. 36, vyp. 2.
12. Gur'yanova, V.N., and A.L. Dodin, Novyye dannyye po geologii Udskoy i Vostochnoy Sayana. [NEW DATA ON THE GEOLOGY OF THE UDSKO-IIYSKIY REGION IN EASTERN SAYAN]: Inform. sb. VSEGEI, No. 40, 1960.
13. Davlyatov, Sh.D., Ob al'piyskoy tektonike zapadnykh okonchaniy Zirabulak-Ziaetdinskikh antiklinal'nykh podnyatiy i ikh svyazi s Kaganskoy grupпой polozhitel'nykh skladok. [ALPINE TECTONICS OF THE WESTERN FRINGES OF THE ZIRABULAK-ZIAETDINSK ANTICLINAL UPLIFTS AND THEIR RELATION TO THE KAGANSK GROUP OF POSITIVE FOLDS]: Uzb. geol. zh., No. 2.
14. Dadashev, F.G., Sopostavleniye razrezov produktivnoy tolschi perspektivnykh ploshchadey Yugo-Vostochnogo Kobystany. [COMPARISON OF THE PRODUCTIVE SERIES SECTIONS IN THE POTENTIAL AREAS OF SOUTHEASTERN KOBYSTAN]: Izv. Akad. Nauk AzerbSSR, ser. geol.-geogr. nauk i nefti, No. 1.
15. Dement'skaya, R.M., Glavnyye cherty stroyeniya zemnoy kory v Antarktike. [PRINCIPAL STRUCTURAL FEATURES OF THE EARTH'S CRUST IN THE ANTARCTIC]: Inform. byul. Sov. Antarkt. eksp., No. 23, 1960.
16. Dobrovolskiy, V.V., Okamenelyye sledy drevnykh landshtaftov. [PETRIFIED TRACES OF ANCIENT LANDSCAPES]: Priroda, No. 4.
17. Dobryanskiy, A.M., K metodike opredeleniya amplitud nebol'shikh razryvnykh naru-sheniy. [METHODS OF DETERMINING MINOR FAULT AMPLITUDES]: Geol. zh. Akad. Nauk UkrSSR, t. 21, vip. 2.
18. Drumya, A.V., et al., Novyye kapital'nyy trud po geologii Ukrainy i Moldavii (Atlas paleogeograficheskikh kart Ukrainy i Moldavskoy SSR). [A NEW MAJOR WORK ON THE GEOLOGY OF THE UKRAINE AND MOLDAVIA (ATLAS OF PALEO-GEOGRAPHIC MAPS OF THE UKRAINIAN AND MOLDAVIAN SSR)]: Geol. zh. Akad. Nauk UkrSSR, t. 21, vip. 2.
19. Yeziashvili, A.G., Novaya metodika detal'nykh geologo-s'yemnykh rabot. [NEW METHOD FOR DETAILED GEOLOGIC SURVEYS]: Izv. Akad. Nauk TurkmSSR, ser. fiz.-tekhn., khim. i geol. nauk, No. 1.
20. Zavarzin, L.G., and M.N. Blagoveshchenskaya, Ordovikiye otlozheniya v bassejne verkhnego techeniya r. Kamenki i nekotoryye osobennosti geologicheskogo stroyeniya etogo rayona. [THE ORDOVICIAN DEPOSITS IN THE BASIN OF THE UPPER KAMENKA RIVER AND CERTAIN PECULIARITIES OF THIS REGION'S GEOLOGIC STRUCTURE]: Inform. sb. VSEGEI, No. 40, 1960.
21. Zamarayev, S.M., O svitakh aldanskogo yarusy otlozheniy nizhnego kembriya yuzhnoy chasti Sibirskoy platformy. [THE ALDAN SERIES OF THE LOWER CAMBRIAN DEPOSITS IN THE SOUTHERN PART OF THE SIBERIAN PLATFORM]: Geol. i geofiz., Sib. otd. Akad. Nauk SSSR, No. 2.
22. Zankina, I.Ye., K voprosu o granitnoy devone i karbone v tsentral'noy chasti Russkoy platformy. [THE PROBLEM OF THE BOUNDARY BETWEEN THE DEVONIAN AND THE CARBONIFEROUS

IN THE CENTRAL PART OF THE
RUSSIAN PLATFORM]: Inform. sb.
VSEGEI, No. 42.

43. Zernetskiy, B. F., and G. I. Nemkov, Nekotoryye zamechaniya po povodu stat'i V. G. Kulichenko "K voprosu o vozraste numulitovykh izvestnyakov yugo-zapadnoy chasti Gornogo Kryma". [A FEW REMARKS ON V. G. KULICHENKO'S ARTICLE: "THE AGE PROBLEM OF THE NUMMULITIC LIMESTONES IN THE SOUTHWESTERN PART OF THE CRIMEAN UPLAND"]: Geol. zh., Akad. Nauk UkrSSR, t. 21, vip. 2.
44. Ibragimov, Z. S., K voprosy o treshchino-
vatosti porod Yugo-Vostochnykh Kyzyl-
kumov. [THE PROBLEM ROCK JOINTS
IN THE SOUTHEASTERN KYZYLKUM
MOUNTAINS]: Uzb. geol. zh., No. 1.
45. Il'yukhina, N. P., Sredne-verkhnepaleo-
zoyskiye otlozheniya srednego techeniya
r. Angary. [THE MIDDLE AND UPPER
PALEOZOIC DEPOSITS IN THE CEN-
TRAL REACHES OF THE ANGARA
RIVER]: Inform. sb. VSEGEI, No. 40,
1960.
46. Kazakova, V. P., and G. P. Leonov, K
voprosu o stratigraficheskom polozhenii
i vozraste tak nazyvayemykh "onkoforo-
vykh sloyev" Severnykh Yergeny.
THE PROBLEM OF THE STRATIGRAPHIC
POSITION AND AGE OF THE SO-CALLED
"ONKOFORIC LAYERS" IN THE NORTH
ERGENI MOUNTAINS]: Byul. MOIP, otd.
geol., t. 36, vyp. 2.
47. Kazenkina, G. A., Veshchestvennyy sostav
donnykh otlozheniy zaliva Proval i
osobennosti osadkoobrazovaniya. [THE
ELEMENTARY COMPOSITION OF THE
BOTTOMSET BEDS IN PROVAL BAY
AND THE PECULIARITIES OF SEDI-
MENTATION]: Geol. i geofiz., Sib.
otd. Akad. Nauk SSSR, No. 3.
48. Kim, V. A., K voprosu o metamorfizme
peschanikov Dzhezkazgana. [THE
PROBLEM OF THE METAMORPHISM
OF THE DZHEZKAZGAN SANDSTONES]:
Izv. Akad. Nauk KazakhSSR, ser. geol.,
vyp. 2.
49. Kovalevskiy, O. P., O verkhneordoviskom
vozraste izvestnyakov g. Akdombak
(khr. Chingiz). [THE UPPER ORDOVI-
CIAN AGE OF MOUNT AKDOMBAK
LIMESTONES (CHINGHIZ RANGE)]:
Inform. sb. VSEGEI, No. 42.
50. Koval'skiy, V. V., K voprosu o vydelenii
nekotorykh kharakternykh tipov kimber-
litov Yakutii. [THE PROBLEM OF THE
IDENTIFICATION OF CERTAIN CHAR-
ACTERISTIC TYPES OF YAKUTIAN
KIMBERLITES]: Geol. i geofiz., Sib.
otd. Akad. Nauk SSSR, No. 2.
51. Kovrigina, Ye. K., and F. P. Kovrigin,
Stratigrafiya dokembriya zapadnogo
sklona Yeniseyskogo kryazha v basseyn-
nakh rek Verkhney i Nizhney Surnikh,
Stolobovoy i Isakovki. [STRATIGRAPHY
OF THE PRECAMBRIAN DEPOSITS ON
THE WESTERN SLOPE OF THE YENISEY
RANGE IN THE BASINS OF THE UPPER
AND LOWER SURNIKH, STOLBOVAYA,
AND ISAKOVKA RIVERS]: Inform. sb.
VSEGEI, No. 40, 1960.
52. Kogan, A. B., Postroyeniye i ispol'zovaniye
kart prevysheniya dlya vyyavleniya pogre-
bennykh platformennykh podnyatiy.
[PLOTING AND UTILIZATION OF
ELEVATION MAPS TO IDENTIFY
BURIED PLATFORM UPLIFTS]: Geol.
nefti i gaza, No. 4.
53. Kononov, Yu. S., O formirovaniy triasovykh
triasovykh struktur v Yuzhno-Emben-
skom rayone. [FORMATION OF TRIASSIC
STRUCTURES IN THE SOUTH-EMBEN
REGION]: Vestn. Akad. Nauk KazakhSSR
No. 4.
54. Kostogryz, N. Ya., O razlome, nameche-
nnom V. N. Veberom v Yuzhnoy Fer-
gane. [ON THE FAULT DESCRIBED
BY V. N. WEBER IN SOUTH FERGANA]:
Novosti neft. i gazovoy tekhniki, No. 3.
55. Kochkin, G. B., K kharakteristike
kembrizmskikh subvulkanicheskikh in-
truziy v Gornom Altaye. [CHARAC-
TERISTIC OF THE CAMBRIAN SUB-
VOLCANIC INTRUSIONS IN THE ALTAY
HIGHLANDS]: Inform. sb. VSEGEI,
No. 41, 1960.
56. Krapivner, R. B., Problema soyedineniya
basseyinov Kamy, Vychehgy i Pechory
v chetvertichnoye vremya i periglyatsial'
nyye otlozheniya basseynov r. Kamy. [THE
PROBLEM OF THE CONNECTION OF
THE KAMA, VYCHEGDA, AND PECHORA
RIVER BASINS IN THE QUATERNARY
AND THE PERIGLACIAL DEPOSITS OF
THE KAMA RIVER BASIN]: Byul. MOIP,
otd. geol., t. 36, vyp. 2.
57. Krivolutskiy, A. Ye., Geologicheskoy
vozrast Kavkazskogo khrebt. [THE
GEOLOGIC AGE OF THE CAUCASUS
MOUNTAIN RANGE]: Izv. Akad. Nauk
SSSR, ser. geogr., No. 2.
58. Kropotkin, P. N., Problema krupnykh
gorizontal'nykh dvizheniy zemnoy kory

BIBLIOGRAPHY

- v svete sovremennykh geofizicheskikh dannykh. [THE PROBLEM OF MAJOR HORIZONTAL MOVEMENTS OF THE EARTH'S CRUST IN THE LIGHT OF MODERN GEOPHYSICAL DATA]: Byul. MOIP, otd. geol., t. 36, vyp. 2.
9. Kudryavtsev, V. A., O tektonike Severo-Vostochnogo Kitaya. [THE TECTONICS OF NORTHEASTERN CHINA]: Byul. MOIP, otd. geol., t. 36, vyp. 2.
 10. Kunin, N. Ya., Tektonicheskoye stroyeniye i perspektivy neftegazonosnosti yugo-zapadnoy chasti Chu-Sarysuyskoy depressii. [THE TECTONIC PATTERN AND THE PROSPECTS FOR GAS AND OIL IN THE SOUTHWESTERN PART OF THE CHU-SARYSUYSK DEPRESSION]: Izv. Akad. Nauk KazakhSSR, ser. geol., vyp. 2.
 11. Ladiyeva, V. D., Dokembriyskiye vulkano-gennyye obrazovaniya Konkskogo rayona Ukrainskogo kristallicheskogo massiva. [PRECAMBRIAN VOLCANIC FORMATIONS IN THE KONSKIY REGION OF THE UKRAINIAN CRYSTALLINE MASSIF]: Geol. zh., Akad. Nauk UkrSSR, t. 21, vip. 2.
 12. Lange, O. K., Aleksandr Nikolayevich Mazarovich—student i molodoy uchenyy. [ALEKSANDR NIKOLAYEVICH MAZAROVICH — STUDENT AND YOUNG SCIENTIST]: Byul. MOIP, otd. geol., t. 36, vyp. 2.
 13. Larchenko, A. Ya., Miotsenovyye otlozheniya Zapadnogo Kopet-Daga. [MIOCENE DEPOSITS IN WESTERN KOPET-DAG]: Izv. Akad. Nauk TurkmSSR, ser. fiz.-tekhn., khim. i geol. nauk, No. 1.
 14. Lebedev, V. G., O printsipakh geomorfologicheskogo rayonirovaniya. [PRINCIPLES OF GEOMORPHOLOGICAL ZONING]: Vestn. MGU, ser. V, geogr., No. 2.
 15. Lyustikh, Ye. N., and A. Ya. Saltykovskiy, K voprosu ob obrazovanii granitnogo sloya zemnoy kory. [THE PROBLEM OF THE FORMATION OF THE GRANITE LAYER IN THE EARTH'S CRUST]: Geokhimiya, No. 4.
 16. Mazarovich, O. A., Formatsionnyye ryady nekotorykh mezhgornyykh vpadin kaledonid SSSR. [THE FORMATIVE SERIES OF CERTAIN INTERMONTANE TROUGHS OF THE CALEDONIDES IN THE USSR]: Byul. MOIP, otd. geol., t. 36, vyp. 2.
 17. Malovitskiy, Ya. P., and S. P. Vartanov, Novyye dannyye o seysmostratigraficheskoy razreze Yugo-Vostochnogo Kaspiya. [NEW DATA ON THE SEISMO-STRATIGRAPHIC SECTION OF THE SOUTHEASTERN CASPIAN]: Novosti nef. i gazovoy tekhniki, No. 3.
 18. Mal'kovskiy, F. S., K voprosu o granitse karbona i permi. [THE PROBLEM OF THE BOUNDARY BETWEEN THE CARBONIFEROUS AND THE PERMIAN]: Byul. MOIP, otd. geol., t. 36, vyp. 2.
 19. Mironova, L. V., O granitse bukharskikh i suzaksikh sloyev. [THE BOUNDARY BETWEEN THE BUKHARIAN AND SUZAKIAN LAYERS]: Inform. sb. VSEGEI, No. 42.
 20. MIKHALEVA, L. A., O pozdnegertsinskikh granitakh kalbinskogo tipa v Gornom Altaye. [THE LATE HERCYNIAN GRANITES OF THE KALBINIAN TYPE IN THE ALTAY HIGHLANDS]: Geol. i geofiz., Sib. otd. Akad. Nauk SSSR, No. 2.
 21. Musatov, D. I., Stratigrafiya kembriyskikh otlozheniy vostochnogo sklona Kuznetskogo Alatau. [THE STRATIGRAPHY OF THE CAMBRIAN DEPOSITS ON THE EASTERN SLOPE OF KUZNETSKIY ALATAU]: Geol. i geofiz., Sib. otd. Akad. Nauk SSSR, No. 2.
 22. Nakaryakov, V. D., Stroyeniye domezozoy-skogo fundamenta Priyeniseyskoy polosy Zapadno-Sibirskoy platformy. [STRUCTURE OF THE PRE-MESOZOIC BASEMENT OF THE YENISEY BELT OF THE WEST-SIBERIAN PLATFORM]: Geol. i geofiz., Sib. otd. Akad. Nauk SSSR, No. 3.
 23. Nedovizin, A. A., K stratigrafii akzhal'skoy svity Chu-Iliyskikh gor. [STRATIGRAPHY OF THE AKZHAL'SKAYA SUITE IN THE CHU-ILIYSKIYE MOUNTAINS]: Izv. Akad. Nauk KazakhSSR, ser. geol., vyp. 2.
 24. Neyman, V. B., and I. V. Kirillov, Gipoteza rasshiryayushcheyas Zemli v yeye geologo-geofizicheskoy sushchnosti. [THE EXPANDING EARTH HYPOTHESIS AND ITS GEOLOGIC-GEOPHYSICAL ESSENCE]: Byul. MOIP, otd. geol., t. 36, vyp. 2.
 25. Nikolayev, V. A., and Mu-dun Chzhan, Kartometricheskoye izucheniye landshaftov. [CARTOMETRIC STUDY OF LANDFORMS]: Vestn. MGU, ser. V, geogr., No. 2.
 26. Ovechkin, N. K., Novyye dannyye o vozraste

- chechanskoy svity. [NEW DATA ON THE AGE OF THE CHECHANSKAYA SUITE]: Inform. sb. VSEGEI, No. 42.
77. Padalka, G.L., Nekotoryye voprosy tektoniki, magmatizma i metallogenii Zabaykal'ya. [A FEW PROBLEMS OF THE TECTONICS, MAGMATISM, AND METALLOGENY IN TRANS-BAYKALIA]: Inform. sb. VSEGEI, No. 41, 1960.
78. Pamyati, A.N., Mazarovicha. [IN MEMORIAM TO A.N. MAZAROVICH]: Byul. MOIP, otd. geol., t. 36, vyp. 2.
79. Petrashkevich, M.I., Ya. A. Voloshchak, A.I. Guridov, and N.N. Demchuk, K voprosu geologicheskogo stroyeniya Zakarpat'ya v svete novykh dannykh bureniya. [THE PROBLEM OF THE GEOLOGIC STRUCTURE OF THE TRANSCARPATHIAN REGIONS IN THE LIGHT OF NEW BORING DATA]: Dopovidi, Akad. Nauk UkrSSR, No. 4.
80. Pogonya-Stefanovich, Yu. F., Novyye dannyye o stratigrafii pozdnetretichnykh otlozheniy vostochnoy okrainy Zapadno-Sibirskoy nizmennosti (mezhdurech'ye Sym-Dubches). [NEW DATA ON THE STRATIGRAPHY OF LATE-TRIASSIC DEPOSITS IN THE EASTERN FRINGE OF THE WEST SIBERIAN LOWLAND (SYM-DUCHBES INTERFLUVE)]: Geol. i geofiz., Sib. otd. Akad. Nauk SSSR, No. 3.
81. Podgornaya, N.S., and A.G. Gur'yev, O nekotorykh osobennostyakh tolshchi konglomeratobrekhiy, razvitykh po r. Vorogovke na severe Yeniseyskogo kryazha. [CERTAIN PECULIARITIES OF THE CONGLOMERATE-BRECCIA SERIES ALONG THE VOROGOVKA RIVER IN THE NORTHERN PART OF THE YENISEY RIDGE]: Inform. sb. VSEGEI, No. 40, 1960.
82. Polak, L.S., Ye. M. Filippov, G.A. Kuznetsov, and V. Ya. Zhavoronkov, Issledovaniye spektra rasseyannogo gamma-izlucheniya primenitel'no k resheniyu nekotorykh zadach geofiziki. [INVESTIGATION OF SCATTERED GAMMA-RADIATION SPECTRA IN THE SOLUTION OF CERTAIN GEOPHYSICAL PROBLEMS]: Geol. i geofiz., Sib. otd. Akad. Nauk SSSR, No. 3.
83. Postnikov, D.V., Litologiya i usloviya obrazovaniya terrigennoy tolshchi devona Zapadnoy Bashkirii. [THE LITHOLOGY AND CONDITIONS OF FORMATION OF THE CONTINENTAL DEVONIAN FORMATION IN WESTERN BASHKIRIA]: Geol. nefi i gaza, No. 4.
84. Ptushkin, E.I., K.V. Tiunov, and G. Khudaynazarov, Tektonika Bol'shogo Balkhana. [THE TECTONICS OF THE GREAT BALKHAN]: Izv. Akad. Nauk TurkmSSR, ser. fiz. -tekh., khim. i geol. nauk, No. 1.
85. P'yankov, V.M., and G.M. Davydova, K voprosu o tak nazyvayemykh "strukturnakh bez korney" stalingradskogo tipa. [THE PROBLEM OF THE SO-CALLED "ROOTLESS STRUCTURES" OF THE STALINGRAD TYPE]: Geol. nefi i gaza, No. 4.
86. Rabkin, M.I., Novyye dannyye po geologii Antarktidy. [NEW DATA ON THE GEOLOGY OF ANTARCTICA]: Inform. byul. Sov. Antarkt. eksp., No. 23, 1960.
87. Rzhonsnitskaya, M.A., K biostratigraficheskomu raschleneniyu devona Severo-Vostoka SSSR. [THE BIOSTRATIGRAPHIC COLUMN OF THE DEVONIAN IN NORTH-EASTERN USSR]: Inform. sb. VSEGEI, No. 42.
88. Rudnik, V.A., and L.M. Alekseyev, Osobennosti proyavleniya dislokatsionnoy metamorfizma v predelakh khr. Dzhagdy-Tukuringra na Dal'nem Vostoke. [PECULIAR MANIFESTATIONS OF DISLOCATION METAMORPHISM WITHIN THE LIMITS OF THE DZHAGDY-TUKURINGRA RANGE IN THE RUSSIAN FAR EAST]: Inform. sb. VSEGEI, No. 41, 1960.
89. Ryzhkov, O.A., K metodike paleotektonicheskogo analiza. [METHOD OF PALEOTECTONIC ANALYSIS]: Uzb. geol. zh., No. 2.
90. Ryabokon', S.M., and V.A. Khomenko, Ob istochnikakh pitaniya devonskikh otlozheniy na yuge Dneprovsko-Donetskoy vpadiny. [SEDIMENTATION SOURCES OF THE DEVONIAN DEPOSITS IN THE SOUTH DNEPR-DONETZ DEPRESSION]: Geol. zh. Akad. Nauk UkrSSR, t. 21, vip. 2.
91. Savel'yev, B.A., Glyatsiologicheskiye issledovaniya Chetvertichnoy antarkticheskoy ekspeditsii v 1959 g. [GLACIOLOGICAL INVESTIGATIONS OF THE QUATERNARY ON THE ANTARTIC EXPEDITION IN 1959]: Byul. MOIP, otd. geol., t. 36, vyp. 2.
92. Saks, V.N., Nekotoryye voprosy stratigrafii i fatsial'noy kharakteristiki mezozoyskikh otlozheniy Zapadno-Sibirskoy nizmennosti. [CERTAIN PROBLEMS OF THE STRATIGRAPHIC AND FACIES CHARACTERISTICS OF THE MESOZOIC

- DEPOSITS IN THE WEST SIBERIAN LOWLANDS]: Geol. i geofiz., Sib. otd. Akad. Nauk SSSR, No. 3.
3. Svireenko, I. P., Granulometricheskiy i mineralogicheskiy sostav donnykh otlozheniy morya Deyvisa. [GRANULOMETRIC AND MINERALOGIC COMPOSITIONS OF THE BOTTOMSET DEPOSITS OF THE DAVIS SEA]: Inform. byul. Sov. Antarkt. eksp., No. 22, 1960.
 4. Seidov, A. G., Usloviya obrazovaniya maykopskikh otlozheniy Azerbaydzhana. [FORMATIVE CONDITIONS OF THE MAYKOP DEPOSITS IN AZERBAIDZHAN]: Izv. Akad. Nauk AzerbSSR, ser. geol. - geogr. nauk i nefti, No. 1.
 5. Semenenko, N. P., Nekotoryye voprosy metodologii v geologii. [CERTAIN PROBLEMS OF METHODOLOGY IN GEOLOGY]: Geol. zh. Akad. Nauk UkrSSR, t. 21, vip. 2.
 6. Serebryakov, V. A., Okhandzhinskiy granitoidnyy massiv i sopostavleniye yego s massivom Ch'yorgo (bassey n verkhov'yev r. Kolymy). [THE OKHANDZHINSKIY GRANITOID MASSIF AND ITS COMPARISON WITH THE CH'ORGO MASSIF (BASIN OF THE UPPER KOLYMA)]: Izv. vyssy. uchebn. zaved., ser. geol. i razvedka, No. 1.
 7. Skotarenko, V. V., Sledy oledeneniya yugo-vostochnoy chasti Aldanskogo nagor'ya i khrebt Dzhugdzhur. [GLACIATION IN THE SOUTHEASTERN PART OF THE ALDAN UPLAND VESTIGES OF THE DZHUDZHUR RANGE]: Byul. MOIP, otd. geol., t. 36, vyp. 2.
 8. Spiridonov, A. I., Problema poverkhnostey vyvyrnivaniya v SSSR. [THE PROBLEM OF PENEPLAINS IN THE USSR]: Byul. MOIP, otd. geol., t. 36, vyp. 2.
 9. Stovas, M. V., Znachenije neravnomernosti vrashcheniya Zemli v obrazovanii planetarnykh glubinnykh razlomov v zemnoy kore. [THE SIGNIFICANCE OF THE EARTH'S ROTATION IN THE FORMATION OF PLANETARY ABYSSAL FRACTURES IN THE EARTH'S CRUST]: Geol. zh. Akad. Nauk UkrSSR, t. 21, vip. 2.
 10. Suyetova, I. A., Karta i ploshchadi drevnego oledeneniya territorii SSSR. [THE MAP AND AREA OF THE ANCIENT GLACIATION IN THE USSR]: Vestn. MGU, ser. V, geogr., No. 2.
 11. Tvaltvadze, G. K., Geologicheskoye stroyeniye Ardaganskogo rayona po seysmicheskim dannym. [THE GEOLOGIC STRUCTURE OF THE ARDAGAN REGION ACCORDING TO SEISMIC DATA]: Soobshch. Akad. Nauk GruzSSR, t. 26, No. 3.
 12. Teplitskiy, V. A., Stratifikatsiya seysmotricheskikh gorizontov v rayonakh srednego techeniya Amu-Dar'yi. [STRATIFICATION OF THE SEISMO-METRIC HORIZONS IN THE REGIONS OF THE AMU-DAR'YA RIVER'S MIDDLE COURSE]: Novosti neft. i gazovoy tekhniki, geologiya, No. 3.
 13. Timofeyev, D. A., Poverkhnosti vyvyrnivaniya Aldano-Olekminskogo mezhdurech'ya. [THE PENEPLAIN OF THE ALDAN-OLEKMINSK INTERFLUVE]: Byul. MOIP, otd. geol., t. 36, vyp. 2.
 14. Tiunov, K. V., K voprosu o prisutstvii na Bol'shom Balkhane turonskogo yarusa. [THE PROBLEM OF THE EXISTENCE OF THE TURONIAN STAGE IN THE GREAT BALKHAN]: Izv. Akad. Nauk TurkmSSR, ser. fiz. - tekhn., khim. i geol. nauk, No. 1.
 15. Tiunov, K. V., Novyye dannyye o vozraste i usloviyakh zaleganiya magmaticheskikh porod na Bol'shom Balkhane. [NEW DATA ON THE AGE AND THE MODE OF OCCURRENCE OF MAGMATIC ROCKS IN THE GREAT BALKHAN]: Novosti neft. i gazovoy tekhniki, geologiya, No. 3.
 16. Tiunov, K. V., Novyye dannyye o geologicheskimi stroyenii Severobalkhanskogo predgornogo progiba. [NEW DATA ON THE GEOLOGIC STRUCTURE OF THE NORTH BALKHAN FOREDEEP]: Izv. Akad. Nauk TurkmSSR, ser. fiz. - tekhn., khim. i geol. nauk, No. 1.
 17. Tiunov, K. V., and M. A. Tkachuk, Novyye dannyye o paleogenovykh otlozheniyakh zapadnoy chasti Bol'shogo Balkhana. [NEW DATA ON THE PALEOGENE DEPOSITS IN THE WESTERN PART OF THE GREAT BALKHAN]: Izv. Akad. Nauk TurkmSSR, ser. fiz. - tekhn. khim. i geol. nauk, No. 1.
 18. Trusova, I. F., Yavleniya polimetamorfizma v kristallicheskikh slantsakh dokembriya Tsentral'nogo Kazakhstana. [MANIFESTATIONS OF POLYMETAMORPHISM IN THE PRECAMBRIAN CRYSTALLINE SCHISTS OF CENTRAL KAZAKHSTAN]: Izv. vyssh. uchebn. zaved., ser. geol. i razvedka, No. 1.
 19. Usmanov, F. A., Dayki Ikhnachskogo rayona i ikh otnosheniye k intruzivnym i skarnovorudnym obrazovaniyam. [DIKES

IN THE IKHNACHSKIY REGION AND THEIR RELATION TO THE METALLIFEROUS INTRUSIVE AND SKARN FORMATIONS]: Uzb. geol. zh., No. 2.

[STUDY OF THE PHENOLOGY OF RELIEF-FORMING PROCESSES]: Byul. MOIP, otd. geol., t. 36, vyp. 2.

110. Ushko, K.A., and B.M. Ulizlo, O metodike vydeleniya razmyvov v pliotsenovykh otlozheniyakh Pribalkhanskoy zony podnyatiy. [METHODS TO IDENTIFY THE WASHOUTS IN PLIOCENE DEPOSITS OF THE BALKHAN UPLIFT ZONE]: Novosti nef. i gazovoy tekhniki, geologiya, No. 3.
111. Frolov, A.I., and Ye.D. Koryakin, Issledovaniye podlednogo rel'yefa v rayone stantsii Lazarev gravimetricheskim metodom. [INVESTIGATION OF SUBGLACIAL RELIEF BY THE GRAVIMETRIC METHOD IN THE AREA NEAR LAZAREV STATION]: Inform. byul. Sov. Antarkt. eksp., No. 23, 1960.
112. Khlobustov, A.A., Ob osnovnykh zakonmernostyakh razvitiya Zemli. [THE PRINCIPAL LAWS OF THE EARTH'S DEVELOPMENT]: Uzb. geol. zh., No. 2.
113. Khudaynazarov, G., and K.V. Tiunov, Nekotoryye rezul'taty izucheniya yurskoy argillitovoy tolshchi Bol'shogo Balkhana po dannym bureniya. [CERTAIN INVESTIGATION RESULTS OBTAINED FROM BORE HOLE DATA IN THE JURASSIC ARGILLITE FORMATIONS OF THE GREAT BALKHAN]: Izv. TurkmSSR, ser. fiz. -tekhn., khim. i geol. nauk, No. 1.
114. Chebanenko, I.I., O rvoobraznykh progibakh platform. [DITCH-LIKE DOWNWARPS IN PLATFORMS]: Dopovidi, Akad. Nauk UkrSSR, No. 4.
115. Chekunov, A.V., Geotektonicheskiye karty Azovo-Kubanskoy vpadiny v miotsene i antropogene. [GEOTECTONIC MAPS OF THE AZOV-KUBAN' DEPRESSION IN THE MIOCENE AND POST-QUARTERNARY]: Geol. zh. Akad. Nauk UkrSSR, t. 21, vyp. 2.
116. Chernyak, N.Yu., A.T. Bogayets, G.M. Voloshina, and F.T. Khadikin, K stratigrafii melovykh i paleogenovykh otlozheniy severnogo sklona Prichernomorskoy vpadiny. [STRATIGRAPHY OF THE CRETACEOUS AND PALEOGENE DEPOSITS ON THE NORTHERN SLOPE OF THE BLACK SEA DEPRESSION]: Geol. zh. Akad. Nauk UkrSSR, t. 21, vip. 2.
117. Chichagov, V.P., K izucheniyu fenologii rel'yefoobrazuyushchikh protsessov. [STUDY OF THE PHENOLOGY OF RELIEF-FORMING PROCESSES]: Byul. MOIP, otd. geol., t. 36, vyp. 2.
118. Shaposhnikov, K.K., and K.I. Shaposhnikov, Nekotoryye voprosy tektonicheskogo rayonirovaniya Severo-Vostochnoy Yakutii po geofizicheskim dannym. [CERTAIN PROBLEMS OF TECTONIC ZONING IN THE NORTHEAST YAKUTIYA ON THE BASIS OF GEOPHYSICAL DATA]: Geol. i geofiz. Sib. otd. Akad. Nauk SSSR, No. 2.
119. Sheynmann, Yu.M., Znachenie okeanov atlanticheskogo tipa dlya razvitiya struktury Zemli. [THE IMPORTANCE OF ATLANTIC-TYPE OCEANS IN THE DEVELOPMENT OF THE EARTH'S STRUCTURE]: Byul. MOIP, otd. geol., t. 36, vyp. 2.
120. Shmulevich, A.D., O psevdomorfozakh sulfidov osadochnogo proiskhozhdeniya. [THE PSEUDOMORPHOSES OF SULFIDES OF SEDIMENTARY ORIGIN]: Uzb. geol. zh., No. 2.
121. Shuvalov, Yu.M., Bogdanovskiy magmaticheskii kompleks v Stepnyaksko-Stalinskom rayone Severnogo Kazakhstana. [THE BOGDANOVSK MAGMATIC COMPLEX IN THE STEPNYAKSKO-STALINSK REGION OF NORTHERN KAZAKHSTAN]: Inform. sb. VSEGEI, No. 41, 1960.
122. Epshteyn, G.M., Po povodu stat'i L.V. Zorina "O paleogeograficheskom znachenii merzlykh sloyev". [WITH REFERENCE TO L.V. ZORIN'S ARTICLE: "THE PALEOGEOGRAPHIC SIGNIFICANCE OF FROZEN LAYERS"] Izv. vyssh. uchebn. zaved., ser. geol. i razvedka, No. 1.
123. Esenov, M.E., O litologii pliotsenovykh otlozheniy Pribalkhanskogo rayona. [THE LITHOLOGY OF PLIOCENE DEPOSITS IN THE BALKHAN REGION]: Izv. Akad. Nauk TurkmSSR, ser. fiz. -tekhn., khim. i geol. nauk, No. 1.
124. Yurevich, A.L., Vulkanicheskiye peply i nizhneapsheroniskikh otlozheniy poluostrova Cheleken. [VOLCANIC ASH FROM THE LOWER-APSHERONIAN DEPOSITS ON THE CHELEKEN PENINSULA]: Izv. Akad. Nauk TurkmSSR, ser. fiz. -tekhn. i geol. nauk, No. 1.
125. Aliyev, Kh.Sh., Novyye vidy radiolyariy nizhnemelovykh otlozheniya Severo-Vostochnogo Azerbaydzhana. [NEW TYPES OF RADIOLARIA IN THE LOWER CRETACEOUS DEPOSITS OF NORTH-

PALEONTOLOGY

- EASTERN AZERBAIDZHAN]: *Izv. Akad. Nauk AzerbSSR, ser. geol. -geogr. nauk i nefti*, No. 1.
6. Ali-zade, Ak. A., *Novyye vidy i filogeneticheskaya skhema razvitiya al'bskikh belemnitov gruppy Neohibolites minimus* List. [NEW TYPES AND THE PHYLOGENETIC DEVELOPMENT PATTERN OF ALBIAN BELEMNITES OF THE NEOHIBOLITES MINIMUS LIST. GROUP]: *Izv. Akad. Nauk AzerbSSR, ser. geol. -geogr. nauk i nefti*, No. 1.
 7. Amurskiy, G. I., *O nakhodke zuba Equus caballus v otlozheniyakh karakumsovsity. [DISCOVERY OF A EQUUS CABALLUS TOOTH IN THE DEPOSITS OF THE KARAKUM SERIES]*: *Izv. Akad. Nauk TurkmSSR, ser. fiz. -tekhn., khim. i geol. nauk*, No. 1.
 8. Astakhova, T. V., *Tret'ye zasedaniye Sibirskogo otdeleniya Vsesoyuznogo paleontologicheskogo obshchestva. [THE THIRD SESSION OF THE SIBERIAN BRANCH OF THE ALL UNION PALEONTOLOGICAL SOCIETY]*: *Geol. i geofiz., Sib. otd. Akad. Nauk SSSR*, No. 2.
 9. Butusova, I. P., *Nekotoryye rannekamenougol'nyye gastropody Kuznetskogo basseyna. [A FEW EARLY CARBONIFEROUS GASTROPODS IN THE KUZNETSK BASIN]*: *Inform. sb. VSEGEI*, No. 42.
 10. Vasilyuk, N. P., *Kamenougol'nyye korally zapadnogo prodolzheniya Donetskogo basseyna. [CARBONIFEROUS CORALS IN THE WESTERN EXTENSION OF THE DONETZ BASIN]*: *Geol. zh. Akad. Nauk UkrSSR*, t. 21, vyp. 2.
 11. Gekker, R. F., *O prizhiznennoy svyazi organizmov geologicheskogo proshlogo. [LIFETIME RELATIONS OF ORGANISMS IN THE GEOLOGIC PAST]*: *Byul. MOIP, otd. geol.*, t. 36, vyp. 2.
 12. Gofman, Ye. A., *Nekotoryye vidy yurskikh foraminifer Yugo-Vostochnogo Kryma. [CERTAIN GENERA OF JURASSIC FORAMINIFERA IN SOUTHEASTERN CRIMEA]*: *Geol. zh. Akad. Nauk UkrSSR*, t. 21, vyp. 2.
 13. Drushits, V. V., *Aptikhi (obzor literaturey). [THE APTYCHI (REVIEW OF PUBLISHED REPORTS)]*: *Byul. MOIP, otd. geol.*, t. 36, vyp. 2.
 14. Ivanova, Ye. A., *Ekologiya i sistematika na primere brakhiopod. [ECOLOGY AND TAXONOMY AS EXEMPLIFIED BY BRACHIOPODS]*: *Byul. MOIP, otd. geol.*, t. 36, vyp. 2.
 135. Kotsyubinskiy, S. P., *Zona Inoceramus labiatus v mele Volyno-Podol'skoy plity. [ZONE OF INOCERAMUS LABIATUS IN THE CRETACEOUS OF THE VOLYNIA-PODOLSK PLATFORM]*: *Geol. zh. Akad. Nauk UkrSSR*, t. 21, vyp. 2.
 136. Likharev, B. K., *K poznaniyu roda Camarophoria King i nekotorykh drugikh sblizhayemykh s nim brakhiopod. [CONTRIBUTION TO THE KNOWLEDGE OF CAMAROPHORIA KING AND CERTAIN OTHER COGNATE BRACHIOPODS]*: *Inform. sb. VSEGEI*, No. 42.
 137. Lomayeva, Ye. T., *Spory i pyl'tsa doverkhnemelovykh osadochnykh otlozheniy Zhitomirskoy oblasti. [SPORES AND POLLEN FROM THE PRE-UPPER CRETACEOUS SEDIMENTARY DEPOSITS IN THE ZHYTOMIR REGION]*: *Geol. zh. Akad. Nauk UkrSSR*, t. 21, vyp. 2.
 138. Lyal'ovich, S. S., and L. P. Markova, *Mikrofauna miotsenovykh otlozheniy Zapadnogo Kopet-Daga. [THE MICROFAUNA OF THE MIOCENE DEPOSITS IN WESTERN KOPET-DAG]*: *Izv. Akad. Nauk TurkmSSR, ser. fiz. -tekhn., khim. i geol. nauk*, No. 1.
 139. Martynova, O. M., *O paleontologicheskikh dokladakh na XI Mezhdunarodnom entomologicheskom kongresse v avguste 1960 g. v Vene. [PALEO-ENTOMOLOGICAL REPORTS AT THE 11TH INTERNATIONAL ENTOMOLOGICAL CONGRESS IN AUGUST 1960, VIENNA]*: *Byul. MOIP, otd. geol.*, t. 36, vyp. 2.
 140. Pasternak, S. I., *Opyt primeneniya diagramm pri opredelenii formy melovykh pektinid. [EXPERIMENT IN THE USE OF DIAGRAMS TO DETERMINE THE FORMS OF CRETACEOUS PECTINIDES]*: *Geol. zh., Akad. Nauk UkrSSR*, t. 21, vip. 2.
 141. Sivertseva, I. A., *Sporovo-pyl'tsevyye komplekсы verkhnepaleozoyskikh otlozheniy yuzhnoy chasti Tungusskogo basseyna. [THE SPORE-POLLEN COMPLEXES IN THE UPPER PALEOZOIC DEPOSITS IN THE SOUTHERN PART OF THE TUNGUSS BASIN]*: *Inform. sb. VSEGEI*, No. 42.
 142. Stukalina, G. A., *Stebli krinoidey iz otlozheniy verkhnego silura gor Aksarly (Tsentral'nyy Kazakhstan). [CRINOID STEMS FROM THE UPPER SILURIAN DEPOSITS OF THE ASKARLY MOUNTAINS (CENTRAL KAZAKHSTAN)]*: *Inform. sb. VSEGEI*, No. 42.

143. Chekhovich, V. D., Ob usloviyakh sushchestv sushchestvovaniya korallov v ludlovskom i zhedinskoy moryakh Tyan'-Shanya. [EXISTENCE CONDITIONS OF THE CORALS IN LUDLOW AND GEDINNIAN SEAS OF TIEN'-SHAN']: Inform. sb. VSEGEI, No. 42.
 144. Sheshegova, L. I., Novyye vidy iskopayemykh rasteniy il'inskoy svity Kuzbassa. [NEW TYPES OF FOSSIL PLANTS IN THE IL'YINSKAYA SUITE IN THE KUZBASS]: Geol. i geofiz., Sib. otd. Akad. Nauk SSSR, No. 3.
- PETROGRAPHY, MINERALOGY,
CRYSTALLOGRAPHY, GEOCHEMISTRY
145. Azizbekov, Sh. A., and T. G. Gadzhiyev, Pervichnyye struktury techeniya Megri-Ordubadskogo batolita. [THE PRIMARY FLOW STRUCTURES OF THE MEGRI-ORDUBAD BATHOLITH]: Izv. Akad. Nauk AzerbSSR, ser. geol.-geogr. nauk i nefti, No. 1.
 146. Alyavdin, V. F., and Ye. S. Alyavdina, O shchelochnykh polevykh shpatakh magmaticheskikh porod i gidrotermal'nykh zhil Chagydanskogo mestorozhdeniya (bassey n. Kolymy). [THE ALKALINE FELDSPARS IN MAGMATIC ROCKS AND HYDROTHERMAL VEINS OF THE CHAGYDANSK DEPOSITS (KOLYMA RIVER BASIN)]: Zap. Vses. min. o-va, ch. 90, vyp. 2.
 147. Ananyan, A. A., and V. P. Dobrovol'skiy, Ob elektroprovodnosti merzlykh gornykh porod. [THE ELECTRIC CONDUCTIVITY OF FROZEN ROCKS]: Geol. i geofiz., Sib. otd. Akad. Nauk SSSR, No. 3.
 148. Baskov, Ye. A., Khloridy magniya v dolomitakh nizhnego kembriya Berezovskogo progiba i ikh vozmozhnyy genezis. [MAGNESIUM CHLORIDES IN THE LOWER CAMBRIAN DOLOMITES OF THE BEREZOVSK TROUGH AND THEIR POSSIBLE ORIGIN]: Inform. sb. VSEGEI, No. 40, 1960.
 149. Belov, N. V., O polozenii kvartsa v sisteme prirodnykh i sinteticheskikh kremnekislorodnykh syoedineniy. [THE POSITION OF QUARTZ IN A SYSTEM OF NATURAL AND SYNTHETIC OXYSILICON FORMATIONS]: Zap. Vses. min. o-va, ch. 90, vyp. 2.
 150. Bryzgalin, O. V., Simpozium po eksperimental'nyy issledovaniyam v oblasti glubinnykh protsessov. [SYMPOSIUM ON THE EXPERIMENTAL INVESTIGATIONS IN THE FIELD OF PLUTONIC PROCESSES]: Geokhimiya, No. 4.
 151. Vadilo, P. S., Gabitus kristallov almaza i otrazheniya usloviy ikh obrazovaniya. [DIAMOND CRYSTAL HABIT AS A REFLECTION OF THE CONDITIONS OF FORMATION]: Zap. Vses. min. o-va, ch. 90, vyp. 2.
 152. Vaynshteyn, E. Ye., and V. I. Lebedev, Novyye opredeleniya Li, Na, K, Rb, Cs, Ca i Sr v standartakh W-1 i G-1 metodami fotometrii plameni. [NEW Li, Na, K, Rb, Cs, Ca, and Sr DETERMINATIONS IN TERMS OF W-1 AND G-1 STANDARDS BY THE METHOD OF FLAME PHOTOMETRY]: Geokhimiya, No. 4.
 153. Ginzburg, I. V., O nekotorykh izmeneniyakh v granitakh na kontakte s daykoy diabaz. [CERTAIN ALTERATIONS IN GRANITES AT THE CONTACT WITH A DIABASE DIKE]: Byul. MOIP, otd. geol., t. 36, vyp. 2.
 154. Gridnev, N. I., and K. A. Nabiyeu, O mineral'nom sostave chetvertichnykh nadvodnodel'tovykh otlozheniy Amu-Dar'yi. [THE MINERAL COMPOSITION OF ABOVE-WATER QUATERNARY DELTAIC DEPOSITS OF THE AMU-DARYA]: Doklady, Akad. Nauk UzbSSR, No. 4.
 155. Grudnev, A. P., Svyaz' khimicheskogo i mineral'nogo sostava porod semeystva kvartsevykh dioritov i granodioritov. [THE RELATIONSHIP BETWEEN THE CHEMICAL AND MINERAL COMPOSITIONS OF ROCKS BELONGING TO THE FAMILY OF QUARTZ-DIORITES AND GRANODIORITES]: Byul. MOIP, otd. geol., t. 36, vyp. 2.
 156. Gurevich, V. I., O rasprostraneni broma v khloridnykh vodakh. [THE DISTRIBUTION OF BROMIDE IN CHLORIDE WATERS]: Razv. i okhrana nedr, No. 1.
 157. Dobrokhotova, Ye. S., Ob amfibolakh Vishnevyykh gor. [THE VISHNEVYYE MOUNTAINS AMPHIBOLES]: Byul. MOIP, otd. geol., t. 36, vyp. 2.
 158. Dolivo-Dobrovol'skiy, V. V., O trekhfaznoy krivoy v sistemakh "porodoobrazuyushchiy silikat-voda". [THE THREE PHASE CURVE IN THE "ROCK-FORMING SILICATE - WATER SYSTEMS"]: Zap. Vses. min. o-va, ch. 90, vyp. 2.
 159. Yermolayev, V. A., Morfologiya kristallov tsirkona iz paleogenovykh otlozheniy okrain Tomskogo vala. [THE MORPHOLOGY OF ZIRCON CRYSTALS FROM THE PALEOGENE DEPOSITS OF THE TOMSK EMBANKMENT SPURS]: Zap. Vses. min. o-va, ch. 90, vyp. 2.

3. Zhilinskiy, G. B., O nekotorykh zakonomenostyakh raspredeleniya germaniya v zhelezorudnykh mestorozhdeniyakh. [CERTAIN REGULARITIES OF THE DISTRIBUTION OF GERMANIUM IN IRON ORE DEPOSITS]: *Izv. Akad. Nauk KazakhSSR, ser. geol.*, vyp. 2.
4. Zalutskiy, V. V., O reliktovykh daykakh i psevdodaykakh nekotorykh rayonov Zabaykal'ya. [RELICT DIKES AND PSEUDO-DIKES OF CERTAIN REGIONS IN TRANSBAIKALIA]: *Geol. i geofiz., Sib. otd. Akad. Nauk SSSR*, No. 2.
5. Zlobin, B. I., M. S. Gorshkova, Pb i Zn v shchelochnykh porodakh i nekotoryye' petrologicheskiye problemy. [Pb AND Zn IN ALKALINE ROCKS AND CERTAIN PETROLOGICAL PROBLEMS]: *Geokhimiya*, No. 4.
6. Isamukhamedov, I. M., and A. A. Kustarnikova, O kaledonskikh intruzivakh pravoberezh'ya r. Angren. [CALEDONIAN INTRUSIVES ON THE RIGHT BANKS OF THE ANGREN RIVER]: *Uzb. geol. zh.*, No. 1.
7. Ismatullayev, Kh. K., Nekotoryye rezul'taty geokhimicheskogo izucheniya yurskikh i melovykh neftegazonosnykh otlozheniy kaganskoy gruppy struktur [CERTAIN RESULTS OF GEOCHEMICAL STUDIES OF JURASSIC AND CRETACEOUS OIL- AND GAS-BEARING DEPOSITS IN THE KAGANSKAYA STRUCTURAL GROUP]: *Uzb. geol. zh.*, No. 1.
8. Karlov, N. N., Iskopayemyye treshcheniy vysykhaniya v tretichnom konglomerate. [FOSSILIZED DESICCATION FISSURES IN TERTIARY CONGLOMERATES]: *Geol. zh. Akad. Nauk UkrSSR*, t. 21, vip. 2.
9. Kayupova, M. M., Silikaty margantsa iz mestorozhdeniy Dzhumart i Kamys v Tsentral'nom Kazakhstane. [MANGANESE SILICATES FROM THE DEPOSITS OF DZUMART AND KAMYS IN CENTRAL KAZAKHSTAN]: *Izv. Akad. Nauk Kazakh SSR, ser. geol.*, vyp. 2.
10. Korel', V. G., and V. N. Kuznetsova, Petrologicheskiy ocherk Ol'ginsko-Ampalykskogo intruziya (severnaya chast' Kuznetskogo Alatau). [A PETROLOGICAL SKETCH OF THE OLGINSKIY-AMPALYKSKIY INTRUSIVE (NORTHERN PART OF KUZNETSK ALATAU)]: *Geol. i geofiz., Sib. otd. Akad. Nauk SSSR*, No. 2.
11. Kudryavtsev, V. Ye., O nekotorykh mezozoysskikh intruziyakh Tuvy. [CERTAIN MESOZOIC INTRUSIVES IN TUVA]: *Inform. sb. VSEGEI*, No. 41, 1960.
12. Kukhareno, A. A., V. I. Skrizhinskaya, E. Ye. Vaynshteyn, and M. M. Kakhana, K geokhimii niobiya i tantala v kompleksakh shchelochno-ul'traosnovnykh porod. [THE GEOCHEMISTRY OF NIOBIUM AND TANTALUM IN COMPLEXES OF ALAKALI-ULTRABASIC ROCKS]: *Zap. Vses. min. o-va, ch. 90*, vyp. 2.
13. Lebedev, V. I., Retsenziya na knigu V. V. Shcherbiny "Osobennosti geokhimii skandiya i tipy yego mestorozhdeniy". Seriya "Geologiya mestorozhdeniy redkikh elementov", vyp. 8, M., Gosgeoltekhizdat, 1960. [REVIEW OF V. V. SHCHERBINA'S BOOK: "THE GEOCHEMICAL FEATURES OF SCANDIUM AND ITS TYPES OF DEPOSITS"]]: *Geokhimiya*, No. 4.
14. Vydayushchiysya sovet'skiy uchenyy i pedagog. (K stoletiyu so dnya rozhdeniya akademika Frantsa Yul'yevicha Levinson-Lessinga). [AN OUTSTANDING SOVIET SCIENTIST AND TEACHER (THE CENTENNIAL OF ACADEMICIAN FRANTS YUL'YEVITCH LEVINSON-LESSING)]: *Izv. Akad. Nauk KazakhSSR, ser. geol.*, vyp. 2.
15. Levskiy, L. K., Kosmogennyye izotopy v Yardymlinskom meteorite. [COSMOGENIC ISOTOPES IN THE YARDYMLINSK METEORITE]: *Geokhimiya*, No. 4.
16. Linetskaya, L. V., Konglomeraty polyanitskoy serii Pokutsko-Bukovinskiy Karpat. [CONGLOMERATES OF THE POLYANITSKAYA SERIES IN THE POKUTSKO-BUKOVINA CARPATHIANS]: *Geol. zh. Akad. Nauk UkrSSR*, t. 21, vip. 2.
17. Malygin, A. A., O magnitnykh svoystvakh gornyykh porod Leninogorskogo rayona. [MAGNETIC PROPERTIES OF THE LENINOGORSKIY REGION ROCKS]: *Izv. Akad. Nauk KazakhSSR, ser. geol.*, vyp. 2.
18. Marfunin, A. S., Shchelochnyye polevyiye shpaty-fazovyye vzaimootnosheniya i opticheskiye svoystva. [ALKALINE FELDSPARS—THEIR PHASE INTERRELATIONS AND OPTICAL PROPERTIES]: *Byul. MOIP, otd. geol.*, t. 36, vyp. 2.
19. Medvedeva, I. Ye., K voprosu genezisa eklogitov Makbal'skogo podnyatiya v Severnoy Kirgizii. [PROBLEM OF ORIGIN OF ECLOGITES IN THE MAKBAL'SKIY UPLIFT OF NORTHERN KIRGHIZIA]: *Byul. MOIP, otd. geol.*, t. 36, vyp. 2.

177. Miroshnichenko, L.A., and T.A. Vele-shina, Selen i tellur v polimetallicheskih mestorozhdeniyakh Tsentral'nogo Kazakhstana. [SELENIUM AND TELLURIUM IN THE POLYMETALLIC DEPOSITS OF CENTRAL KAZAKHSTAN]: Vestn. Akad. Nauk KazakhSSR, No. 1.
178. Mukanov, K.M., M.Z. Fursova, and M.K. Yanulova, Plattnerit iz zony okisleniya svintsovo-baritovogo mestorozhdeniya Karagayly. [PLATTNERITE FROM THE OXIDE ZONE OF THE KARAGAYLY LEAD-BARITE DEPOSITS]: Vestn. Akad. Nauk KazakhSSR, No. 1.
179. Myaz', N.I., Nekotoryye osobennosti formirovaniya khrustalenosnykh kvartsevykh zhil Tsentral'nogo Kazakhstana. [SOME FORMATION PECULIARITIES OF THE CRYSTAL QUARTZ VEINS IN CENTRAL KAZAKHSTAN]: Zap. Vses. min. o-va, ch. 90, vyp. 2.
180. Novoselova, L.N., Amfibol gastingsit-dashkesanitovogo ryada Akatuyevskogo mestorozhdeniya (Vostochnoye Zabaykal'ye). [AMPHIBOLE OF THE HASTINGSITE-DASHKENSANITE SERIES IN THE AKATUYEVSKOYE DEPOSITS (EASTERN TRANSBAYKALIA)]: Zap. Vses. min. o-va, ch. 90, vyp. 2.
181. N'yuton, R., and G.F. Round, Diffuziya geliya v osadochnykh porodakh. [HELIUM DIFFUSION IN SEDIMENTARY ROCKS]: Geokhimiya, No. 4.
182. Pis'mennyy, V.A., K voprosu o lozhnykh formakh kristallov. [THE PROBLEM OF SPURIOUS CRYSTAL FORMS]: Zap. Vses. min. o-va, ch. 90, vyp. 2.
183. Plotnikov, L.M., O migratsii veshchestva pri dinamometamorfizme siniyskikh otlozheniy Yeniseyskogo kryazha. [MIGRATION OF MATTER DURING DYNAMIC METAMORPHISM OF THE SINIAN DEPOSITS OF THE YENISEY RIDGE]: Inform. sb. VSEGEI, No. 40, 1960.
184. Povarennykh, A.S., K voprosu o vliyaniye uchastvuyushchikh v svyazi elektronov na svoystva kristallov. [THE PROBLEM OF THE EFFECT OF ELECTRONS NOT PARTICIPATING IN THE CIRCUIT ON THE PROPERTIES OF CRYSTALS]:
185. Povilaytis, M.M. Yavleniya fel'dshpatizatsii v rudnom pole granitnogo massiva Kuu (Tsentral'nyy Kazakhstan). [THE PHENOMENON OF FELDSPATHIZATION IN THE MINERALIZED ZONE OF THE KUU GRANITE MASSIF (CENTRAL KAZAKHSTAN)]: Geol. rudn. mestorozhd., No. 2.
186. Polunina, L.A., Novyye dannyye ob ul'traosnovnykh porodakh Chadobetskogo podnyatiya. [NEW DATA ON THE ULTRABASIC ROCKS OF THE CHADOBETZKO UPLIFT]: Inform. sb. VSEGEI, No. 40, 1960.
187. Pospelov, G.A., P.I. Kaushanskaya, and S.S. Lapin, Ob obrazovanii zhilopodobnykh brekchiyevykh form mineralizatsii vne treshchin. [FORMATION OF VEIN-LIKE AND BRECCIATED FORMS OF MINERALIZATION OUTSIDE OF FISSURES]: Geol. rudn. mestorozhd., No. 2.
188. Rodionov, D.A., K voprosu o logarifmicheskoy normal'nom raspredelenii sodernzhanii elementov v izverzhennykh gornykh porodakh. [THE PROBLEM OF THE LOGARITHMICALLY NORMAL DISTRIBUTION OF ELEMENT CONTENTS IN IGNEOUS ROCKS]: Geokhimiya, No. 4.
189. Rozina, B.B., Uskorennyy metod kachestvennogo opredeleniya germaniya. [ACCELERATED METHOD FOR QUALITATIVE DETERMINATION OF GERMANIUM]: Inform. sb. VSEGEI, No. 41, 1960.
190. Ronov, A.B., E.Ye. Vaynshteyn, and A.M. Tuzova, K. geokhimii hafniya, tsirkoniya i nekotorykh drugikh elementov gidrolizatsii v glinakh. [THE GEOCHEMISTRY OF HAFNIUM, ZIRCONIUM AND CERTAIN OTHER HYDROLYSATE ELEMENTS IN CLAYS]: Geokhimiya, No. 4.
191. Ryabchikov, I.D., and B.A. Solov'yeva, K geokhimii rubidiya i litiya v slyudonostnykh pegmatitakh Severnoy Karelii. [THE GEOCHEMISTRY OF RUBIDIUM AND LITHIUM IN THE MICACEOUS PEGMATITES OF NORTHERN KARELIA]: Geokhimiya, No. 4.
192. Samoylova, A.P., Metod koeffitsiyentov v biogeokhimicheskikh issledovaniyakh. [THE COEFFICIENTS METHOD IN BIOGEOCHEMICAL INVESTIGATIONS]: Razv. i okhrana nedr, No. 1.
193. Sveshnikova, Ye.V., and L.I. Shabynin, O zelenykh klinopiroksenakh iz skarnovykh porod i nekotorykh drugikh metasomaticheskikh obrazovaniy. [THE GREEN CLINOPYROXENES FROM SKARNS AND CERTAIN OTHER METASOMATIC FORMATIONS]: Zap. Vses. min. o-va, ch. 90, vyp. 2.
194. Slepnev, Yu.S., Sootnosheniya talliya s rubidiyem, tseziyem i kaliyem v

- metamorficheskikh porodakh, granitakh i redkometal'nykh pegmatitakh Sayan. [THE RATIO OF THALLIUM TO RUBIDIUM, CESIUM, AND POTASSIUM IN METAMORPHIC ROCKS, GRANITES, AND RARE-METAL PEGMATITES OF THE SAYAN]: *Geokhimiya*, No. 4.
5. Solov'yev, S. P., Tvorcheskiy put' V. A. Nikolayev. [THE CREATIVE LIFE OF V. A. NIKOLAYEV]: *Zap. Vses. min. o-va*, ch. 90, vyp. 2.
 6. Stavrov, O. D., and T. A. Bykova, Osobennosti raspredeleniya ryada redkikh i letuchikh elementov v porodakh i pegmatitakh Korosten'skogo plutona. [THE DISTRIBUTION PATTERN OF A NUMBER OF RARE AND VOLATILE ELEMENTS IN THE ROCKS AND PEGMATITES OF THE KOROSTEN' PLUTON]: *Geokhimiya*, No. 4.
 7. Starobinets, I. S., and Kh. Z. Sabirov, K geokhimii prirodnikh gazov Yuzhnoy Fergany. [THE GEOCHEMISTRY OF THE NATURAL GASES IN SOUTH FERGANA]: *Doklady, Akad. Nauk UzbSSR*, No. 4.
 8. Tikhonenkov, I. P., O graficheskikh strukturakh zameshcheniya v shchelochnykh porodakh. [GRAPHIC REPLACEMENT STRUCTURES IN ALKALINE ROCKS]: *Zap. Vses. min. o-va*, ch. 90, vyp. 2.
 9. Troitskiy, V. I., Protsessy autigen'nogo mineraloobrazovaniya v nizhnemezozoy-skikh formatsiyakh Yuzhnogo Uzbekistana. [PROCESSES OF AUTOGENETIC MINEROGENESIS IN THE LOWER MESOZOIC FORMATIONS OF SOUTHERN UZBEKISTAN]: *Uzb. geol. zh.*, No. 2.
 10. Ushakova, A. M., K radioaktivnosti porod Perzhanskogo intruzivnogo kompleksa. [RADIOACTIVITY OF THE ROCKS IN THE PERZHANSKIY INTRUSIVE COMPLEX]: *Geokhimiya*, No. 4.
 11. Fadeyev, M. I., K voprosu o proiskhozhdenii satkinskikh kristallicheskikh magnezitov. [THE GENETIC PROBLEM OF THE SATKINSK CRYSTALLINE MAGNESITES]: *Zap. Vses. min. o-va*, ch. 90, vyp. 2.
 12. Futergendler, S. I., and V. A. Frank-Kamenetskiy, Orientirovannyye vrostki olivina, granata i khromshpinelida v almazakh. [ORIENTED OLIVINE, GARNET, AND CHROME-SPINELLID IN-GROWTHS IN DIAMONDS]: *Zap. Vses. min. o-va*, ch. 90, vyp. 2.
 203. Khayrutdinov, D. Kh., K voprosu o privnosenose veshchestv pri obrazovanii vtorichnykh kvartsitov. [THE PROBLEM OF THE DELIVERY AND EVACUATION OF MATTER DURING THE FORMATION OF SECONDARY QUARTZITES]: *Vestn. Akad. Nauk KazakhSSR*, No. 4.
 204. Chesnokov, O. F., and V. S. Sukhnevich, Spektral'nyy metod opredeleniya selena v geologicheskikh probakh. [SPECTRAL METHOD FOR DETERMINATION OF SELENIUM IN GEOLOGIC SAMPLES]:
 205. Shchukin, S. I., O povedenii zheleza pri gidrotermal'nom metamorfizme kizlykh effuzivov Yuzhnogo Gissara (Yuzhnyy Tyan'-Shan'). [BEHAVIOR OF IRON DURING HYDROTHERMAL METAMORPHISM OF THE ACID EFFUSIVES IN SOUTHERN GISSAR (SOUTHERN TIEN-SHAN)]: *Inform. sb. VSEGEI*, No. 41, 1960.
 206. Yushchko, S. A., Mineraly svintsia i serebra v rudakh svintsovo-tsinkovogo orudeniya khrebta Karatau. [LEAD AND SILVER MINERALS IN THE ORES FROM THE LEAD-ZINC DEPOSITS IN THE KARATAU RANGE]: *Izv. vyssh. uchebn. zaved., ser. geol. i razvedka*, No. 1.
 207. MINERAL RESOURCES
 207. Albul, S. P., O gidrogeokhimicheskikh poiskakh rudnykh mestorozhdeniy. [HYDROGEOCHEMICAL PROSPECTING FOR ORE DEPOSITS]: *Razv. i okhrana neдр.*, No. 4.
 208. Albul, S. R., O fiziko-matematicheskoy toerii polya rasseyaniya pri gidrogeokhimicheskikh poiskakh rudnykh mestorozhdeniy. [THE PHYSICO-MATHEMATICAL SCATTERING FIELD THEORY IN APPLICATION TO HYDROGEOCHEMICAL PROSPECTING OF ORE DEPOSITS]: *Byul. MOIP, otd. geol.*, t. 36, vyp. 2.
 209. Al'bov, M. N., Retsenziya na knigu A. A. Yakzhina "Poiski i razvedka mestorozhdeniy poleznykh iskopayemykh". [REVIEW OF A. A. YAKZHIN'S BOOK: "PROSPECTING AND EXPLORATION OF ORE DEPOSITS"]:
Razv. i okhrana neдр., No. 1.
 210. Akhmedzhanov, M. A., O stat'ye Ye. L. Abramovich "O genezise polimetallicheskogo orudneniya plastovogo tipa v dolomitakh Kalkanaty". [ON YE. L. ABRAMOVICH'S ARTICLE: "THE GENESIS OF SHEET-TYPE POLYMETALLIC MINERALIZATION IN THE DOLOMITES OF KALKANATY"]:
Uzb. geol. zh., No. 1.

211. Babayev, A.G., O poznanii neftegazonosnosti Zapadnogo Uzbekistana (po povodu vyskazyvaniy N.A. Kudryavtseva i N.P. Tuayeva). [THE PERCEPTION OF THE OIL AND GAS POTENTIALITIES OF WESTERN UZBEKISTAN (ON THE SUBJECT OF STATEMENTS MADE BY N.A. KUDRYAVTSEVA AND N.P. TUAYEV)]: Uzb. geol. zh., No. 1.
212. Babayev, A.G., O sravnitel'noy otsenke perspektiv neftegazonosnosti rayonov Zapadnogo i Yuzhnogo Uzbekistana. [COMPARATIVE APPRAISAL OF THE PROSPECTS FOR OIL AND GAS IN WESTERN AND SOUTHERN UZBEKISTAN]: Novosti nef. i gazovoy tekhniki, geologiya, No. 3.
213. Badalov, S.T., O brusitovom mramore Almalyka. [BRUCITE MARBLE IN ALMALYK]: Uzb. geol. zh., No. 1.
214. Stepan Dmitriyevich Batishchev-Tarasov, Nekrolog. [STEPAN DIMITRIYEVICH BATISHCHEV-TARASOV. OBITUARY]: Vestn. Akad. Nauk KazakhSSR, No. 4.
215. Bezsmertnyy, V.V., Pervaya Vsesoyuznaya konferentsiya po geologii i metallogenii Tikhookeasskogo rudnogo poyasa. [THE FIRST ALL-UNION CONFERENCE ON THE GEOLOGY AND METALLOGENY OF THE PACIFIC ORE BELT]: Izv. vyssh. uchebn. zaved., ser. geol. i razvedka, No. 1.
216. Borisov, O.M., O klassifikatsii rudno-petrograficheskikh provintsiy. [THE CLASSIFICATION OF MINERALIZED PETROGRAPHIC PROVINCES]: Uzb. geol. zh., No. 1.
217. Borodayevskaya, M.B., and N.A. Perizhnyak, Nekotoryye voprosy struktury Gayskogo mestorozhdeniya mednokolchedannykh rud. [CERTAIN STRUCTURAL PROBLEMS OF THE GAYSKY CHALCOPYRITE ORE DEPOSITS]: Geol. rudn. mestorozhd., No. 2.
218. Bykhover, N.A., Osnovnyye bazy mineral'nogo syr'ya kapitalisticheskikh stran. [MAIN CENTERS OF MINERAL RESOURCES IN THE CAPITALISTIC COUNTRIES]: Razv. i okhrana nedr, No. 4.
219. Varentsov, M.I., and K.N. Kravchenko, Sravnitel'naya kharakteristika neftegazonosnykh basseynov vostochnoy chasti Sredney Azii i Zapadnogo Kitaya. [COMPARATIVE CHARACTERISTICS OF THE OIL- AND GAS-BEARING BASINS IN THE EASTERN PART OF CENTRAL ASIA AND WESTERN CHINA]: Novosti nef. i gazovoy tekhniki, geologiya, No. 3.
220. Vartanyan, S.U., Nekotoryye dannyye o poiskovom znachenii oreola rasseyaniya i fiziko-mekhanicheskikh svoystv porod medno-kolchedannykh mestorozhdeniy Severnoy Armenii. [CERTAIN DATA ON THE PROSPECTING VALUE OF DISPERSION AUREOLES AND THE PHYSICO-MECHANICAL PROPERTIES OF ROCKS IN THE CHALCOPYRITE ORE DEPOSITS OF NORTHERN ARMENIA]: Izv. vyssh. uchebn. zaved., ser. geol. i razvedka, No. 1.
221. Vol'vovskiy, I.S., and V.Z. Ryaboy, Primeneniye metodov matematicheskoy statistiki dlya geologicheskoy interpretatsii gravitatsionnogo polya v Ferganskoj vpadine. [APPLICATION OF THE METHODS OF MATHEMATICAL STATISTICS TO THE GEOLOGIC INTERPRETATION OF THE GRAVITY FIELD IN THE FERGANA DEPRESSION]: Novosti nef. i gazovoy tekhniki, geologiya, No. 1.
222. Genkin, A.D., Z.V. Vasil'yeva, and T.A. Yakovlevskaya, Usloviya nakhozhdeniya apatita v medno-nikelevykh sulfidnykh rudakh Noril'skogo mestorozhdeniya. [CONDITIONS FOR THE DISCOVERY OF APATITE IN THE COPPER-NICKEL SULFIDE ORES OF THE NORIL'SK DEPOSITS]: Geol. rudn. mestorozhd., No. 2.
223. Golovin, V.A., A.V. Kantsel', and M.V. Shumilin, Nekotoryye osobennosti strukturnogo kontrolya ekzogennogo uranovogo orudneniya v osadochnykh i tufogenykh porodakh. [CERTAIN STRUCTURAL CONTROL FEATURES OF EXOGENIC URANIUM MINERALIZATION IN SEDIMENTARY AND TURAGENIC ROCKS]: Geol. rudn. mestorozhd., No. 2.
224. Grudnev, A.P., Staticheskaya svyaznost' rudnykh tel. [STATIC COHESION OF OREBODIES]: Izv. vyssh. uchebn. zaved., ser. geol. i razvedka, No. 1.
225. Dushko, Ye.P., Z.G. Muromtseva, and S.S. Tsurul'nikov, Primeneniye kompleksnogo karotazha pri razvedke zheleznykh rud v Krivbasse. [COMPLEX LOGGING IN IRON-ORE PROSPECTING IN KRIVBASS]: Geol. zh. Akad. Nauk UkrSSR, t. 21, vip. 2.
226. Yermakov, V.I., and Yu.A. Romanov, Sopostavleniye rezultatov vozdukhnykh issledovaniy gamma-polya Gazlinskogo gazovogo mestorozhdeniya i Shiikskoy antiklinali. [COMPARISON OF RESULTS OBTAINED IN AERIAL INVESTIGATION OF THE GAMMA-FIELD IN THE GAZLINSKIY GAS FIELDS AND THE

- SHIHKHASKAYA ANTICLINE]: *Novosti neft. i gazovoy tekhniki*, No. 3.
7. Ismailov, M.I., Vollastonitovyye mestorozhdeniya Uzbekistana. [WOLLASTONITE DEPOSITS IN UZBEKISTAN]: *Uzb. geol. zh.*, No. 1.
 8. Kaminskaya, V.I., Geologicheskiye osobennosti Kondomskogo zhelezorudnogo rayona. [GEOLOGIC FEATURES OF THE KONDOMSKIY IRON ORE DISTRICT]: *Geol. i geofiz.*, Sib. otd. Akad. Nauk SSSR, No. 3.
 9. Komarov, P.V., Magnezial'nyye skarny Teyskogo mestorozhdeniya. [MAGNESIAN SKARNS IN THE TEYSKOYE DEPOSITS]: *Geol. rudn. mestorozhd.*, No. 2.
 10. Korenevskiy, S.M., Otsenka kaliyenosnosti galogennykh otlozheniy Prikaspiyskoy vpadiny. [ESTIMATES OF POTASSIUM CONTENTS IN THE HALOGENOUS DEPOSITS IN THE KASPIAN DEPRESSION]: *Razv. i okhrana nedr.*, No. 4.
 11. Koristashevskaya, T.I., and A.A. Popov, Opyt primeneniya sredne- i nizkochtotoy seysmorazvedki v Dzhezkazganskom rayone. [AN EXPERIMENT IN THE APPLICATION OF MEDIUM AND LOW FREQUENCY SEISMIC PROSPECTING IN THE DZHEZ KAZGANSKIY REGION]: *Izv. Akad. Nauk KazakhSSR*, ser. geol., vyp. 2.
 12. Kosals, Ya.A., O gel'vinosnom tipe skarnovykh mestorozhdeniy. [HELVITE-BEARING SKARN-TYPE DEPOSITS]: *Geol. i geofiz.*, Sib. otd. Akad. Nauk SSSR, No. 2.
 13. Kotlyar, V.N., Ob osobennostyakh obrazovaniya nekotorykh blizpoverkhnostnykh poslemagmatischeskikh mestorozhdeniy. [THE FORMATION CHARACTERISTICS OF SOME NEAR-SURFACE POST-MAGMATIC ORE DEPOSITS]: *Izv. vyssh. uchebn. azved.*, ser. geol. i razvedka, No. 1.
 14. Kuz'min, A.M., Verkhnepaleozoyskoye zolotoye orudneniye v okrestnostyakh g. Tomsk. [THE UPPER-PALEOZOIC AURIFEROUS MINERALIZATION IN THE VICINITY OF TOMSK]: *Geol. rudn. mestorozhd.*, No. 2.
 15. Lebzin, Ye.V., O tektonicheskom stroenii i neftegazonosnosti severozapadnoy chasti Bukhara-Khivinskoy vpadiny. [THE TECTONIC STRUCTURE AND OIL- AND GAS-POTENTIALITIES OF THE NORTHWESTERN PART OF THE BUKHARA-KHIVA DEPRESSION]: *Uzb. geol. zh.*, No. 1.
 16. LEVITSKIY, O.D., Nekrolog. [OBITUARY]: *Geol. rudn. mestorozhd.*, No. 2.
 17. Matsokina-Voronich, T.M., and O.M. Borisov, Petrometallogenicheskiye issledovaniya v Laboratorii metallogenii Instituta geologii Akad. Nauk UzSSR v 1960 g. [THE PETROMETALLOGENIC INVESTIGATIONS IN THE METALLOGENY LABORATORY OF THE GEOLOGICAL INSTITUTE OF THE USBEK ACADEMY OF SCIENCES IN 1960]: *Uzb. geol. zh.*, No. 2.
 18. Merkur'yev, N.D., and A.I. Suderkin, Spetsificheskiye osobennosti, poiskov i razvedki mestorozhdeniy p'yezoopticheskogo syr'ya. [THE SPECIFIC FEATURES OF EXPLORATION AND PROSPECTING FOR DEPOSITS OF PIEZO-OPTICAL RAW MATERIALS]: *Razv. i okhrana nedr.*, No. 1.
 19. Mirkhodzhayev, I.M., Okolorudnyy metamorfizm i yego rol' v tipizatsii polimetallicheskikh mestorozhdeniy Kuraminskogo khrebt. [MANIFESTATIONS OF METAMORPHISM AROUND OREBODIES AND ITS ROLE IN PRE-DETERMINING THE PATTERN OF POLY-METALLIC DEPOSITS IN THE KURAMINSKIY MOUNTAIN RANGE]: *Uzb. geol. zh.*, No. 2.
 20. Moralve, V.M., and S.V. Nuzhnov, Perspektivy neftenosnosti pozdnedokembriyskikh otlozheniy v basseynе srednego techeniya r. Maya. [PROSPECTS OF OIL LOCATION IN THE LATE PRECAMBRIAN DEPOSITS IN THE LOWER-COURSE BASIN OF THE MAYA RIVER]: *Izv. vyssh. uchebn. zaved.*, ser. geol. i razvedka, No. 1.
 21. Muromtsev, V.S., Kuznetskiy basseyn—novyy gazoneftenosnyy rayon Zapadnoy Sibiri. [THE KUZNETSK BASIN — A NEW GAS- AND OIL-BEARING REGION IN WESTERN SIBERIA]: *Geol. i geofiz.*, Sib. otd. Akad. Nauk SSSR, No. 3.
 22. Nekrasov, I.Ya., and K.V. Kablokov, Osnovnyye cherty metallogenii khrebt Ulakhan-Sis na Severo-Vostoke Yakutii. [THE BASIC METALLOGENIC FEATURES OF THE ULAKHANSIY RANGE IN NORTHWESTERN YAKUTIYA]: *Geol. rudn. mestorozhd.*, No. 2.
 23. Nikol'skiy, A.P., and A.N. Yefimov, Olovyanaya mineralizatsiya Del'begetyskogo granitnogo massiva na Kalbe.

- [STANNIFEROUS MINERALIZATION IN THE DEL'BEGETEY GRANITE MASSIF ON THE KALBA]: Inform. sb. VSEGEI, No. 41, 1960.
244. Obshinnikov, A.M., and P.A. Udodov, Soveshchaniye po gidrogeokhimitskim metodam poiskov rudnykh mestorozhdeniy. [CONFERENCE ON THE HYDRO-CHEMICAL METHODS OF PROSPECTING FOR ORE DEPOSITS]: Izv. vyssh. uchebn. zaved., ser. geol. i razvedka, No. 1.
 245. Pamyati, O.D., Levitskogo. [IN MEMORIAM OF O.D. LEVITSKIY]: Geol. i geofiz., Sib. otd. Akad. Nauk SSSR, No. 3.
 246. Petrovskaya, N.V., O mineral'nykh assotsiatsiyakh i nekotorykh usloviyakh formirovaniya kolchedannykh mestorozhdeniy Yuzhnogo Urala. [THE MINERAL ASSOCIATIONS AND CERTAIN CONDITIONS OF FORMATION OF CHALCOPYRITIC ORE DEPOSITS IN THE SOUTHERN URALS]: Geol. rudn. mestorozhd., No. 2.
 247. Popov, A.A., Ob izmenenii bokovykh porod na polimetallicheskom mestorozhdenii Uch-Ochak. [ALTERATIONS OF WALL ROCKS IN THE POLYMETALLIC DEPOSITS OF UCH-OCHAK]: Geokhimiya, No. 4.
 248. Popova, Z.D., Tsink v zone okisleniya plastovykh fosforitov mestorozhdeniya Sarysay. [ZINC IN THE OXIDIZED ZONE OF PHOSPHORITIC SHEET DEPOSITS IN SARYSAY]: Vestn. Akad. Nauk KazakhSSR, No. 1.
 249. Rabinovich, B.I., and Yu. Ya. Mikhaylov, Nekotoryye voprosy differentsial'nykh preobrazovaniy krivyykh VEZ. [CERTAIN PROBLEMS OF DIFFERENTIAL TRANSFORMATION OF VEZ CURVES]: Geol. i geofiz., Sib. otd. Akad. Nauk SSSR, No. 3.
 250. Rozanov, Yu.A., Poristost' gornyykh porod i yeye rol' v lokalizatsii endogenogo orudeneniya. [THE POROSITY OF ROCKS AND ITS ROLE IN THE LOCALIZATION OF ENDOGENIC MINERALIZATION]: Geol. rudn. mestorozhd., No. 2.
 251. Ryzhkov, O.A., Yu. N. Zuyev, S. N. Nazarov, and T. T. Tadzhiyev, O raschlenenii podekzogiroykh otlozheniy Fergany v svyazi s perspektivami ikh neftegazonosnosti. [DIFFERENTIATION OF THE SUB-EXOGENOUS DEPOSITS IN FERGANA IN CONNECTION WITH THEIR PROSPECTIVE OIL CAPABILITY]: Novosti neft. i gazovoy tekhniki, geologiya, No. 3.
 252. Samedov, F.I., and L. A. Buryakovskiy, Gazonosnost' produktivnoy tolshchi yuzhnovostochnoy chasti Apsheron'skogo arkhipelaga. [GAS POTENTIALITIES OF THE PRODUCING SERIES IN THE SOUTH-EASTERN PART OF THE APSHERON ARCHIPELAGO]: Geol. nefti i gaza, No. 4.
 253. Svishchev, M. F., Osobennosti litologicheskogo sostava i neftenosnost' ugleonosnogo gorizonta mestorozhdeniy Bol'shekinelskogo vala. [THE PECULIARITIES OF THE LITHOLOGICAL COMPOSITION AND THE OIL CAPABILITY OF THE COAL-BEARING HORIZON IN THE DEPOSITS OF THE BOL'SHEKINEL'SKIY RAMPART]: Geol. nefti i gaza, No. 4.
 254. Semenov, L.S., O svyazi postmagmaticheskogo orudeneniya s intruziyami trapov povyshennoy shchelochnosti. [THE RELATIONSHIP BETWEEN POST-MAGMATIC MINERALIZATION AND HIGHLY-ALKALINE TRAP INTRUSIONS]: Inform. sb. VSEGEI, No. 40, 1960.
 255. Smirnov, V.I., O knige G. N. Shcherbiny "Formirovaniye redkometal'nykh mestorozhdeniy Tsentral'nogo Kazakh'stana". [ON G. N. SHCHERBINA'S BOOK: "THE FORMATION OF RARE-METAL DEPOSITS IN CENTRAL KAZAKHSTAN"]: Geol. rudn. mestorozhd., No. 2.
 256. Smolin, P.P., Zakonomernosti razmeshcheniya magnezial'no-silikatnykh poleznykh iskopayemykh. [THE EMPLACEMENT MECHANISM OF MAGNESIAN-SILICATE MINERALS]: Geol. rudn. mestorozhd., No. 2.
 257. Titov, V.N., and V.M. Babenko, Osnovnyye cherty struktury mestorozhdeniya Severnogo Kantau v Kansayskom rudnom pole. [THE PRINCIPAL STRUCTURAL FEATURES OF THE SEVERNYY KANTAU ORE DEPOSITS IN THE KANSAYSKIY ORE FIELD]: Geol. rudn. mestorozhd., No. 2.
 258. Tkachuk, L.G., and V.I. Koltun, K voprosu o roli stillolitov v formirovaniye mestorozhdeniy sery Pridnestrov'ya. [THE PROBLEM OF THE ROLE OF STYLOLITES IN THE FORMATION OF SULFUR DEPOSITS IN THE DNYESTER RIVER REGION]: Geol. zh. Akad. Nauk UkrSSR, t. 21, vip. 2.
 259. Tuganova, Ye.V., O rudonosnykh intruziyakh basseyna r. Kureyki. [THE ORE-BEARING INTRUSIONS IN THE

- BASIN OF THE KUREYKA RIVER]: Inform. sb. VSEGEI, No. 40, 1960.
0. Chaynikov, V.I., Assotsiatsiya kassiterit-sul'fidnogo i rtutnogo orudneniya s osnovnymi izverzhennymi porodami v Terneyskom rayone (Primor'ye). [THE ASSOCIATION OF CASSITERITE-SULFIDE AND MERCURY MINERALIZATION WITH BASIC IGNEOUS ROCKS IN THE TERNEYSKIY REGION (PRIMOR'YE)]: Geol. rudn. mestorozhd., No. 2.
 1. Shaposhnikov, G.N., Rudonosnyye skarny Vostochnogo Tannu-Ola (Tuva). [THE ORE-BEARING SKARNS IN EASTERN TANNU-OLA (TUVA)]: Inform. sb. VSEGEI, No. 41, 1960.
 2. Shatalov, Ye.T., O rabote Komissii po geologicheskoy karte mira i rabote sektsii po geneticheskoy probleme rud na XXI sessii Mezhdunarodnogo geologicheskogo kongressa. [THE WORK OF THE COMMISSION ON THE GEOLOGIC MAP OF THE WORLD AND ON THE WORK OF THE SECTION ON THE GENETIC PROBLEM OF ORES AT THE 21ST SESSION OF THE INTERNATIONAL GEOLOGICAL CONGRESS]: Byul. MOIP, otd. geol., t. 36, vyp. 2.
 3. Shvarts, L.Ya., Korrelyatsiya razrezov produktivnoy i krasnotsvetnoy tolshch v svyazi s perspektivami neftenosnosti Yugo-Zapadnoy Turkmenii. [CORRELATION OF THE SECTIONS OF THE PRODUCING AND RED-BED SERIES IN CONNECTION WITH THE PROSPECTIVE OIL CAPABILITY OF SOUTHWESTERN TURKMENIA]: Novosti neft. i gazovoy tekhniki, geologiya No. 3.
 64. Shvarts, T.V., Svyaz' formirovaniya i razmeshcheniya neftnykh zalezhey Zapadno-Turkmenskoy nizmennosti s osobennostyami yeye geologicheskogo razvitiya. [THE CONNECTION BETWEEN THE FORMATION AND EMPLACEMENT OF OIL DEPOSITS IN THE WEST-TURKMENIAN LOWLAND AND THE PECULIARITIES OF THEIR GEOLOGIC DEVELOPMENT]: Novosti neft. i gazovoy tekhniki, geologiya, No. 3.
 65. Shirkov, A.S., Metodika geofizicheskikh rabot prirazvedke poleznykh iskopayemykh. [GEOPHYSICAL SURVEY METHODS IN PROSPECTING FOR ORE DEPOSITS]: Razv. i okhrana nedr, No. 4.
 266. Shnyukov, Ye.F., Ob Inguletskom margantsevom mestorozhdenii. [THE INGULETSK MANGANESE ORE DEPOSITS]: Geol. rudn. mestorozhd., No. 2.
 267. Shcherbakov, Yu.G., Mezhdovedstvennoye soveshchaniye po metallogenii zolota v Zapadnoy Sibiri i Krasnoyarskom kraye. [THE INTER-DEPARTMENTAL CONFERENCE ON THE METALLOGENY OF GOLD IN WESTERN SIBERIA AND KRASNOYARSKIY KRAY]: Geol. i geofiz., Sib. otd. Akad. Nauk SSSR, No. 2.
 268. Shcherbakov, Yu.G., Nekotoryye zakonomernosti zolotoorudneniya na Sinyukhinskom mestorozhdenii. [CERTAIN REGULARITIES OF GOLD MINERALIZATION IN THE SINYUKHIN DEPOSITS]: Geol. i geofiz., Sib. otd. Akad. Nauk SSSR, No. 2.
- HYDROGEOLOGY, ENGINEERING,
GEOLOGY
269. Akhmedsafin, U.M., I.B. Dal'yan, and Zh.S. Sydykov, Artezianskiye vody Vostochnogo Priaral'ya i usloviya ikh formirovaniya. [THE ARTESIAN WATERS IN EASTERN ARAL AREA AND THE CONDITIONS OF THEIR FORMATION]: Izv. Akad. Nauk KazakhSSR, ser. geol., vyp. 2.
 270. Bagirov, T.U., K voprosu ob inzhenerno-geologicheskoy rayonirovaniy severo-zapadnoy chasti Apsheron'skogo polustrova. [THE PROBLEM OF ENGINEERING-GEOLOGIC ZONING OF THE NORTHWESTERN PART OF THE APSHERON PENINSULA]: Izv. Akad. Nauk AzerbSSR, ser. geol.-geogr. nauk i nefti, No. 1.
 271. Volkova, M.A., Novyy uglekislyy istochnik. [A NEW SOURCE OF CARBONIC-ACID]: Priroda, No. 4.
 272. Gafurov, V.G., Inzhenerno-geologicheskoye rayonirovaniye Golodnoy stepi stsel'yu orosheniya. [ENGINEERING-GEOLOGIC ZONING OF THE GOLODNAYA STEPPE FOR PURPOSES OF IRRIGATION]: Uzb. geol. zh., No. 2.
 273. Golubkov, V.N., O laboratornom opredelenii prosadochnosti lessovykh gruntov. [LABORATORY DETERMINATION OF THE TENDENCY OF LOESS TO SAG]: Izv. vyssh. uchebn. zaved., ser. geol. i razvedka, No. 1.
 274. Gurevich, M.S., and N.I. Tostikhin, Skhema khimicheskoy klassifikatsii podzemnykh vod. [DIAGRAM OF THE CHEMICAL CLASSIFICATION OF SUBTERRANEAN WATERS]: Izv. vyssh. uchebn. zaved., ser. geol. i razvedka, No. 1.

275. Gyul', A.K., O vodoprochnosti glin apsheronского yarusa na Apsheronском poluostrove. [THE IMPERMEABILITY OF THE APSHERONIAN-STAGE CLAYS ON THE APSHERON PENINSULA]: Izv. Akad. Nauk AzerbSSR, ser. geol.-geogr. nauk i nefi, No. 1.
276. Gyul', A.K., O prirode fiziko-mekhanicheskikh svoystv glin apsheronского yarusa (na primere Apsheronского poluostrova). [THE NATURE OF THE PHYSICAL AND MECHANICAL PROPERTIES OF THE CLAYS OF THE APSHERONIAN STAGE (AS EXEMPLIFIED ON THE APSHERON PENINSULA)]: Doklady, Akad. Nauk AzerbSSR, t. 17, No. 2.
277. Davydov, I. Ya., O roli vodonepronsityemykh linz peschanykh massivov v pitanii gluboko zalegayushchikh presnykh gruntovykh vod. [THE ROLE OF IMPERMEABLE LENSES OF SAND MASSES IN SUPPLYING DEEP-SEATED FRESH GROUND-WATER SOURCES]: Izv. Akad. Nauk TurkmSSR, ser. fiz.-tekhn., i geol. nauk, No. 1.
278. Devdariani, A.S., and S.D. Voronkevich, O mekhanizme obrazovaniya suffozionnykh polostey v lessovykh porodakh. [THE FORMATION MECHANISM OF SUFFOSION CAVITIES IN LOESS ROCKS]: Uzb. geol., zh., No. 1.
279. Kaldarov, M.K., O rodnike v rayone Darpan-Ata. [THE SPRING IN THE REGION OF DARPAN-ATA]: Izv. Akad. Nauk TurkmSSR, ser. fiz.-tekhn., i geol. nauk, No. 1.
280. Kaldarov, M.K., Ob uchastii Amu-Dar'yi v pitanii gruntovykh vod vostochnoy okrainy Zaunguz'ya. [THE PARTICIPATION OF THE AMU-DAR'YA IN THE FEEDING OF GROUND WATER IN THE EASTERN OUTSKIRTS OF AZUNGUZ'-YE]: Izv. Akad. Nauk TurkmSSR, ser. fiz.-tekhn., i geol. nauk, No. 1.
281. Kats, D.M., Zonal'nost' gruntovykh vod v aridnykh oblastyakh Sredney Azii na primere oazisov Bukhary. [ZONALITY OF GROUND WATER IN THE ARID REGIONS OF CENTRAL ASIA AS EXEMPLIFIED BY THE OASES OF BUKHARA]: Byul. MOIP, otd. geol., t. 36, vyp. 2.
282. Kolodiy, V.V., O khimicheskoy kharakteristike plastovykh vod razvedochnoy ploshchadi Farab. [THE CHEMICAL CHARACTERISTICS OF SHEET WATER IN PROSPECTING THE FARAB AREA]: Izv. Akad. Nauk TurkmSSR, ser. fiz.-tekhn., i geol. nauk, No. 1.
283. Mal'tsev, L.M., and A.L. Bubol'ts, Skvazhina-geyzer na Boya-Dage. [THE GEYSER HOLE IN BOYA-DAG]: Izv. Akad. Nauk TurkmSSR, ser. fiz.-tekhn., i geol. nauk, No. 1.
284. Minkin, Ye.L., O sootnosheniyakh p'yezometricheskikh urovney vodonosnykh gorizontov kamennougol'nykh otlozheniy Moskvoyskogo artezianskogo basseyna. [THE RELATIONS BETWEEN THE PIEZOMETRIC LEVELS OF AQUIFER LEVELS IN THE COAL MEASURES OF THE MOSCOW ARTESIAN BASIN]: Razv. i okhrana nedr, No. 4.
285. Mukhin, Yu.V., O dialekticheskom ponimani gidrogeologicheskoy roli glinistykh otlozheniy kak vodoupornykh razdelov. [THE DIALECTICAL CONCEPT OF THE HYDROGEOLOGIC ROLE OF ARGILLACEOUS DEPOSITS AS IMPERMEABLE DIVIDES]: Byul. MOIP, otd. geol., t. 36, vyp. 2.
286. Ogil'vi, N.A., and O.M. Myaskovskiy, O novom metode poiskov linz presnykh vod v Kara-Kumakh. [A NEW METHOD OF PROSPECTING FOR FRESH WATER LENSES IN THE KARA-KUM]: Izv. Akad. Nauk TurkmSSR, ser. fiz.-tekhn., i geol. nauk, No. 1.
287. Ostrovskiy, V.N., O rezul'tatakh issledovaniya svyazi rastitel'nosti i podzemnykh vod v Dzhezkazgan-Ulutauskom rayone Tsentral'nogo Kazakhstana. [RESULTS OF AN INQUIRY INTO THE RELATIONSHIP BETWEEN VEGETATION AND SUBTERRANEAN WATERS IN THE DZHEZKAZGAN-ULUTAU REGION]: Vestn. Akad. Nauk KazakhSSR, No. 4.
288. Plotnikov, N.I., K metodike razvedki napornykh vod artezianskikh basseynov. [METHODS OF INVESTIGATING THE SOURCES OF PRESSURE WATER IN ARTESIAN BASINS]: Razv. i okhrana nedr, No. 1.
289. Sinyagin, G.P., K stoletiyu so dnya rozhdeniya N.F. Pogrebova. [THE CENTENNIAL OF N.F. POGREBOV]: Izv. vyssh. uchebn. zaved., ser. geol. i razvedka, No. 1.
290. Skaballanovich, I.A., Opredeleniye koeffitsiyenta fil'tratsii peskov po granulometricheskomu sostavu pri otsenke obvodnennosti mestorozhdeniy. [DETERMINATION OF THE SAND FILTRATION FACTOR ON THE BASIS OF GRANULOMETRIC COMPOSITION IN ESTIMATING THE WATER RESOURCES OF ORE DEPOSITS]: Geol. zh., Akad. Nauk UkrSSR, t. 21, vip. 2.

1. Subbota, M.I., Lëssovy karst v nizov'-yakh r. Syr-Dar'yi. [LOESSIAL KARST IN THE LOWER REACHES OF THE SYR-DAR'YA RIVER]: Izv. Akad. Nauk KazakhSSR, ser. geol., vyp. 2.
 2. Sultakhodzhayev, A.N., and G.V. Kulikov, O vodonosnosti chetvertichnykh otlozheniy Severnoy Fergany. [THE ACQUIFEROUS CAPABILITY OF THE QUATERNARY DEPOSITS IN NORTHERN FERGANA]: Uzb. geol. zh., No. 2.
 3. Khodzhibayev, N.N., Uravneniya dlya rascheta balansa gruntovykh vod. [EQUATIONS FOR COMPUTING GROUND WATER BALANCES]: Uzb. geol. zh., No. 1.
 4. Shapiro, S.M., Treshchinnyye vody granitoidov Severo-Zapadnogo Pribalkhash'ya. [FISSURE WATER IN THE GRANITOIDS OF NORTHWESTERN BALKHASH AREA]: Izv. Akad. Nauk KazakhSSR, ser. geol., vyp. 2.
 5. Shevchenko, A.I., O vozmozhnosti ispol'zovaniya napornykh vod konusa vynosov r. Angren na orosheniye. [POSSIBILITY OF UTILIZING THE ALLUVIAL FAN HEADWATERS OF THE ANGERN RIVER FOR IRRIGATION PURPOSES]: Uzb. geol. zh. No. 2.
 6. Shor, G.M., K voprosu o poiskovom znachenii mikrokomponentov v podzemnykh vodakh besstochnykh kotlovin na primere Yugo-Zapadnykh Kyzylkumov. [THE PROBLEM OF THE PROSPECTING VALUE OF MICROCOMPONENTS IN THE SUBSURFACE WATERS OF CLOSED DRAINAGE BASINS AS EXEMPLIFIED IN THE SOUTHWESTERN KYZYLKUM]: Inform. sb. VSEGEI, No. 41, 1960.
- B. Articles in "Materials", "Trudy", "Uchenyye zapiski", and "Sborniks"
1. Geografichniy zbirnik Ukrains'kogo geografichnogo tovaristva, [Geographical Symposium of the Ukrainian Geographic Association]: Vip. 4, Kiev, 1961, 216 pages. Partial contents: Tsys', P.N., The principal genetic types of relief in the West Ukraine; Gerenchuk, K.I., The asymmetry of the river-valley slopes in the Russian lowlands; Marinich, A.M., The karst relief forms of the Volynian Poles'ye; Stepanov, V.V., Processes of rock destruction in the Odessa region; Natarov, V.D., The main historical features of river-valley formation in the Krivoy Rog iron-ore basin; Roslyy, I.M., Alluvial deposits in the Ingulets River valley (within the confines of the Black-Sea lowland); Grubrin, Yu.L., Geomorphological structure of the Ikva river valley; L'vova, Ye. V., The geomorphology and neotectonics of the steppe regions of the Crimea; Mulika, A.M., The structure and origin of the basement of the left-bank regions of the Lower Dnieper; Pogrebnyak, P.S., The paleogeography of Olezh'ye; Roslyy, I.M., and Ye. T. Lomayev, The paleogeography of the Quaternary period in the rightbanks of the Lower Dniepr; Tsys', P.N., Certain observations regarding the land forms of Europe, Africa, and South America; Aleksandrovskiy, T.B., The problem of the geomorphological structure of the Kerch Peninsula; Gubanov, I.G., Landslides on the coasts of the Kerch Peninsula; Podgorodetskiy, P.D., Modern tectonic movements on the Tarkhankutskiy Peninsula; Shtogrin, Ye. D., Quaternary glaciation in the Cis-Carpathia and the northern slope of the Eastern Carpathians; Klevtsov, T.A., Physico-geographic zones of the Dniepropetrovsk area.
 2. Materialy po geologii i poleznym iskopayemykh Severo-Vostoka Yevropeyskoy chasti [Materials on the Geology and Ore Minerals in the North-Western European Part]: SSSR, vyp. 1, Moscow, 1961, 112 pages. Contents: Afanas'yev, B.L., G.M. Yaroslavl'tsev, and V.I. Yatsuk, Formation conditions of coal-bearing deposits in foredeeps, as exemplified by the Pechora basin; Knyazev, S.A., and B.I. Tarbayev, New data on the geology of the Chernysheva Ridge; Yenokyan, V.S., The coal measures in the Bechora coal basin; Abramov, V.P., and G.I. Dembskaya, New information on the Mesozoic deposits in the northern part of the Pechora depression; Miklukho-Maklay, A.D., Volcanism manifestation in the area of the Pechora coal basin and on the western slope of the Polar Urals; Belyakov, L.N., and L. T. Belyakova, Certain features of the geologic structure of the junction of the Urals and Pay-Khoy; Gesse, V.N., Tentative results of geologic-survey operations in the basin of the Paga River (Sub-Arctic Urals); Gribov, B.V., The stratigraphy of the Sabli Range; Vollosovich, K.K., The stratigraphy of the Quaternary deposits of the Northern Europe; Kaletskaya, M.S., The neotectonics of the Pechora coal basin and the adjacent areas of the Northern Urals; Dombrovskaya, Kh. R., A few samaropsises of the Vorkuta series in the Pechora basin and their stratigraphic significance.
 3. Pitaniya geologii ta rozvidki nafrovikh i gazovikh rodovish Karpats'koi zoni, [Problems of Geology and Prospecting

of Oil and Gas Deposits in the Carpathian Zone]: Kiev, 1961, 152 pages. (In Ukrainian) Partial contents: Vyalov, O.S., Various problems of the Miocene stratigraphy in Cis-Carpathia; Dolenko, G.N., The principal prospects and the direction of geologic prospecting for oil and gas in the Cis-Carpathian depression; Kopistynskiy, R.S., The problem of the time of formation of the Carpathian oil deposits; Krayushkin, V.O., Structural features of the Bitkivskiy oil- and gas-field formation. Vitrik, S.P., G.N. Dolenko and B.I. Yarosh, The tectonics and oil potentialities of the Dolinskiy deposits; Kitik, V.I., The problem of the industrial oil- and gas-potentialities of the salt structures in Trans-Carpathia; Gofshteyn, I.D., and M.B. Ripun, The significance of the heavy-fraction minerals in the correlation of the oil deposits in the Carpathians; Vyalov, O.S., I.V. Venglinskiy, and V.M. Utrobin, The problem of differentiation and the presence of oil in the section of borehole No. 1 in the Zalush region; Gavrilenko, K.S., The hydrodynamic factors of oil migration; Gofshteyn, I.D., The problem of the structure of the folded basement between the Russian platform and the Carpathians; Gontoviy, I.Z., The methods of geophysical study of the upper part sequence in the interior zone of the Cis-Carpathian depression; Petkevich, G.I., The seismological characteristics of the interior zone of the Cis-Carpathian depression.

4. Trudy Vostochno-Sibirskogo filiala Akad. Nauk SSSR, vyp. 16, ser. geol. [Transactions of the Eastern Siberian Branch of the USSR Academy of Sciences, No. 16, Geologic Series]: Moscow, 1961, 277 pages. Contents: Bogidayeva, M.V., The Ospinsio-Kitoyskiy ultrabasic massif (eastern Sayan); Kukhrinkova, N.V., The basalts of the Tunkinskaya trough system; Gon'shakova, V.I., The traps of the earliest (Pre-Upper Jurassic) and the latest (Post-Lower Jurassic) phases of intrusive volcanism on the Siberian platform; Belov, I.V., The diabasic-granophyre complex of Aban; Feoktistov, G.D., Materials on the petrography of the traps in the region of Badarminsk Narrows Angara River; Vladimirov, B.M., The Padunskiy diabasic sill; Dvorkin-Samarskiy, V.A., The Mamsk granite pegmatites and their mica content; Rafiyenko, N.I., Certain structural features and localization conditions of the mineralization in the Kyzyl-Chadrsk ore field; Korzhinskiy, A.F., Thermo-optical investigations of certain amphiboles in Eastern Siberia.

5. Trudy Instituta geologicheskikh nauk Akad. Nauk Kazakh SSR, [Transactions of the Geological Institute of the Kazakh Academy of Sciences]: t. 4, Almaty, 1961, 142 pages. Contents: Abdrakhmanov, K.A., and V.G. Korinevskiy, Pseudo-leucitic rocks of the Irisu massif (Tallasskiy Alatau); Nurlybayev, A.N., and V.I. Stravrov, Petrography of the intrusive rocks in the Karatas and Kokzabay mining districts of Northwestern Pribalkhash'ye; Kudryashov, A.V., Transformation in the anorthoclases of the intrusive rocks in Central Kazakhstan; Srayylov, T., The types of secondary quartzites in the north eastern part of Central Kazakhstan; Kolmogorov, Yu.A., The distribution and origin of the secondary quartzites in Central Kazakhstan (on the basis of geophysical data); Masgutov, R.V., One case of pegmatite formation; Tabyldiyev, K.T., Metamorphism of phosphorites and the country rocks of the Chulaktau deposits; Al'mukhanbetov, D., Application of a magnetic survey for mapping of closed structures under conditions prevailing in the Ishim areas. Gul'nitskiy, V.L., and Yu.S. Naboko, The problem of the determination of the magnetic susceptibility of intrusive bodies; Bekmukhametov, A.Y., Stages of endogenic mineralization in the Benkalinskoye ore deposits; Nuralin, N.N., Certain localization peculiarities of ore mineralization in the Dzhezkazgan; Tretyakov, V.B., and S.R. Nikonov, New data on the potentialities of the Shubartobinsk zone in the Chinghiz region; Dolgikh, S.A., and T.A. Oshakpayev, The problem of the Structure of the salt dome cores in the Caspian region; Diarov, M., Effect of the lithology of halogen rocks on the formation of cap rock; Laumulin, T.M., Application of the vector-analysis method to the study of the tectonic conditions of stockwork formation; Zhaparkhanov, S., Subsurface water - an important source of water supply in the Shetskiy region; Dzhabasov, M.Kh., Conditions of ground-water accumulation in the Quaternary deposits of the upper part of the Tokrau river basin; Dal'yan, I.B., and Zh. Sydykov, Iodine-bromine water in the Lower Permian deposits of the Aktyubinsk region of the Urals; Ostrovskiy, V.N., and V.P. Oleksenko, Experimental application of the geobotanic method for mapping river terraces in the Dzhezkazgan-Ulatau region of Central Kazakhstan; Verkhovod, B.N., M.O. Kozhbanova, M.P. Dedeshko, and N.V. Vyatchennikova, Spectrochemical determination of certain rare earths in the DFS-3 spectograph; Medoyev, A.G., The ancient Russian names of minerals.

- Trudy Mineralogicheskogo muzeya Akad. Nauk SSSR, [Transactions of the Mineralogical Museum of the USSR Academy of Sciences]: vyp. 11, Moscow, 1961, 232 pages. Contents: Ginsburg, A.I., and M.Ye. Yakovleva, Certain phenomena of spodumene redeposition in pegmatites; Ginsburg, I.V., Hastingsite in the zone of alkali-granitic metasomatosis and isomorphism in monoclinic amphiboles; Ginsburg, I.V., The origin of oriented textures of spodumene and lepidolite-spodumene pegmatites; Godovikov, A.A., Diagnosis of cobalt, nickel, and iron arsenides; Nesterova, Yu.S., The chemical composition of sphalerites; Orlov, Yu.L., A.I. Ginsburg, and N.G. Pinevich, Paragenetic interrelations of beryllium minerals in certain pegmatite lodes; Slepnev, Yu.S., Genetic characteristics and classification problems of lovorrite-rinkolite pegmatites; Yakhontova, L.K., Cobalt's behavior in the oxidized zone of the Dashkesansk iron ore deposits; Yakhontova, L.K., Composition and property problems relative to the minerals of the erythrite-annabergite series; Brodin, B.V., Wurtzite from the Ken-Shanyk deposits; Ginsburg, I.V., The composition of rhombic amphiboles and the isomorphic replacements in them; Kornetova, V.A., and A.I. Ginsburg, Herderite-hydroxyl from the pegmatites of Eastern Transbaykal; Kornetova, V.A., and Z.V. Vasil'yeva, The pink apatite from a pegmatitic lense; Makarochkina, B.A., K.A. Gonibesova, and M.S. Makarochkina, The "new mineral" perrierite; Orlov, Yu.L., and N.N. Mart'yanov, The rare earth vesuvianite from the Yenisey ridge; Sakharova, M.S., and A.I. Tsvetkova, Hydromica from the ore veins of Gornaya Racha; Semenov, Ye.I., and M.Ye. Kazakov, Ancylyte in the alkali pegmatites of the Kola Peninsula; Semenov, Ye.I., New data on nordite; Semenov, Ye.I., A.P. Khomyakov, and A.V. Bykova, Supergene bastnaesite in the residuum of an alkaline massif; Chukhrov, F.V., V.M. Senderova, and M.T. Yanchenko, Lead and copper content in the bismuthines from North-Koundarsk deposits; Yushko, S.A., and T.I. Ifantopulo, Cinnabar in the oxidized lead-zinc ores of Southern Kazakhstan; Yushko, S.A., Sulvanite in the lead-zinc ores of the Karatau Range; Sol'skiy, D.I., A historical sketch on the Mineralogical Museum of the USSR Academy of Sciences (Pre-revolutionary period).
7. Trudy Nauchno-issledovatel'skogo instituta geologii Arktiki M-va geologii i okhrany nedr SSSR, [Transactions of the Scientific-Research Institute of Arctic Geology of the Ministry of Geology and Conservation of Mineral Resources of the USSR]: t. 117, Moscow, 1961, 146 pages. Contents: Rabkin, M.I., and M.G. Ravich, The Precambrian of the Soviet Arctic; Markov, F.G., and B.V. Tkachenko, The Paleozoic of the Soviet Arctic; Saks, V.N., and S.A. Strelkov, The Mesozoic and Cenozoic of the Soviet and Foreign Arctic; Atlasov, I.P., and V.N. Sokolov, The principal development features of the central part of the Soviet Arctic; Kovaleva, G.A., New data on the basaltic conglomerates in the Sinian deposits in Central and Eastern Taymyr; Kaban'kov, V.Ya., The rhythmic peculiarities of the Middle and Upper Cambrian deposits in the northeastern Siberian platform and the conditions of sedimentation of the rhythmic beds; Zhukov, V.V., The rhythmicity of the carbonate deposits of the Middle and Upper Cambrian on the northeastern wall of the Sukhansk trough; Kopylova, T.N., and V.G. Korchagin, Lithologic and stratigraphic differentiation of the Permian deposits in the Pur-Olensk region; Erlikh, E.N., and V.V. Zhukov, The tectonic structure of the Sukhansk trough; Mezhvilk, A.A., The Quaternary deposits of northern Kharaulakh; Gor, Yu.G., The formative conditions of the coal-bearing deposits of the Kulyumba-Kureyka interfluvium; Yermolov, V.V., Ya.I. Pol'kin, and S.A. Strelkov, The peculiarities of the geologic interpretation of aerial photographs in the region of the Far North; Obidin, N.I., Methods of investigation of the thermal regime of rocks and of sampling aquifers in the Noril'sk territory.

